

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 2003

[Full Table of Content](#)

Results of the  
Classical  
and other  
Long-term Experiments  
2003

Rothamsted Research

---

## 03/W/RN/12 - Organic Manuring

### Rothamsted Research

Rothamsted Research (2004) *03/W/RN/12 - Organic Manuring* ; Yields Of The Field Experiments 2003, pp 46 - 48 - DOI: <https://doi.org/10.23637/ERADOC-1-260>

03/W/RN/12

**ORGANIC MANURING**

**Object:** To study, from crop yields and soil analyses, the effects of a range of types of organic matter - Woburn, Stackyard B.

**Sponsor:** P.R. Poulton. A.J.Macdonald

The 39th year, w. rye, grass/clover ley.

For previous years see 'Details' 1973 and 74-02/W/RN/12.

**Design:** 4 blocks of 8 plots.

**Whole plot dimensions:** 8.0 x 29.5 (8.0 x 26.5 on Block III).

**Treatments:** From 1966 to 1971 the experiment had a preliminary period designed to build up organic matter from different sources. An arable rotation was started on two blocks in 1972 and the remaining two blocks in 1973. After a period of testing the residues, a further period of accumulation was started; on two blocks (which included ley sown in 1979) in 1981 and on the other two (which included ley sown in 1980) in 1982. A second test phase began when leys on the first pair of blocks were ploughed for the 1st test crop in 1987 and on the second pair for the 1st test crop in 1988. From 1988 two blocks, and 1989 the other two, to 1994, plots were split into 6 sub-plots to test five levels of nitrogen and nil. From 1995 to 1997 residual effects of that nitrogen were measured. In 1998 to 2000 yields were taken from whole plots only. In 2001 plots were split into half-plots to test two rates of N. For 2003\* the experiment was modified to test further inputs of organic matter. An arable rotation (w. rye, s. barley, w. beans, w. wheat, forage maize) was started on seven plots within each block; the eighth was sown to a grass/clover ley.

Whole plots

1. **TREATMNT** (Not necessarily applied each year):

1966-1971/2	1979/82-1986/7	Since 2003
Fd	Fd	F
Ln	Lc6	F
St	St	St
Gm	Lc8	CC
Pt	Lc8	Co
Fs	Fs	Dg10
Dg	Dg	Dg25
Lc	Lc6	Lc

F: no organic amendment. St: chopped straw at 7.5t/ha. CC: cover crop prior to spring sown crops. Co: compost at 40t/ha. Dg10: FYM at 10t/ha. Dg25: FYM at 25t/ha. Dg: FYM at 50t/ha. Fd: fertilizers equivalent to FYM. Fs: fertilizers equivalent to straw (+P). Lc/Lc6/Lc8: grass/clover leys. Ln: grass ley + N. Gm: green manure. Pt: peat.

\* ACCORDING TO EXPT. DIARY, OM ADDITION DID NOT RESUME UNTIL SEPT 2003 (IF IN THE 2003-4 CROPPING YEAR; see BOOK for 2004)



03/W/RN/12

**GRAIN TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

<b>TREATMNT</b>	
F (Fd)	6.82
F (Ln, Lc6)	7.39
St (St)	6.74
CC (Gm, Lc8)	7.22
Co (Pt, Lc8)	3.92
Dg10 (Fs)	6.87
Dg25 (Dg)	3.91
Mean	6.13

\*\*\* Standard errors of differences of means \*\*\*

<b>TREATMNT</b>
0.211

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
Blocks.Plots	18	0.298	4.9
GRAIN MEAN DM%	89.6		

**STRAW TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

<b>TREATMNT</b>	
F (Fd)	4.27
F (Ln, Lc6)	4.54
St (St)	4.39
CC (Gm, Lc8)	4.52
Co (Pt, Lc8)	2.60
Dg10 (Fs)	4.45
Dg25 (Dg)	2.64
Mean	3.92

STRAW MEAN DM% 92.7

PLOT AREA HARVESTED 0 005290