Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Results of the Classical and Other Long-term Experiments 2006

Results of the
Classical
and other
Long-term Experiments

Rothamsted Resear

Full Table of Content

06/R/EX/4 - Exhaustion Land

Rothamsted Research

Rothamsted Research (2007) 06/R/EX/4 - Exhaustion Land; Results Of The Classical And Other Long-Term Experiments 2006, pp 19 - 21 - DOI: https://doi.org/10.23637/ERADOC-1-263

06/R/EX/4

EXHAUSTION LAND

Object: To study the residual effects of manures applied 1856-1901, and of additional phosphate applied since 1986, on the yield of continuous s. barley up to 1991, w. wheat since - Hoosfield.

The 151st year, w. wheat.

For previous years see 'Details' 1977, 1973 and 74-05/R/EX/4.

Treatments: All combinations of:-

Whole plots (P test)

```
1. OLD RES
                     Residues of manures applied annually 1876-1901:
   0
                     None
   D
                     Farmyard manure at 35 t
   N
                     96 kg N as ammonium salts
   P
                     34 kg P as superphosphate
   NPKNAMG
                     N and P as above plus 137 kg K as sulphate of potash,
                        16 kg Na as sulphate of soda, 11 kg Mg as sulphate
                         of magnesia
2. P
                     Maintenance P (20 kg P) applied annually from 2000 to
                        maintain existing levels of available P in the
                        soil. (P1) (P2) and (P3) are residues of P applied
                        annually 1986-1992:
                     2000-04
                                   1986-92
  0
                     None
                                   None
  P(P1)
                     20 kg P
                                   44 kg P
87 kg P
                     20 kg P
  P(P2)
  P(P3)
                                   131 kg P
                     20 kg P
```

NOTE: P treatments were applied at 61.5 kg P in error in 2000.

plus

Whole plots (K test, previously N test until 1991)

OLD RES	Residues of manures applied annually 1876-1901:
O D N*	None Farmyard manure at 35 t 96 kg N as nitrate of soda
PK	34 kg P as superphosphate, 137 kg K as sulphate of potash
N*PK	N, P and K as above

Whole plots

Nitrogen: 50 kg N as ammonium sulphate (to supply sufficient S) during first two weeks in March, 200 kg N as ammonium nitrate at GS31/mid-April (whichever comes first) and 50 kg N as ammonium nitrate at GS37 (not later than mid-May)

Experimental diary:

```
28-Sep-05 : T : K : K basal/124.5 kg (muriate of potash at 250 kg)*,
                            plots 1, 3, 5, 7 & 9.
All plots
                        : Keiserite (30 kg Mg)
   29-Sep-05 : B :
                          Ploughed 25 cm wide furrows.
   04-Oct-05 : B :
                        : Cultipress.
                        : Cultipressed, combination drilled, Xi19, tr.
   07-Oct-05 : B :
                            Redigo Twin at 350 seeds/m<sup>2</sup>, rolled.
   17-Oct-05 : B :
                       : Ice at 4.0 1 in 200 1.
                      : Hallmark with Zeon Technology at 50 ml in 200 1.
   05-Dec-05 : B :
                        : Ammonium sulphate (21% N) at 238 kg
   13-Mar-06 : B :
   18-Apr-06 : B :
                       : 34.5% N at 580 kg.
                       : tm)Opus at 0.75 \bar{1} in 200 1.
   21-Apr-06 : B :
                       : tm)Bravo 500 at 1.0 l in 200 l.
: tm)Flexity at 0.2 l in 200 l.
             : B :
              : B :
   17-May-06 : B :
                       : 34.5% N at 145 kg.
                       : tm)Bravo 500 at 1.0 1 in 200 1.
   21-May-06 : B :
                       : tm)Opus at 0.75 1 in 200 1. : tm)Vivid at 0.4 1 in 2001.
              : B :
              : B :
              : B :
                       : tm)Starane 2 at 0.75 1 in 200 1.
                       : tm)Ally Max SX at 42 g in 200 l.
              : B :
                       : Combine harvested discards, baled straw.
   06-Aug-06 : B :
                        : Combine harvested, plots for yield, straw swathed.
   23-Aug-06 : B :
                       : Straw sampled and weighed.
              : B :
                       : Combine harvested discards, swathed and baled
   25-Aug-06 : B :
                             straw.
```

* this is wrongly entered for the 2001 - 2004 Yield Books which incorrectly stated that the basal K was 100 kg K.

NOTE: Samples of grain and straw were taken for chemical analysis.

NOTE:

STRAW DRY MATTER

Due to an error while weighing sub samples (fresh and dry) an average dry matter percent was calculated. The value was 87.8% (P TEST) and 87.7% (K TEST).

P TEST

GRAIN TONNES/HECTARE

**** Tables of means ****

P_RES	0	P(P1)	P(P2)	P(P3)	Mean
OLD_RES					
0	1.74	7.88	8.34	8.01	6.49
D	4.72	8.85	8.45	8.36	7.60
N	1.20	7.77	8.92	8.12	6.50
P	4.87	8.56	8.66	8.00	7.52
NPKNAMG	3.75	8.38	7.82	8.80	7.19
Mean	3.26	8.29	8.44	8.26	7.06

GRAIN MEAN DM% 85.1

ESTIMATED STRAW TONNES/HECTARE

**** Tables of means ****

P_RES O P(P1) P(P2) P(P3) Mean

OLD_RES					
0	0.85	5.17	6.00	6.21	4.56
D	2.83	6.19	6.43	6.51	5.49
N	0.77	5.19	6.29	6.10	4.59
P	2.89	6.10	6.19	5.90	5.27
NPKNAMG	2.30	6.17	6.19	6.37	5.26
Mean	1.93	5.77	6.22	6.22	5.03

STRAW MEAN DM% 87.8

PLOT AREA HARVESTED 0.00525 06/R/EX/4

K TEST

GRAIN TONNES/HECTARE

**** Tables of means ****

OLD_RES	
0	6.51
D	6.92
N*	7.18
PK	7.39
N*PK	7.08
Mean	7.02

GRAIN MEAN DM% 84.9

ESTIMATED STRAW TONNES/HECTARE

***** Tables of means *****

OLD_RES

O 4.49

D 4.90

N* 4.93

PK 5.50

N*PK 4.88

Mean 4
STRAW MEAN DM% 87.7

PLOT AREA HARVESTED 0.00525

4.94