Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 2012



Full Table of Content

W/RN/3 Ley Arable

Rothamsted Research

Rothamsted Research (2013) *W/RN/3 Ley Arable ;* Yields Of The Field Experiments 2012, pp 38 - 48 - **DOI: https://doi.org/10.23637/ERADOC-1-222**

LEY/ARABLE

Object: To compare the effects on soil fertility of rotations with or without leys – Woburn, Stackyard D.

Sponsors: A. J. Macdonald

The 75th year, leys, w. beans, w. wheat, w. rye

For previous years see 'Details' 1967 & 1973 and Yield Books for 74-11/W/RN/3.

Design: 5 series of 8 plots, split for treatments other than rotations.

Whole plot dimensions: 8.53 x 40.7

Treatments: All phases of four five-course rotations were originally present:

ROTATION

| LEY | Clover/grass ley: | L, L, L, P, W |
|-----|--------------------|---|
| CLO | All legume ley: | SA, SA,SA, P, W until 1971 then CL, CL, |
| | | CL, P, W. |
| А | Arable with roots: | P, R, C, P, W until 1971 then P, B, B, P, |
| | | W. |
| ΑH | Arable with hay: | P, R, H, P, W until 1971 then P, B, H, P, |
| | | W |

P = potatoes, R = w. rye, C = carrots, W= w. wheat, B = s. barley, H = hay, L = clover/grass ley, SA = sainfoin ley, CL = red clover ley.

Rotations themselves followed different cycles:

On four plots in each block the rotations were repeated. On four plots in each block arable rotations alternated each five years with ley rotations. From 1976 all the rotations were changed on all phases except for the first and second test crops in 1976:

- LN 3 (Previous LEY) LN1, LN2, LN3, W, R
- LC 3 (Previous CLO) LC1, LC2, LC3, W, R
- AF (Previous A) F, F, BE, W, R
- AB (Previous A H) B, B, BE, W, R

From 1988 rotations AF and AB are replaced by AM and ABe respectively. Phased in at the beginning of each treatment crop sequence. R, BE, M, W, R

AM R, BE, M, W, R ABe R, M, BE, W, R

LN1 to LN3 = three year grass ley with N, 1^{st} year to 3^{rd} year,

```
LC= clover/grass ley, no N, BE = beans (s. oats until 1980), F = fallow,
```

M = forage maize

Plots hitherto in alternating rotations were changed to test eight-year
leys and two test crops:LLNLLN1, LLN2, LLN3, LLN4, LLN5, LLN6, LLN7, LLN8, W, RLLCLLC1, LLC2, LLC3, LLC4, LLC5, LLC6, LLC7, LLC8, W, RLLN1 to LLN8 = eight year grass ley with nitrogen, first year to eighth year, similarly for
LLC - clover/grass ley, no nitrogen

The new scheme started by sowing these new leys in spring 1976 on four phases and in spring 1977 on the fifth phase (2nd test crop in 1976).

In 1992 w. rye (R) replaced s. barley (B) as the second test crop. Yields are taken from the leys, arable treatment crops and the test crops.

From 2007 plots previously in the 1st cycle of testing eight-year leys followed by two arable test crops (i.e. those plots which were changed to eight-year ley treatments in 1976 or 1977) changed to a three-year arable rotation followed by two arable test crops. Plots were "phased in" but joined the relevant point in the rotation. From 2008 the second cycle 8-yr grass and grass/clover leys changed to 3-yr grass or grass/clover leys respectively. They were phased in between 2008 and 2012.

LLN/AO (Previously 1st cycle, 8-yr grass ley) R, BE, O, W, R LLC/ABe (Previously 1st cycle, 8-yr grass/clover ley) R, O, BE, W, R LLC/LC3 (Previously 2nd cycle, 8-yr grass ley) Lc 1, Lc 2, Lc 3, W, R LLN/LN3 (Previously 2nd cycle, 8-yr grass/clover ley) Ln 1, Ln 2, Ln 3, W, R

From 2009 W oats (O) replaced forage maize (M) in the AM and ABe rotations on block III and were phased in on blocks V, IV, II and I in subsequent years. The AM treatment was renamed AM/AO.

Treatments to first test crop w. wheat, all combinations of:

Whole plots:

1. ROTATION R

Rotations before wheat:

LLN 8 LN 3 LLC 8 LC 3 LLC/LC3 not yet in phase LLN/LN3 not yet in phase LLN/AO not yet in phase LLC/ABe not yet in phase AM/AO ABe

1/2 plots:

2. NSPLIT(FYM res)

Nsplit (noFYM) Nsingle(FYM) 1/8 plots: Farmyard manure residues, last applied 1960s: Split N v single N dressing to wheat, tested 2001-5

39

| 3. | Ν | Nitrogen fertil (kg N) as 34.5 | izer as split dressings in spring 2012 5% N: |
|----|-----|-----------------------------------|---|
| | 0 | 0 | |
| | 80 | 40 + 40 |) to be applied |
| | 160 | 40 + 120 |) late-February/early-March |
| | 240 | 40+ 200 |) and mid-April |

Treatments to second test crop w.rye, all combinations of:

Whole plots:

| 1. | ROTATION LLN8 LN 3 LLC 8 LC 3 LLC/LC3 not yet in phase LLN/LN3 not yet in phase LLN/AO not yet in phase LLC/ABe not yet in phase AM/AO ABe | |
|----|--|--|
|----|--|--|

1/2 plots:

 NSPLIT(FYM res) Nsplit to wheat (no FYM) Nsingle to wheat (FYM) 1/8 plots:
Farmyard manure residues, last applied 1960s:

3. N Nitrogen fertilizer in spring 2009 (kg N) as 34.5%:

0 50 100

150

Treatments to leys:

FYM RES Farmyard manure residues:

NONE

FYM 38 t on each occasion, last applied 1960s.

NOTE: Corrective K dressings (kg K₂O ha⁻¹) as muriate of potash, applied where necessary to first test crop w. wheat and long-term leys in the wheat block, applied 2011 (actual date not recorded).

| Continuous rotations | No FYM | FYM Res |
|----------------------|------------|------------|
| Before wheat | Half plots | Half plots |
| Abe/Be | 270 | 340 |
| AO/O | 200 | 270 |
| LLn/AO | 0 | 30 |
| Ln/Ln | 90 | 70 |
| None to other plots. | | |

Experimental Diary

| Date | | Application | Rate | Units |
|------------------|------|---|------|---------|
| A 11 | | | | |
| All 18-Oct-11 | а | Plough all plots at 14" except those remaining in grass. | - | |
| 18-Oct-11 | а | Ploughed - thrown east on 14" furrows. Grass plots not ploughed. | - | |
| 01-Nov-11 | а | Cambridge Rolled - block 1 only. | - | |
| 23-Apr-12 | а | Cut Paths | - | |
| 22-May-12 | а | Cut paths | - | |
| 16-Jul-12 | а | Cut paths | - | |
| Grass ley and c | love | r/grass leys (first year leys) | | |
| 21-Oct-11 | f | Applied Nitram - applied to plots 11,12 and 13,14 only - new grass plots. | 50 | kg N/ha |
| 21-Oct-11 | f | Applied Nitram - applied to plots 3,4 and 7,8 only - new grass plots | 25 | kg N/ha |
| 21-Oct-11 | f | Applied Triple Super Phosphate - applied to 1st year leys plots 3,4,7,8,11,12,13,14 | 213 | kg/ha |
| 21-Oct-11 | f | Applied Potassium Sulphate - applied to first year leys plots 3,4,7,8,11,12,13,14 | 140 | kg/ha |
| 01-Nov-11 | S | Drilled Grass plots - Ln1 plots only, plots 11, 12, 13 and 14. | 30 | kg/ha |
| 01-Nov-11 | S | Drilled Grass plots - Lc1 plots only; plots 3, 4, 7 and 8. | 30 | kg/ha |
| 24-Nov-11 | а | Cut and weighed grass for yield (Yield cut 2 in 2011) | - | |
| 30-Nov-11 | а | Cut and removed grass (yield cut 2 in 2011) | - | |
| 01-Dec-11 | а | Topped a second time to tidy up. | - | |
| 17-Jul-12 | а | First cut - cut and weighed grass plots for yield | - | |
| 24-Jul-12 | а | Grass areas topped | - | |
| 25-Jul-12 | а | Turned hay | - | |
| 28-Jul-12 | а | Rowed up hay | - | |
| 28-Jul-12 | а | Baled and Removed | - | |
| 18-Aug-12 | а | Sprayed Samurai - Ln3 & Lc3 plots | 4 | l/ha |
| 05-Nov-12 | а | Second cut - cut and weighed grass plots for yield | - | |
| Grass ley and c | love | r/grass leys (second and third year leys) | | |
| 21-Oct-11 | f | Applied Triple Super Phosphate to 2nd and 3rd year leys plots 23,24,25,26,29,30,31,32 | 213 | kg/ha |
| 21-Oct-11 | f | Applied Potassium Sulphate to 2nd and 3rd year leys plots 23,24,25,26,29,30,31,32,55,59,57,58,56,60,51,62 | 140 | kg/ha |

| 24-Nov-11 | а | Cut and weighed grass for yield (yield cut 2 in 2011) | - | |
|-----------|---|---|---|------|
| 30-Nov-11 | а | Cut and removed grass (yield cut 2 in 2011) | - | |
| 01-Dec-11 | а | Topped a second time to tidy up. | - | |
| 17-Jul-12 | а | First cut - cut and weighed grass plots for yield | - | |
| 24-Jul-12 | а | Grass areas topped | - | |
| 25-Jul-12 | а | Turned hay | - | |
| 28-Jul-12 | а | Rowed up hay | - | |
| 28-Jul-12 | а | Baled and Removed | - | |
| 18-Aug-12 | р | Sprayed Samurai - Ln3, Lc3 plots only | 4 | l/ha |
| 05-Nov-12 | а | Second cut - cut and weighed grass plots for yield | - | |

W Beans

| 21-Oct-11 | f | Applied Triple Super Phosphate in accordance with plan | 127 | kg/ha |
|-----------|---|--|-----|---------------------|
| 26-Oct-11 | S | Drill W Bean - Wizzard | 25 | seed/m ² |
| 28-Nov-11 | р | Sprayed Crawler | 3.5 | kg/ha |
| 18-Aug-12 | p | Sprayed Samurai | 4 | l/ha |
| 03-Sep-12 | а | Combined for yields | - | |
| 11-Sep-12 | а | Baled and Removed | - | |
| | | | | |
| W Wheat | | | | |
| 21-Oct-11 | f | Applied Muriate of Potash to plots 65, 66, 67, 68, 72, 73 and 74, as per plan | - | |
| 21-Oct-11 | f | Applied Triple Super Phosphate in accordance with plan | 127 | kg/ha |
| 22-Oct-11 | а | Cultivate using pigtail cultivator to 13cm to incorporate fertilisers | - | |
| 26-Oct-11 | S | Drill Wheat - Solstice @ 400 seeds/m2 (Solstice substituted the Glasgow, ok'd with AM) | - | |
| 30-Dec-11 | р | Sprayed Lexus class @ 200 lt/ha water volume. | 60 | g/ha |
| 23-Apr-12 | f | Applied 1 st Nitrochalk 27%N according to plan | 148 | kg/ha |
| 3-May-12 | f | Applied remaining Nitrochalk 27%N by hand to | - | |
| | | blocks 5 according to plan. | 148 | ka/ba |
| | | N1 plots N2 plots | 444 | kg/ha kg/ha |
| | | N3 plots | 740 | kg/ha |
| 17-May-12 | р | spray fungicide T1 + herbicide; Thor @20g/ha + Ignite @1I/ha in 200I water | ha | g/ha |
| 24-May-12 | р | spray fungicides T2; Rubric @0.8I + Comet @0.5I + Rover @1I/ha in 200I water RN3 - wheat only | ha | l/ha |
| 03-Sep-12 | а | Combined for yields | - | |
| 11-Sep-12 | а | Baled and Removed | - | |
| W Rye | | | | |
| 21-Oct-11 | а | Applied Chalk to plots 33 - 48. | 174 | |
| 21-Oct-11 | f | Applied Triple Super Phosphate | 127 | kg/ha |
| | | · · · · | | - |

| 26-Oct-11 | S | Drill W.Rye - Humbolt | 400 | seed/m ² |
|-----------|---|---|-----|---------------------|
| 01-Nov-11 | S | Drilled Rye - Humbolt - Block 1 - plots 1, 2, 5, 6, 9, 10, 15 and 16. | 400 | seed/m ² |
| 30-Dec-11 | р | Sprayed Lexus class in 200 l/ha of water | 60 | g/ha |
| 23-May-12 | а | Nitrochalk 27%N applied by hand to block 3 | - | |
| | | N1 plots | 180 | kg/ha |
| | | N2 plots | 364 | kg/ha |
| | | N3 plots | 545 | kg/ha |
| 03-Sep-12 | а | Combined for yields | - | |
| 11-Sep-12 | а | Baled and Removed | - | |
| W Oats | | | | |
| 21-Oct-11 | f | Applied Triple Super Phosphate in accordance with plan | 127 | kg/ha |
| 28-Oct-11 | S | Drilled oat plots with Mascani @400seeds/m2 | 400 | seed/m ² |
| 30-Dec-11 | р | Sprayed Lexus class in 200 l/ha of water | 60 | g/ha |
| 03-Sep-12 | а | Combined for yields | - | |
| 11-Sep-12 | а | Baled and Removed | - | |
| | | | | |

NOTE: Herbage and grain samples taken for chemical analyses

LEYS

1^{st} Cut (17 /6/12) dry matter tonnes/hectare

***** TABLES OF MEANS *****

| FYM_RES LEY | NONE | FYM | Mean |
|----------------|------|------|------|
| | 4 00 | | 4 70 |
| LC1 | 4.89 | 4.67 | 4.78 |
| LC2 | 3.25 | 3.68 | 3.46 |
| LC3 | 5.89 | 6.80 | 6.35 |
| LN1 | 3.17 | 3.48 | 3.32 |
| LN2 | 3.96 | 3.80 | 3.88 |
| LN3 | 4.04 | 4.06 | 4.05 |
| (LLC/LC)LC1 | 5.57 | 4.30 | 4.93 |
| (LLC/LC)LC2 | 4.73 | 4.23 | 4.48 |
| (LLC/LC)LC3 | 5.73 | 5.05 | 5.39 |
| (LLN/LN)LN1 | 4.96 | 6.25 | 5.60 |
| (LLN/LN)LN2 | 4.39 | 3.29 | 3.84 |
| (LLN/LN)LN3 | 3.86 | 4.78 | 4.32 |
| Mean | 4.54 | 4.53 | 4.53 |

1st CUT MEAN DM% 31.2

LEYS 2^{ND} CUT (05/11/12) DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

| FYM_RES LEY | NONE | FYM | Mean |
|----------------|------|------|------|
| LC1 | 0.22 | 0.40 | 0.31 |
| LC2 | 0.42 | 0.68 | 0.55 |
| LC3 | 0.00 | 0.00 | 0.00 |
| LN1 | 0.55 | 0.61 | 0.58 |
| LN2 | 0.36 | 0.28 | 0.32 |
| LN3 | 0.00 | 0.00 | 0.00 |
| (LLC/LC)LC1 | 0.25 | 0.20 | 0.22 |
| (LLC/LC)LC2 | 1.05 | 1.01 | 1.03 |
| (LLC/LC)LC3 | 0.00 | 0.00 | 0.00 |
| (LLN/LN)LN1 | 1.10 | 1.56 | 1.33 |
| (LLN/LN)LN2 | 0.60 | 0.19 | 0.40 |
| (LLN/LN)LN3 | 0.00 | 0.00 | 0.00 |
| Mean | 0.38 | 0.41 | 0.40 |
| | | | |

2ND CUT MEAN DM% 33.4

TOTAL OF 2 CUTS DRY MATTER TONNES/HECTARE

***** TABLES OF MEANS *****

| FYM_RES | NONE | FYM | Mean |
|-------------|------|------|------|
| LEY | | | |
| LC1 | 5.11 | 5.07 | 5.09 |
| LC2 | 3.67 | 4.35 | 4.01 |
| LC3 | 5.89 | 6.80 | 6.35 |
| LN1 | 3.72 | 4.08 | 3.90 |
| LN2 | 4.32 | 4.09 | 4.20 |
| LN3 | 4.04 | 4.06 | 4.05 |
| (LLC/LC)LC1 | 5.82 | 4.49 | 5.15 |
| (LLC/LC)LC2 | 5.78 | 5.24 | 5.51 |
| (LLC/LC)LC3 | 5.73 | 5.05 | 5.39 |
| (LLN/LN)LN1 | 6.05 | 7.81 | 6.93 |
| (LLN/LN)LN2 | 5.00 | 3.49 | 4.24 |
| (LLN/LN)LN3 | 3.86 | 4.78 | 4.32 |
| Mean | 4.92 | 4.94 | 4.93 |

TOTAL OF 2 CUTS MEAN DM% 32.2

W. WHEAT (1st TEST CROP)

Grain tonnes/hectare

***** Tables of means *****

| FYMRES ROTATION | none | FYM | Mean | | |
|--|---|---|--|---|---|
| (AO) W | 5.73 | 6.88 | 6.31 | | |
| (ABe)W | 7.02 | 7.31 | 7.16 | | |
| (LLn/AO)W | 6.92 8.28 | 7.03 | 6.97 8.60 | | |
| (LLc/ABe)W (Ln)W | 8.28 7.16 | 8.91 7.05 | 8.60 7.10 | | |
| (LLN/Ln)W | 7.13 | 7.24 | 7.10 | | |
| (Lc)W | 8.18 | 8.04 | 8.11 | | |
| (LLc/Lc)W | 8.46 | 8.37 | 8.42 | | |
| Mean | 7.36 | 7.60 | 7.48 | | |
| N | 0 | 80 | 160 | 240 | Mean |
| ROTATION | 4 4 0 | 6 0 4 | 7 60 | 7 07 | C 01 |
| (AO)W (ABe)W | 4.40 3.92 | 6.04 6.57 | 7.52 8.73 | 7.27 9.44 | 6.31 7.16 |
| (LLn/AO)W | 3.45 | 7.45 | 8.20 | 8.79 | 6.97 |
| (LLc/ABe)W | 5.28 | 9.18 | 10.13 | 9.80 | 8.60 |
| (Ln)W | 4.39 | 7.52 | 8.38 | 8.13 | 7.10 |
| (LLN/Ln)W | 4.20 | 7.75 | 8.46 | 8.33 | 7.19 |
| (LC)W | 7.28 | 7.72 | 8.08 | 9.37 | 8.11 |
| (LLc/Lc)W | 6.61 | 9.17 | 8.91 | 8.98 | 8.42 |
| Mean | 4.94 | 7.68 | 8.55 | 8.76 | 7.48 |
| | | | | | |
| N | 0 | 80 | 160 | 240 | Mean |
| FYMRES | | | | | |
| FYMRES none | 4.52 | 7.79 | 8.50 | 8.62 | 7.36 |
| FYMRES | | | | | |
| FYMRES none FYM | 4.52 | 7.79 7.56 | 8.50 8.60 | 8.62 8.90 | 7.36 7.60 |
| FYMRES none FYM | 4.52 5.36 4.94 | 7.79 7.56 7.68 | 8.50 8.60 8.55 | 8.62 8.90 8.76 | 7.36 7.60 7.48 |
| FYMRES none FYM Mean | 4.52 5.36 4.94 N FYMRES none | 7.79 7.56 7.68 0 2.27 | 8.50 8.60 8.55 80 6.63 | 8.62 8.90 8.76 160 7.07 | 7.36 7.60 7.48 240 6.97 |
| FYMRES none FYM Mean ROTATION (AO)W | 4.52 5.36 4.94 N FYMRES none FYM | 7.79 7.56 7.68 0 2.27 6.53 | 8.50 8.60 8.55 80 6.63 5.46 | 8.62 8.90 8.76 160 7.07 7.97 | 7.36 7.60 7.48 240 6.97 7.57 |
| FYMRES none FYM Mean ROTATION | 4.52 5.36 4.94 N FYMRES none | 7.79 7.56 7.68 0 2.27 | 8.50 8.60 8.55 80 6.63 | 8.62 8.90 8.76 160 7.07 | 7.36 7.60 7.48 240 6.97 |
| FYMRES none FYM Mean ROTATION (AO)W | 4.52 5.36 4.94 N FYMRES none FYM none | 7.79 7.56 7.68 0 2.27 6.53 3.82 | 8.50 8.60 8.55 80 6.63 5.46 6.07 | 8.62 8.90 8.76 160 7.07 7.97 8.27 | 7.36 7.60 7.48 240 6.97 7.57 9.90 |
| FYMRES none FYM Mean ROTATION (AO)W (ABe)W (LLn/AO)W | 4.52 5.36 4.94 N FYMRES none FYM none FYM | 7.79 7.56 7.68 0 2.27 6.53 3.82 4.01 3.19 3.70 | 8.50 8.60 8.55 80 6.63 5.46 6.07 7.07 7.10 7.10 7.81 | 8.62 8.90 8.76 160 7.07 7.97 8.27 9.19 8.40 8.00 | 7.36 7.60 7.48 240 6.97 7.57 9.90 8.97 8.98 8.61 |
| FYMRES none FYM Mean ROTATION (AO)W (ABe)W | 4.52 5.36 4.94 N FYMRES none FYM none FYM none FYM none | 7.79 7.56 7.68 0 2.27 6.53 3.82 4.01 3.19 3.70 4.72 | 8.50 8.60 8.55 80 6.63 5.46 6.07 7.07 7.10 7.10 7.81 8.99 | 8.62 8.90 8.76 160 7.07 7.97 8.27 9.19 8.40 8.00 10.03 | 7.36 7.60 7.48 240 6.97 7.57 9.90 8.97 8.98 8.61 9.38 |
| FYMRES none FYM Mean ROTATION (AO)W (ABe)W (LLn/AO)W (LLc/ABe)W | 4.52 5.36 4.94 N FYMRES none FYM none FYM none FYM | 7.79 7.56 7.68 0 2.27 6.53 3.82 4.01 3.19 3.70 4.72 5.83 | 8.50 8.60 8.55 80 6.63 5.46 6.07 7.07 7.10 7.81 8.99 9.38 | 8.62 8.90 8.76 160 7.07 7.97 8.27 9.19 8.40 8.00 10.03 10.23 | 7.36 7.60 7.48 240 6.97 7.57 9.90 8.97 8.98 8.61 9.38 10.21 |
| FYMRES none FYM Mean ROTATION (AO)W (ABe)W (LLn/AO)W | 4.52 5.36 4.94 N FYMRES None FYM none FYM none FYM none FYM none FYM | 7.79 7.56 7.68 0 2.27 6.53 3.82 4.01 3.19 3.70 4.72 5.83 4.66 | 8.50 8.60 8.55 80 6.63 5.46 6.07 7.07 7.10 7.81 8.99 9.38 7.41 | 8.62 8.90 8.76 160 7.07 7.97 8.27 9.19 8.40 8.00 10.03 10.23 8.57 | 7.36 7.60 7.48 240 6.97 7.57 9.90 8.97 8.98 8.61 9.38 10.21 8.00 |
| FYMRES none FYM Mean ROTATION (AO)W (ABe)W (LLn/AO)W (LLc/ABe)W (Ln)W | 4.52 5.36 4.94 N FYMRES none FYM none FYM none FYM none FYM | 7.79 7.56 7.68 0 2.27 6.53 3.82 4.01 3.19 3.70 4.72 5.83 4.66 4.12 | 8.50 8.60 8.55 80 6.63 5.46 6.07 7.07 7.10 7.81 8.99 9.38 7.41 7.62 | 8.62 8.90 8.76 160 7.07 7.97 8.27 9.19 8.40 8.00 10.03 10.23 8.57 8.19 | 7.36 7.60 7.48 240 6.97 7.57 9.90 8.97 8.98 8.61 9.38 10.21 8.00 8.25 |
| FYMRES none FYM Mean ROTATION (AO)W (ABe)W (LLn/AO)W (LLc/ABe)W | 4.52 5.36 4.94 N FYMRES None FYM none FYM none FYM none FYM none FYM | 7.79 7.56 7.68 0 2.27 6.53 3.82 4.01 3.19 3.70 4.72 5.83 4.66 | 8.50 8.60 8.55 80 6.63 5.46 6.07 7.07 7.10 7.81 8.99 9.38 7.41 | 8.62 8.90 8.76 160 7.07 7.97 8.27 9.19 8.40 8.00 10.03 10.23 8.57 | 7.36 7.60 7.48 240 6.97 7.57 9.90 8.97 8.98 8.61 9.38 10.21 8.00 |
| FYMRES none FYM Mean ROTATION (AO)W (ABe)W (LLn/AO)W (LLc/ABe)W (Ln)W | 4.52 5.36 4.94 N FYMRES none FYM none FYM none FYM none FYM none FYM | 7.79 7.56 7.68 0 2.27 6.53 3.82 4.01 3.19 3.70 4.72 5.83 4.66 4.12 4.17 | 8.50 8.60 8.55 80 6.63 5.46 6.07 7.07 7.10 7.81 8.99 9.38 7.41 7.62 7.87 7.64 8.93 | 8.62 8.90 8.76 160 7.07 7.97 8.27 9.19 8.40 8.00 10.03 10.23 8.57 8.19 7.95 | 7.36 7.60 7.48 240 6.97 7.57 9.90 8.97 8.98 8.61 9.38 10.21 8.00 8.25 8.52 |
| FYMRES none FYM Mean ROTATION (AO)W (ABe)W (LLn/AO)W (LLc/ABe)W (Ln)W (LLN/Ln)W (LC)W | 4.52 5.36 4.94 N FYMRES none FYM none FYM none FYM none FYM none FYM | 7.79 7.56 7.68 0 2.27 6.53 3.82 4.01 3.19 3.70 4.72 5.83 4.66 4.12 4.17 4.22 7.29 7.27 | 8.50 8.60 8.55 80 6.63 5.46 6.07 7.07 7.10 7.81 8.99 9.38 7.41 7.62 7.87 7.64 8.93 6.51 | 8.62 8.90 8.76 160 7.07 7.97 8.27 9.19 8.40 8.00 10.03 10.23 8.57 8.19 7.95 8.97 7.93 8.22 | 7.36 7.60 7.48 240 6.97 7.57 9.90 8.97 8.98 8.61 9.38 10.21 8.00 8.25 8.52 8.52 8.15 8.56 10.18 |
| FYMRES none FYM Mean ROTATION (AO)W (ABe)W (LLn/AO)W (LLc/ABe)W (Ln)W (LLN/Ln)W | 4.52 5.36 4.94 N FYMRES none FYM none FYM none FYM none FYM none FYM none FYM | 7.79 7.56 7.68 0 2.27 6.53 3.82 4.01 3.19 3.70 4.72 5.83 4.66 4.12 4.17 4.22 7.29 | 8.50 8.60 8.55 80 6.63 5.46 6.07 7.07 7.10 7.81 8.99 9.38 7.41 7.62 7.87 7.64 8.93 | 8.62 8.90 8.76 160 7.07 7.97 8.27 9.19 8.40 8.00 10.03 10.23 8.57 8.19 7.95 8.97 7.93 | 7.36 7.60 7.48 240 6.97 7.57 9.90 8.97 8.98 8.61 9.38 10.21 8.00 8.25 8.52 8.52 8.15 8.56 |

W.WHEAT - GRAIN MEAN DM% 85.3

PLOT AREA HARVESTED 0.00192

RYE EXTRA

GRAIN (85% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

| FYMRES ROTATION | NONE | FYM | Mean |
|--------------------|------|------|------|
| (ABe)R | 1.99 | 1.65 | 1.82 |
| (AO) R | 1.49 | 1.24 | 1.36 |
| (LLn/AO)R | 2.31 | 2.65 | 2.48 |
| (LLc/ABe)R | 2.99 | 2.55 | 2.77 |
| | | | |
| Mean | 2.19 | 2.02 | 2.11 |

GRAIN MEAN DM% 81.8

PLOT AREA HARVESTED 0.00413

W. RYE (2^{ND} TEST CROP)

12/W/RN/3 W. RYE

Grain tonnes/hectare

***** Tables of means *****

| FYMRES ROTATION | none | FYM | Mean | | |
|--|--|--|--|--|--|
| (AO) R | 2.37 | 2.74 | 2.55 | | |
| (ABe)R (LLn/AO)R | 2.37 2.83 | 2.91 2.96 | 2.64 2.89 | | |
| (LLc/ABe)R | 3.07 | 2.50 | 2.87 | | |
| (Ln)R | 3.38 | 2.97 | 3.17 | | |
| (LLn/Ln)R | 3.27 | 3.44 | 3.36 | | |
| (Lc)R | 3.33 | 3.45 | 3.39 | | |
| (LLc/Lc)R | 3.17 | 3.07 | 3.12 | | |
| Mean | 2.98 | 3.03 | 3.00 | | |
| | | | | | |
| N | 0 | 50 | 100 | 150 | Mean |
| N ROTATION | 0 | 50 | 100 | 150 | Mean |
| | 0 2.03 | 50 2.55 | 100 2.76 | 150 2.88 | Mean 2.55 |
| ROTATION | | | | | |
| ROTATION (AO) R | 2.03 | 2.55 | 2.76 | 2.88 | 2.55 |
| ROTATION (AO)R (ABe)R | 2.03 | 2.55 | 2.76 3.01 | 2.88 | 2.55 |
| ROTATION (AO)R (ABe)R (LLn/AO)R | 2.03 2.02 2.19 | 2.55 2.65 3.12 | 2.76 3.01 2.93 | 2.88 2.89 3.33 | 2.55 2.64 2.89 |
| ROTATION (AO)R (ABe)R (LLn/AO)R (LLc/ABe)R | 2.03 2.02 2.19 2.37 | 2.55 2.65 3.12 3.06 | 2.76 3.01 2.93 3.00 | 2.88 2.89 3.33 3.06 | 2.55 2.64 2.89 2.87 |
| ROTATION (AO)R (ABe)R (LLn/AO)R (LLc/ABe)R (LLn)R | 2.03 2.02 2.19 2.37 3.36 | 2.55 2.65 3.12 3.06 3.07 | 2.76 3.01 2.93 3.00 3.31 | 2.88 2.89 3.33 3.06 2.96 | 2.55 2.64 2.89 2.87 3.17 |
| ROTATION (AO)R (ABe)R (LLn/AO)R (LLc/ABe)R (Ln)R (LLn/Ln)R | 2.03 2.02 2.19 2.37 3.36 3.21 | 2.55 2.65 3.12 3.06 3.07 3.63 | 2.76 3.01 2.93 3.00 3.31 3.18 | 2.88 2.89 3.33 3.06 2.96 3.41 | 2.55 2.64 2.89 2.87 3.17 3.36 |

| N FYMRES | 0 | 50 | 100 | 150 | Mean |
|-------------|--------|------|------|------|------|
| none | 2.59 | 3.08 | 3.16 | 3.07 | 2.98 |
| FYM | 2.87 | 2.99 | 3.12 | 3.12 | 3.03 |
| Mean | 2.73 | 3.04 | 3.14 | 3.09 | 3.00 |
| Mean | 2.15 | 3.04 | 3.14 | 5.09 | 3.00 |
| | N | 0 | 50 | 100 | 150 |
| ROTATION | FYMRES | | | | |
| (AO) R | none | 1.71 | 2.35 | 2.58 | 2.85 |
| | FYM | 2.35 | 2.74 | 2.95 | 2.90 |
| (ABe)R | none | 1.73 | 2.31 | 2.73 | 2.73 |
| | FYM | 2.32 | 2.99 | 3.29 | 3.04 |
| (LLn/AO)R | none | 1.98 | 3.00 | 2.99 | 3.34 |
| | FYM | 2.40 | 3.24 | 2.87 | 3.32 |
| (LLc/ABe)R | none | 2.35 | 2.95 | 3.45 | 3.53 |
| | FYM | 2.39 | 3.16 | 2.54 | 2.59 |
| (Ln)R | none | 3.42 | 3.62 | 3.39 | 3.10 |
| | FYM | 3.30 | 2.51 | 3.23 | 2.82 |
| (LLn/Ln)R | none | 2.87 | 3.87 | 3.15 | 3.21 |
| , | FYM | 3.55 | 3.40 | 3.22 | 3.61 |
| (Lc)R | none | 3.56 | 3.27 | 3.82 | 2.68 |
| / | FYM | 3.48 | 3.05 | 3.69 | 3.56 |
| (LLc/Lc)R | none | 3.12 | 3.25 | 3.22 | 3.08 |
| | FYM | 3.15 | 2.86 | 3.15 | 3.13 |
| | | | | | |

GRAIN MEAN DM% 83.9 PLOT AREA HARVESTED 0.00192

BEANS

GRAIN (85% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

| FYMRES ROTATION | NONE | FYM | Mean |
|--------------------|------|------|------|
| (AO)Be | * | * | * |
| (LLn/AO)Be | * | * | * |
| (LLc/ABe)Be | 0.12 | 0.36 | 0.24 |
| (ABe)Be | 0.40 | 1.62 | 1.01 |
| Mean | 0.26 | 0.99 | 0.63 |

GRAIN MEAN DM% 80.2

0.00413 PLOT AREA HARVESTED

Note: Due to wet weather and severe weed infestation no yields were obtained for (AO)Be and (LLn/AO)Be treatment plots.

OATS

GRAIN (85% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

| FYMRES ROTATION | NONE | FYM | Mean |
|--------------------|------|------|------|
| ABe | 1.28 | 1.42 | 1.35 |
| AO | 2.36 | 2.07 | 2.21 |
| LLc/ABe | 1.71 | 1.73 | 1.72 |
| LLn/AO | 3.30 | 3.18 | 3.24 |
| Mean | 2.16 | 2.10 | 2.13 |

GRAIN MEAN DM% 82.9

PLOT AREA HARVESTED 0.00413