Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 2013



Full Table of Content

# R/EX/4 Exhaustion Land

# **Rothamsted Research**

Rothamsted Research (2014) *R/EX/4 Exhaustion Land*; Yields Of The Field Experiments 2013, pp 22 - 25 - **DOI:** https://doi.org/10.23637/ERADOC-1-223

#### 13/R/EX/4

#### **EXHAUSTION LAND**

**Object**: To study the residual effects of manures applied 1856 - 1901, and of additional phosphate applied since 1986 (P test) and of additional potassium since 2007 (K test); on the yield of continuous s. barley up to 1991, w. wheat since – Hoosfield.

The 158th year, w. wheat.

For previous years see 'Details' 1977, 1973 and Yield Books for 74-12/R/EX/4

Treatments: All combinations of:-

Whole plots (P test)

OLD RES	Residues of manures applied annually 1876 – 1901:			
	None Farmyard manure at 35 t 96 kg N as ammonium salts 34 kg P as superphosphate N and P as above plus 137 kg K as sulphate of potash, 16 kg Na as sulphate of soda, 11 kg Mg as sulphate of magnesia			
P	Maintenance P (20 kg P) applied annually from 2000 to maintain existing levels of available P In the soil. (P1) (P2) and (P3) are residues of P applied annually 1986–1992:			
O P (P1) P (P2) P (P3)	2000-12 None 20 kg P 20 kg P 20 kg P	1986-92 None 44 kg P 87 kg P 131 kg P		
	D N P NPKNAMG  P O P (P1) P (P2)	O None D Farmyard manure at 35 to 10 None P 34 kg P as superphosphators Nand P as above plus 10 potash, 16 kg Na as sulph sulphate of magnesia  P Maintenance P (20 kg P) to maintain existing lever (P1) (P2) and (P3) are resulted to 1986–1992:  2000-12 O None P (P1) 20 kg P P (P2) 20 kg P		

**NOTE**: P treatments were applied at 61.5 kg P in error in 2000.

Plus

Whole plots (K test, previously N test until 1991

1. OLD RES	Residues of manures applied annually 1876 – 1901:		
0	None		
D	Farmyard manure at 35 t		
N*	96 kg N as nitrate of soda		
PK	34 kg P as superphosphate, 137 kg K as sulphate of potash		
N*PK	N, P and K as above		

# 13/R/EX/4

2. K	Potassium applied annually from 2007 as muriate of p	otash

O None

 $\begin{array}{ccc} K1 & 75 \text{ kg } \text{K}_2\text{O } (62.2 \text{ kg K}) \\ K2 & 150 \text{ kg } \text{K}_2\text{O } (124.5 \text{ kg K}) \end{array}$ 

Whole plots

Nitrogen: 50 kg N as ammonium sulphate (to supply sufficient S) during first two weeks

in March, 200 kg N as ammonium nitrate at GS31/mid-April (whichever comes first) and 50 kg N as ammonium nitrate at GS37 (not later than mid-

May)

# **Experimental diary**

Date		Application	Rate	Units
20-Sep-12	р	Sprayed Whole field w/ Weedazol EW	20	l/ha
28-Sep-12	f	Spread MOP Fertilizer on plots - 103,83,63,43,23	125	kg/ha
28-Sep-12	f	Spread MOP Fertilizer as on sheet on plots 104-94,87-73,64-54,44-34,24-14	250	kg/ha
28-Sep-12	f	Spread TSP as on sheet on plots 101-93,81-73,61-53,41-33,21-13	75	kg/ha
15-Oct-12	а	Drilled WW Xi19	400	seeds m <sup>2</sup>
15-Nov-12	р	Sprayed Liberator and PDM, also o+e's at each end of EX/4	Li@0.6 PDM@2.7	l/ha l/ha
15-Nov-12	p	Spread Slug Pellets (Gusto)	5	kg/ha
11-Mar-13	f	Applied Kieserite	80	kg/ha
12-Mar-13	f	Applied Ammonium Sulphate Fertiliser	238	kg/ha
25-Apr-13	f	Applied Nitram	580	kg/ha
16-May-13	p	Sprayed AllyMax, Kingdom, Bravo500, NewCycocel, HatchetExtra	AM@42 Ki@1.25	g/ha l/ha
			Br500@1.0	l/ha
			NCy@2.0 Hat@1.0	l/ha l/ha
21-May-13	f	Applied Nitram	145	kg/ha
06-Jun-13	р	Sprayed Ingite and Comet	ign@1.2 com@0.4	l/ha l/ha
18-Jun-13	р	Sprayed Cello	0.55	l/ha
19-Jun-13	а	Cut Paths		_
04-Jul-13	а	Cut Paths	_	_
19-Jul-13	а	Cut Paths.		_
12-Aug-13	а	Claas - Harvested	_	_
13-Aug-13	а	Sampo - Harvested	_	_
13-Aug-13	а	baled weighed and sampled plots	_	_
20-Aug-13	а	Claas - Harvested, OE's	_	_

Note: Samples of grain and straw were taken for chemical analysis. The yield strips on plots 031, 034, 071, 074, 091 & 094 were made smaller this year to avoid areas where the crop had already been sampled by S. McGrath et al.

P TEST

Grain tonnes/hectare

\*\*\*\* Tables of means \*\*\*\*

P_RES OLD_RES	0	P1	P2	Р3	Mean
_ 0	1.83	4.32	4.45	5.45	4.01
D	3.24	5.49	6.00	5.87	5.15
N	0.89	4.78	5.15	5.81	4.16
P	3.24	5.25	6.10	6.05	5.16
NPKNAMG	3.60	5.35	5.83	6.64	5.35
Mean	2 56	5 04	5 51	5 96	4 77

Grain mean DM% 86.3

Straw tonnes/hectare

\*\*\*\*\* Tables of means \*\*\*\*\*

P_RES	0	P1	P2	Р3	Mean
OLD_RES					
_ o	0.51	2.21	2.17	2.98	1.97
D	0.80	2.72	3.25	3.47	2.56
N	0.29	2.07	2.32	2.65	1.83
P	1.15	2.55	2.85	3.36	2.47
NPKNAMG	1.24	2.46	3.26	4.01	2.74
Mean	0.80	2.40	2.77	3.29	2.32

Straw mean DM% 87.1

Plot area harvested 0.00538, 0.00252.

#### 13/R/EX/4

#### K TEST

#### Grain tonnes/hectare

#### \*\*\*\* Tables of means \*\*\*\*

K_Test	K0	K1	К2	Mean
OLD_RES				
_ o	4.99	6.10	6.51	5.64
D	5.89	6.59	6.76	6.28
N*	4.92	5.26	5.93	5.26
PK	6.04	6.57	6.67	6.33
N*PK	5.58	6.45	7.06	6.17
Mean	5.48	6.19	6.59	5.94

Grain mean DM% 88.1

#### Straw tonnes/hectare

# \*\*\*\*\* Tables of means \*\*\*\*\*

<b>K_Test</b>	K0	K1	K2	Mean
OLD_RES				
_ 0	1.92	2.66	2.83	2.34
D	2.38	3.15	3.27	2.80
N*	1.67	2.23	2.45	2.01
PK	2.81	3.06	2.82	2.87
N*PK	2.36	3.47	3.39	2.89
Mean	2.23	2.91	2.95	2.58

Straw mean DM% 92.4 Plot area harvested 0.00538