

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 2015



Results of the
Classical and other
Long-term Experiments
2015

[Full Table of Content](#)

R/EX/4 Exhaustion Land

Rothamsted Research

Rothamsted Research (2016) *R/EX/4 Exhaustion Land* ; Yields Of The Field Experiments 2015, pp 24 - 27 - DOI: <https://doi.org/10.23637/ERADOC-1-225>

15/R/EX/4

EXHAUSTION LAND

Object: To study the residual effects of manures applied 1856 - 1901, and of additional phosphate applied since 1986 (P test) and of additional potassium since 2007 (K test); on the yield of continuous s. barley up to 1991, w. wheat since – Hoosfield.

The 160th year, w. wheat.

For previous years see 'Details' 1977, 1973 and Yield Books for 74-14/R/EX/4

Treatments: All combinations of:-

Whole plots (P test)

1. **OLD RES** Residues of manures applied annually 1876 – 1901:

O	None
D	Farmyard manure at 35 t
N	96 kg N as ammonium salts
P	34 kg P as superphosphate
NPKNAMG	N and P as above plus 137 kg K as sulphate of potash, 16 kg Na as sulphate of soda, 11 kg Mg as sulphate of magnesia

2. **P** Maintenance P (20 kg P) applied annually from 2000 to maintain existing levels of available P in the soil. In 2009 maintenance P applications were changed from 20 kg P/ha to 15 kg P/ha. This was not recorded in the yield books for 2009-13. (P1) (P2) and (P3) are residues of P applied annually 1986–1992:

	2009-Present	2000-08	1986-92
O	None	None	None
P (P1)	15 kg P	20 kg P	44 kg P
P (P2)	15 kg P	20 kg P	87 kg P
P (P3)	15 kg P	20 kg P	131 kg P

NOTE: P treatments were applied at 61.5 kg P in error in 2000.

Plus

Whole plots (K test, previously N test until 1991)

1. **OLD RES** Residues of manures applied annually 1876 – 1901:

O	None
D	Farmyard manure at 35 t
N*	96 kg N as nitrate of soda
PK	34 kg P as superphosphate, 137 kg K as sulphate of potash
N*PK	N, P and K as above

2. **K** Potassium applied annually from 2007 as muriate of potash

O	None
K1	75 kg K ₂ O (62.2 kg K)
K2	150 kg K ₂ O (124.5 kg K)
Whole plots	
Nitrogen:	50 kg N as ammonium sulphate (to supply sufficient S) during first two weeks in March, 200 kg N as ammonium nitrate at GS31/mid-April (whichever comes first) and 50 kg N as ammonium nitrate at GS37 (not later than mid-May)

Experimental diary

Date		Application	Rate	Units
02/10/2014	f	Applied TSP	75	kg/ha
02/10/2014	f	Applied MOP onto plots 103 83 63 43 + 23	125	kg/ha
02/10/2014	f	Applied MOP onto plots 104-24, 91-11, 92-12, 93-13, 94-14.	250	kg/ha
08/10/2014	a	Ploughed ground (thrown North)	-	-
28/10/2014	s	Drilled Crusoe trt Redigo Deter	400	seeds/m ²
01/11/2014	a	Ring rolled all new drilling	-	-
10/11/2014	p	Sprayed Crystal	4	l/ha
04/12/2014	p	Sprayed Hallmark	50	ml/ha
23/03/2015	f	Applied Ammonia Sulphate (21%N) onto plots	238	kg/ha
05/04/2015	p	Sprayed Artemis	1	l/ha
05/04/2015	p	Sprayed Bravo 500	1	l/ha
05/04/2015	p	Sprayed Chlormequat 750	1.25	l/ha
05/04/2015	p	Sprayed Moddus	150	ml/ha
09/04/2015	f	Applied Nitram fertiliser (34.5%N) to Winter Wheat Plots	580	kg/ha
16/04/2015	f	Applied Kieserite to Winter Wheat Plots	80	kg/ha
27/04/2015	p	Sprayed Kingdom	1.25	l/ha
27/04/2015	p	Sprayed Balear720	700	ml/ha
30/04/2015	f	Applied Nitram fertiliser (34.5%N) to Winter Wheat Plots	145	kg/ha
27/05/2015	p	Sprayed Ally Max	30	gm/ha
27/05/2015	p	Sprayed Bassoon	750	ml/ha
27/05/2015	p	Sprayed Gemstone	1	l/ha
27/05/2015	p	Sprayed Bravo500	1	l/ha
15/06/2015	p	Sprayed Proline onto WW	500	ml/ha
06/08/2015	a	Topped Paths	-	-
07/09/2015	a	Harvested All Commercial WW; Swathed Straw	-	-
13/09/2015	a	Harvested All Plots for Grain Yield	-	-
13/09/2015	a	Sampled Baled and Weighed	-	-
17/09/2015	a	removed round bales from field	-	-

Note: Samples of grain and straw were taken for chemical analysis. The yield strips on plots 031, 034, 071, 074, 091 & 094 were made smaller this year to avoid areas where the crop had already been sampled by S. McGrath et al.

15/R/EX/4

P TEST

Grain tonnes/hectare

***** Tables of means *****

P_RES	O	P1	P2	P3	Mean
OLD_RES					
O	3.56	8.59	8.38	8.18	7.18
D	5.58	9.77	9.80	9.62	8.69
N	3.08	7.98	8.95	8.98	7.25
P	*7.31	*12.03	*10.62	*10.85	*10.20
NPKNAMG	5.15	9.37	10.07	10.86	8.86
Mean	4.94	9.56	9.57	9.65	8.43

Grain mean DM% 85.1

*Note: Yields estimated using grain/straw ratios for plots 072,073 & 074, because of a problem with the combine when harvesting.

Straw tonnes/hectare

***** Tables of means *****

P_RES	O	P1	P2	P3	Mean
OLD_RES					
O	1.91	3.99	4.17	4.17	3.56
D	2.21	4.59	4.78	4.71	4.07
N	2.05	4.29	4.55	3.94	3.71
P	3.18	6.33	5.59	5.71	5.20
NPKNAMG	2.24	4.82	5.41	5.52	4.50
Mean	2.32	4.80	4.90	4.81	4.21

Straw mean DM% 88.3

Plot area harvested 0.00538, 0.00252.

K TEST

Grain tonnes/hectare

***** Tables of means *****

K_Test	K0	K1	K2	Mean
OLD_RES				
O	7.61	8.90	9.14	8.31
D	7.57	9.77	9.90	8.70
N*	8.03	8.60	8.87	8.38
PK	9.70	9.96	10.13	9.87
N*PK	8.54	10.08	10.27	9.36
Mean	8.29	9.46	9.66	8.93

Grain mean DM% 85.0

15/R/EX/4

Straw tonnes/hectare

***** Tables of means *****

K_Test	K0	K1	K2	Mean
OLD_RES				
O	3.80	4.69	5.13	4.35
D	3.37	4.37	4.39	3.88
N*	3.98	4.34	4.34	4.16
PK	5.40	5.40	5.53	5.44
N*PK	4.36	5.13	5.2	4.76
Mean	4.18	4.78	4.92	4.52

Straw mean DM% 87.4 Plot area harvested 0.00538