

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments

[Full Table of Content](#)



R/EX/4 Exhaustion Land

Rothamsted Research

Rothamsted Research (2016) *R/EX/4 Exhaustion Land* ; Yields Of The Field Experiments , pp 24 - 27
- DOI: <https://doi.org/10.23637/ERADOC-1-249>

16/R/EX/4

EXHAUSTION LAND

Object: To study the residual effects of manures applied 1856 - 1901, and of additional phosphate applied since 1986 (P test) and of additional potassium since 2007 (K test); on the yield of continuous s. barley up to 1991, w. wheat since – Hoosfield.

The 161st year, w. wheat.

For previous years see 'Details' 1977, 1973 and Yield Books for 74-14/R/EX/4

Treatments: All combinations of:-

Whole plots (P test)

1. **OLD RES** Residues of manures applied annually 1876 – 1901:

O	None
D	Farmyard manure at 35 t
N	96 kg N as ammonium salts
P	34 kg P as superphosphate
NPKNAMG	N and P as above plus 137 kg K as sulphate of potash, 16 kg Na as sulphate of soda, 11 kg Mg as sulphate of magnesia

2. **P** Maintenance P (20 kg P) applied annually from 2000 to maintain existing levels of available P in the soil. In 2009 maintenance P applications were changed from 20 kg P/ha to 15 kg P/ha. This was not recorded in the yield books for 2009-13. (P1) (P2) and (P3) are residues of P applied annually 1986–1992:

	2009-Present	2000-08	1986-92
O	None	None	None
P (P1)	15 kg P	20 kg P	44 kg P
P (P2)	15 kg P	20 kg P	87 kg P
P (P3)	15 kg P	20 kg P	131 kg P

NOTE: P treatments were applied at 61.5 kg P in error in 2000.

Plus

Whole plots (K test, previously N test until 1991)

1. **OLD RES** Residues of manures applied annually 1876 – 1901:

O	None
D	Farmyard manure at 35 t
N*	96 kg N as nitrate of soda
PK	34 kg P as superphosphate, 137 kg K as sulphate of potash
N*PK	N, P and K as above

2. **K** Potassium applied annually from 2007 as muriate of potash

O None
 K1 75 kg K₂O (62.2 kg K)
 K2 150 kg K₂O (124.5 kg K)
 Whole plots
 Nitrogen: 50 kg N as ammonium sulphate (to supply sufficient S) during first two weeks in March, 200 kg N as ammonium nitrate at GS31/mid-April (whichever comes first) and 50 kg N as ammonium nitrate at GS37 (not later than mid-May)

Experimental diary

Date		Application	Rate	Units
25/08/2015	p	Sprayed Firebrand	1	lt/ha
25/08/2015	p	Sprayed Samurai	4	lt/ha
01/10/2015	f	Applied MOP - Plots 103, 83, 63, 43 + 23	125	kg/ha
01/10/2015	f	Applied MOP - Plots 094-014, 092-012, 104-024, 093-013, 091-011	250	kg/ha
01/10/2015	f	Applied TSP - plots 101-021, 102-022, 103-023, 104-024, 091-011 + 092-012	75	kg/ha
09/10/2015	a	Ploughed; thrown south	-	-
09/10/2015	a	Topped all field and exp - Batwing	-	-
15/10/2015	a	Power harrowed - Seed bed preparation	-	-
15/10/2015	s	Drilled Crusoe trt Redigo	350	seed/m ²
18/10/2015	a	Ring Rolled all new drilling	-	-
19/10/2015	p	Sprayed WW - Liberater	0.6	lt/ha
19/10/2015	p	Sprayed WW - Stomp Aqua	1.75	lt/ha
12/11/2015	p	Sprayed Hallmark	40	ml/ha
03/12/2015	p	Applied TDS Major	5	kg/ha
12/12/2015	p	Sprayed Hallmark	40	ml/ha
21/03/2016	f	Applied Sulphate of Ammonia	238	kg/ha
04/04/2016	f	Applied Nitram @ 34.5%N	840	kg/ha
08/04/2016	p	Sprayed Moddus	150	ml/ha
08/04/2016	p	Sprayed BASF 3C Chlormequat750	1.25	lt/ha
08/04/2016	p	Sprayed Odin	500	ml/ha
08/04/2016	p	Sprayed Mirage 40ec	500	ml/ha
08/04/2016	p	Sprayed Bravo500	1	lt/ha
13/04/2016	f	Applied Kieserite - all plots	80	kg/ha
05/05/2016	p	Sprayed Keystone - wheat only	800	ml/ha
05/05/2016	p	Sprayed Keystone - wheat only	700	ml/ha
24/05/2016	p	Sprayed Vortex	1.5	lt/ha
24/05/2016	p	Sprayed VortexCorbel	250	ml/ha
09/07/2016	p	Sprayed Cello	550	ml/ha
09/07/2016	p	Sprayed Hallmark	50	ml/ha
22/08/2016	a	Cut Paths using Iseki and Mower	-	-
23/08/2016	a	Mopped Up remaining crop	-	-
25/08/2016	a	Harvested All Plots	-	-

26/08/2016 a Combined all crop using Tucano - -
 25/08/2015 p Sprayed Firebrand 1 lt/ha

Note: Samples of grain and straw were taken for chemical analysis. The yield strips on plots 031-034, 071- 074, 091- 094 (12 plots in total) were made smaller this year to avoid areas where the crop had already been sampled by S. McGrath et al.

16/R/EX/4

P TEST

Grain tonnes/hectare

***** Tables of means *****

P_RES	O	P1	P2	P3	Mean
OLD_RES					
O	2.48	7.72	8.51	9.12	6.96
D	5.66	10.35	10.85	11.49	9.59
N	2.33	8.07	9.90	9.59	7.47
P	4.23	9.89	10.93	11.27	9.08
NPKNAMG	3.96	7.98	11.11	11.83	8.72
Mean	3.73	8.80	10.26	10.66	8.36

Grain mean DM% 85.2

Straw tonnes/hectare

***** Tables of means *****

P_RES	O	P1	P2	P3	Mean
OLD_RES					
O	2.49	6.84	7.76	7.49	6.14
D	5.33	9.5	9.49	8.94	8.32
N	1.90	7.56	8.45	7.68	6.40
P	3.59	7.56	8.47	8.05	6.92
NPKNAMG	3.92	7.22	9.13	8.80	7.27
Mean	3.45	7.74	8.66	8.19	7.01

Straw mean DM% 92.1

Plot area harvested 0.00538, 0.00252.

K TEST

Grain tonnes/hectare

***** Tables of means *****

16/R/EX/4

K_Test	K0	K1	K2	Mean
OLD_RES				
O	9.14	10.50	10.48	9.82
D	8.76	10.92	10.86	9.83
N*	9.44	10.52	9.84	9.81
PK	10.54	10.51	10.18	10.44
N*PK	9.56	10.75	11.14	10.25
Mean	9.49	10.64	10.50	10.03

Grain mean DM% 85.5

Straw tonnes/hectare

***** Tables of means *****

K_Test	K0	K1	K2	Mean
OLD_RES				
O	6.70	8.34	8.40	7.53
D	6.55	8.05	8.12	7.32
N*	6.08	7.53	7.64	6.83
PK	7.22	7.08	7.19	7.18
N*PK	6.31	7.10	7.40	6.78
Mean	6.57	7.62	7.75	7.13

Straw mean DM% 96.6 Plot area harvested 0.00538