

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Results of the Classical and Other Long-term Experiments - 2017



Results of the
Classical and other
Long-term Experiments
2017

[Full Table of Content](#)

17/R/HB/2 Hoos Barley (Hoosfield)

Rothamsted Research

Rothamsted Research (2018) *17/R/HB/2 Hoos Barley (Hoosfield)* ; Results Of The Classical And Other Long-Term Experiments - 2017, pp 12 - 18 - DOI: <https://doi.org/10.23637/ERADOC-1-251>

Results of the Classical and other Long-term Experiments 2017

17/R/HB/2 HOOS BARLEY (Hoosfield)

Object: To study the effects of organic manures and inorganic fertilizers on continuous spring barley. From 1968 to 1978 a rotation of potatoes, beans and spring barley was practised on parts of the experiment. The rotation was discontinued in 1979 and the whole experiment reverted to continuous spring barley. The experiment was modified for 2003. The main plots continue as previously. The Silicate Test plots continue but are not split to test rates of N (basal N is applied). The remaining plots are to be used to study the effect on yield of P residues, (basal N applied).

The 166th year, spring barley.

For previous years see 'Details' 1967 and 1973, Station Report for 1966 and Yield Books for 74-16/R/HB/2.

Main plots

Treatments:

Whole plots

MANURE	Plot	Fertilizers and Organic Manures:-		
		Form of N 1852-1966	Additional treatments 1852-2002	Treatments since 2003
---	11	None	-	-
-P-	21	None	P	(P)
--K	31	None	K (Na) Mg	K(Mg)
-PK	41	None	PK (Na) Mg	(P) K (Mg)
A--	12	A	-	-
AP-	22	A	P	(P)
A-K	32	A	K (Na) Mg	K(Mg)
APK	42	A	PK (Na) Mg	(P) K (Mg)
D1852	72	None	D	D
(D)	71	None	(D)	(D)
(A)	62	None	(Ashes)	(Ashes)
-	61	None	-	-
D2001 ^(a)	73 ^(a)	-	D	D
P2KMg ^(a)	63 ^(a)	-	P2KMg	P2KMg

^(a) Plots 63 and 73 started in 2001

- Form of N: A, sulphate of ammonia to supply 48kg N
- P: 35 kg P as triple superphosphate in 1974 and from 1988 to 2002, single superphosphate in other years
- (P): (none), P application to be reviewed for 2018
- P2: 44kg P as triple superphosphate
- K: 90 kg K as sulphate of potash
- (Na): (none), 16 kg Na as sulphate of soda until 1973
- Mg: 35kg Mg as kieserite every third year since 1974 (applied at 30 kg in 1992, 1995 and 1998) (sulphate of magnesia annually until 1973). Annually to new plot 63.
- (Mg): (none), Mg application to be reviewed for 2021

Results of the Classical and other Long-term Experiments 2017

- D1852: Farmyard manure at 35t since 1852
 D2001: Farmyard manure at 35t since 2001
 (D): Farmyard manure 1852 – 1871 only
 (Ashes): Weed ash 1852-1916, furnace ash 1917-1932, none since

Sub-Plots

- (2) N Nitrogen fertilizer (kg N), as 'Nitro-Chalk', since 1968 (cumulative N applications until 1973, on a cyclic system since 1974):
 0
 48
 96
 144

Silicate Test plots

Treatments:

Whole plots

MANURE	Plot	Fertilizers:- Additional treatment 1852-1979	Changes since 1980	Treatments since 2003
N----	131	-	-	N3
NP---	231	P	-	N3 (P)
N-K--	331	K(Na)Mg	-	N3 K(Mg)
NPK--	431	PK(Na)Mg	-	N3(P)K(Mg)
N—S-	134	Si	Si omitted	N3 (Si)
NP-S-	234	P Si	Si omitted	N3(P) (Si)
N-KS-	334	K(Na)MgSi	Si omitted	N3 K(Mg)(Si)
NPKS-	434	PK(Na)MgSi	Si omitted	N3(P)K(Mg)(Si)
N---S	132	-	Si added	N3 Si
NP--S	232	P	Si added	N3(P) Si
N-K-S	332	K(Na)Mg	Si added	N3 K(Mg) Si
NPK-S	432	PK(Na)Mg	Si added	N3(P)K(Mg) Si
N--SS	133	Si	-	N3 Si
NP-SS	233	P Si	-	N3(P) Si
N-KSS	333	K(Na)MgSi	-	N3 K(Mg) Si
NPKSS	433	PK(Na)MgSi	-	N3(P)K(Mg) Si

- N: From 1852-1966 whole plots received 48kg N as nitrate of soda. Between 1968-2002 whole plots were split to test 4 rates of N as "Nitro-chalk" (cumulative applications until 1973, on a cyclic system from 1974).
 N3: Basal N, 144kg as "Nitro-chalk" since 2003
 Si: Silicate of soda at 450kg (Note: S also refers to silicate of soda)
 (Si): Silicate of soda omitted since 1980
 P, (P), K, Mg, (Mg), (Na): as above

Results of the Classical and other Long-term Experiments 2017

Phosphorus Test plots

Treatments:

Since 2003 the remaining plots [ex-Castor meal (plots 14, 24, 34 & 44) and those testing combinations of NPK with and without Mg (plots 55, 56, 57 & 58)] have been used to study the effect of P residues on yield. Previous treatments have resulted in different levels of available P in the soil. Large dressings of K were applied to some plots to increase levels of exchangeable K in the soil such that K should not limit yield; plots 141 and 241 were sacrificed and used as discard areas so that the K application did not encroach on adjacent no K plots on the Silicate Test. Other plots received the normal rate of K. The level of exchangeable Mg in the soil is such that Mg should not limit yield; the need to apply Mg was reviewed for 2017.

Whole plots

Manure

Plot	Treatment since
	2003
142	N3K*
143	N3K*
144	N3K*
242	N3K*
243	N3K*
244	N3K*
341	N3K
342	N3K
343	N3K
344	N3K
441	N3K
442	N3K
443	N3K
444	N3K
551	N3K
552	N3K
561	N3K
562	N3K
571	N3K*
572	N3K*
581	N3K*
582	N3K*

N3: Basal N, 144kg as "Nitro-chalk"

K: 90kg K as sulphate of potash

K*: 450kg K as sulphate of potash

In 2005 the extra dressings of K (i.e. K*) was stopped and all of the P test plots reverted to K

Results of the Classical and other Long-term Experiments 2017

Experimental Diary

Date		Application	Rate	Units
11/10/2016	a	Ploughed - thrown South	-	-
15/10/2016	p	Sprayed Buffalo Elite	1	lt/ha
15/10/2016	p	Sprayed Samurai	3	lt/ha
31/10/2016	f	Applied SOP to Plots 141 - 144, 241 - 244, 311 - 344, 411 - 444, 551 - 582, 631 - 634	217	kg/ha
31/10/2016	f	Applied Kieserite to Plots 631 - 634	233	kg/ha
31/10/2016	f	Applied TSP to Plots 631 - 634	215	kg/ha
08/11/2016	f	Applied Silicate of Soda to Plots 433, 333, 233, 133, 432, 332, 232 and 132	450	kg/ha
10/11/2016	f	Applied FYM to Plots 721 to 734	35	t/ha
07/03/2017	p	Sprayed Buffalo Elite	1	lt/ha
07/03/2017	p	Sprayed Samurai	3	lt/ha
12/03/2017	a	Cousin Combi Harrowed	-	-
15/03/2017	s	Drilled KWS Irina trt With Raxil Star	350	seeds/m ²
17/03/2017	a	Ring Rolled	-	-
12/05/2017	a	Rotavated Paths	-	-
22/05/2017	p	Sprayed Hallmark	50	ml/ha
23/05/2017	f	Applied Nitrochalk (27% N) to Plots 113, 124, 211, 222, 313, 321, 412, 421, 611, 621, 631, 712, 721, 732	48	kg/ha
23/05/2017	f	Applied Nitrochalk (27% N) to Plots 112, 123, 212, 223, 314, 324, 414, 422, 613, 624, 634, 711, 722, 731	96	kg/ha
23/05/2017	f	Applied Nitrochalk (27% N) to Plots 114, 122, 213, 224, 312, 323, 411, 424, 612, 622, 632, 714, 723, 733	144	kg/ha
24/05/2017	a	Rotavated Paths	-	-
24/05/2017	f	Applied Nitram to Series AA (except 621 to 624, 721 to 724), C and Strip 5	417	kg/ha
30/05/2017	p	Sprayed Refine Max	75	g/ha
30/05/2017	p	Sprayed Kingdom	1.25	lt/ha
30/05/2017	p	Sprayed Hurler	700	ml/ha
14/07/2017	a	Cut Paths	-	-
28/07/2017	a	Pulled 7 Wild Oats from within plots	-	-
29/08/2017	a	Harvested plots for yield & baled all discard and remaining swaths.	-	-

Results of the Classical and other Long-term Experiments 2017

Yields

Main Plots

Grain Yield, tonnes/hectare

Table of means

	N	0	48	96	144	Mean
MANURE						
---		1.43	1.13	0.95	1.14	1.16
-P-		2.50	3.94	3.78	3.92	3.53
--K		1.45	1.45	1.76	1.49	1.54
-PK		2.41	3.47	3.80	4.81	3.62
A--		0.95	1.36	1.10	1.08	1.12
AP-		2.86	3.94	4.44	4.13	3.84
A-K		0.95	0.97	1.26	1.12	1.07
APK		2.51	3.56	4.01	4.88	3.74
FYM1852onwards		8.04	8.07	7.92	7.91	7.99
FYM1852-1871		2.87	2.99	3.11	5.61	3.64
(A)		1.66	2.03	3.38	1.98	2.26
-		1.27	1.40	1.52	1.41	1.40
FYM2001onwards		6.66	6.92	7.09	6.93	6.90
P2K		3.62	4.78	4.86	5.63	4.72
Mean		2.80	3.28	3.5	3.72	3.33
Grain mean DM%		85.9				

Straw Yield, tonnes/hectare

Table of means

	N	0	48	96	144	Mean
MANURE						
---		0.32	0.25	0.19	0.17	0.23
-P-		0.38	0.69	0.72	0.79	0.65
--K		0.25	0.25	0.29	0.28	0.27
-PK		0.39	0.73	1.00	1.30	0.86
A--		0.20	0.22	0.24	0.21	0.22
AP-		0.61	0.84	1.05	0.95	0.86
A-K		0.16	0.18	0.19	0.20	0.18
APK		0.65	0.82	1.00	1.46	0.98

Results of the Classical and other Long-term Experiments 2017

FYM1852onwards	3.86	3.35	3.81	3.16	3.54
FYM1852-1871	0.64	0.70	0.63	1.73	0.92
(A)	0.26	0.41	0.61	0.13	0.35
-	0.32	0.27	0.20	0.36	0.29
FYM2001onwards	3.30	2.98	2.90	2.69	2.97
P2K	1.21	1.52	1.42	1.76	1.48
Mean	0.90	0.94	1.02	1.08	0.99

Grain mean DM% 85.8

PHOSPHATE PLOTS

Grain Yield, tonnes/hectare

Tables of means

PLOTS

142	3.66
143	3.57
144	3.27
242	5.13
243	4.82
244	4.84
341	3.50
342	4.12
343	4.23
344	4.68
441	4.55
442	5.23
443	5.04
444	5.08
551	5.24
552	4.81
561	4.77
562	4.46
571	4.19
572	4.69
581	0.99
582	1.08
Mean	4.18
Grain Mean DM%	86.7
Plot area Harvested	0.00244

Results of the Classical and other Long-term Experiments 2017

SILICATE PLOTS

Grain Yield, tonnes/hectare

Tables of means

	PK	N3--	N3P-	N3-K	N3PK	Mean
Silicate						
(-)-		1.34	4.26	1.24	4.99	2.96
(Si)-		1.89	4.99	2.63	5.11	3.65
(-)Si		2.54	5.11	2.34	5.14	3.78
(Si)Si		2.81	5.00	3.06	4.81	3.92
Mean		2.15	4.84	2.32	5.01	3.58
Grain Mean DM%		86.3				
Plot area harvested		0.00244				