

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Results of the Classical and Other Long-term Experiments - 2017



Results of the
Classical and other
Long-term Experiments
2017

[Full Table of Content](#)

17/R/EX/4 Exhaustion Land (Hoosfield)

Rothamsted Research

Rothamsted Research (2018) *17/R/EX/4 Exhaustion Land (Hoosfield)* ; Results Of The Classical And Other Long-Term Experiments - 2017, pp 20 - 23 - DOI: <https://doi.org/10.23637/ERADOC-1-251>

Results of the Classical and other Long-term Experiments 2017

17/R/EX/4 EXHAUSTION LAND (Hoosfield)

Object: To study the residual effects of manures applied 1856 - 1901, and of additional phosphate applied since 1986 (P test) and of additional potassium since 2007 (K test); on the yield of continuous spring barley up to 1991, winter wheat since – Hoosfield.

The 162nd year, winter wheat.

For previous years see 'Details' 1977, 1973 and Yield Books for 74-16/R/EX/4

Treatments: All combinations of:

Whole plots (P test)

1. **OLD RES** Residues of manures applied annually 1876 – 1901:

Main plot

- | | | |
|----|---------|---|
| 01 | O | None |
| 03 | D | Farmyard manure at 35 t |
| 05 | N | 96 kg N as ammonium salts |
| 09 | P | 34 kg P as superphosphate |
| 07 | NPKNaMg | N and P as above plus 137 kg K as sulphate of potash,
16 kg Na as sulphate of soda, 11 kg Mg as sulphate of magnesia |

2. P Maintenance P (20 kg P) applied annually from 2000

to maintain existing levels of available P in the soil. In 2009 maintenance P applications were changed from 20 kg P/ha to 15 kg P/ha. This was not recorded in the yield books for 2009-13. (P1) (P2) and (P3) are residues of P applied annually. From 2016 onward P was withheld from the P(P1) sub-plots.

1986–1992:

	2016-Present	2009-2015	2000-08	1986-92
O	None	None	None	None
P (P1)	None	15 kg P	20 kg P	44 kg P
P (P2)	15 kg P	15 kg P	20 kg P	87 kg P
P (P3)	15 kg P	15 kg P	20 kg P	131 kg P

NOTE: P treatments were applied at 61.5 kg P in error in 2000.

Plus

Whole plots (K test, previously N test until 1991)

1. **OLD RES** Residues of manures applied annually 1876 – 1901:

Main Plot

- | | | |
|----|------|---|
| 02 | O | None |
| 04 | D | Farmyard manure at 35 t |
| 06 | N* | 96 kg N as nitrate of soda |
| 10 | PK | 34 kg P as superphosphate, 137 kg K as sulphate of potash |
| 08 | N*PK | N, P and K as above |

Results of the Classical and other Long-term Experiments 2017

2.	K	Potassium applied annually from 2007 as muriate of potash
	O	None (2 sub-plots within each treatment strip)
	K1	75 kg K ₂ O (62.2 kg K)
	K2	150 kg K ₂ O (124.5 kg K)

Whole plots

Nitrogen: 50 kg N as ammonium sulphate (to supply sufficient S) during first two weeks in March, 200 kg N as ammonium nitrate at GS31/mid-April (whichever comes first) and 50 kg N as ammonium nitrate at GS37 (not later than mid-May).

Experimental Diary

Date		Application	Rate	Units
26/09/2016	a	Topped all stubble	-	-
26/09/2016	f	Applied TSP to plots 101-104, 081-084, 061-064, 041-044, 021-024, 091-092, 071-072, 051-052, 031-032 and 011-012	75	kg/ha
26/09/2016	f	Applied MOP to plots 103, 083, 063,043, 023	125	kg/ha
26/09/2016	f	Applied MOP onto plots 104-094, 084-074, 064-054, 044-034 and 024-014	250	kg/ha
28/09/2016	a	Ploughed - thrown North	-	-
06/10/2016	a	Cultipressed - all ground and immediate surrounds	-	-
10/10/2016	a	Ring Rolled All New Drilling	-	-
10/10/2016	s	Drilled Crusoe Treated w/Redigo Pro + Deter	350	seeds/m ²
10/10/2016	a	Cultipressed - all sites and surrounds	-	-
15/10/2016	p	Sprayed Liberator	600	ml/ha
15/10/2016	p	Sprayed Defy	3	lt/ha
15/10/2016	p	Sprayed Deploy	400	ml/ha
02/12/2016	p	Sprayed Hallmark	50	ml/ha
14/03/2017	f	Applied Sulphate of Ammonia (21%N 60%SO ₃) to all plots	238	kg/ha

Results of the Classical and other Long-term Experiments 2017

21/03/2017	f	Applied Kieserite	80	kg/ha
04/04/2017	f	Applied Nitram	580	kg/ha
05/04/2017	p	Sprayed Artemis	1	lt/ha
05/04/2017	p	Sprayed Claw 500	1	lt/ha
05/04/2017	p	Sprayed 3C Chlormequat 750	2	lt/ha
27/04/2017	p	Sprayed keystone	600	ml/ha
27/04/2017	p	Sprayed epic	400	ml/ha
27/04/2017	p	Sprayed balear 720sc	700	ml/ha
05/05/2017	f	Applied Nitram (34.5% N)	145	kg/ha
08/06/2017	a	Cut Paths	-	-
19/06/2017	p	Sprayed Fezan (Tebuconazole)	750	ml/ha
26/06/2017	a	cut all paths	-	-
27/07/2017	a	cut all paths	-	-
01/09/2017	a	Harvested All Plots	-	-
02/09/2017	a	Completed Straw Samples and Weights	-	-
06/09/2017	a	harvested leftover wheat of harvested trials and surrounds	-	-
07/09/2017	a	Baled all remaining commercial swath	-	-

Yields

P TEST

Grain Yield, tonnes/hectare

Tables of means

P_RES OLD_RES	O	(P1)	(P2)	(P3)	Mean
O	1.87	4.18	5.69	6.14	4.47
D	3.49	6.23	7.49	7.68	6.22
N	1.76	5.19	6.56	7.74	5.31
P	2.44	5.66	7.28	7.31	5.67
NPKNAMG	3.22	5.86	6.47	7.12	5.67
Mean	2.56	5.43	6.70	7.20	5.47
Grain mean DM%	86.4				

Results of the Classical and other Long-term Experiments 2017

Straw Yield, tonnes/hectare

Tables of means

P_RES	O	(P1)	(P2)	(P3)	Mean
OLD_RES					
O	1.01	1.76	2.65	3.05	2.12
D	1.78	2.86	3.58	3.36	2.89
N	1.23	2.81	3.66	2.78	2.62
P	0.93	2.15	2.68	3.02	2.20
NPKNAMG	1.40	2.37	2.69	3.16	2.40
Mean	1.27	2.39	3.05	3.07	2.45

Straw mean DM% 96.2

Plot area harvested 0.00512.

K TEST

Grain Yield, tonnes/hectare

Tables of means

K_Test	K0	K1	K2	Mean
OLD_RES				
O	6.54	7.80	7.52	7.10
D	7.02	8.52	8.07	7.66
N*	7.04	7.96	8.19	7.56
PK	8.37	7.86	8.23	8.21
N*PK	7.81	7.48	8.14	7.81
Mean	7.36	7.92	8.03	7.67
Grain mean DM%	86.6			

Straw Yield, tonnes/hectare

Tables of means

K_Test	K0	K1	K2	Mean
OLD_RES				
O	2.31	3.65	3.68	2.99
D	2.62	3.55	3.51	3.08
N*	2.65	3.73	3.77	3.2
PK	3.20	3.15	3.30	3.21
N*PK	2.99	1.99	2.95	2.73
Mean	2.76	3.21	3.44	3.04

Straw mean DM% 97.0

Plot area harvested 0.00512