Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Results of the Classical and Other Long-term Experiments - 2017



Full Table of Content

17/R/PG/5 Park Grass

Rothamsted Research

Rothamsted Research (2018) *17/R/PG/5 Park Grass ;* Results Of The Classical And Other Long-Term Experiments - 2017, pp 24 - 29 - **DOI: https://doi.org/10.23637/ERADOC-1-251**

17/R/PG/5 PARK GRASS

Object: To study the effects of organic manures and inorganic fertilisers and lime on old grass for hay.

The 162nd year, hay. For previous years see 'Details' 1977 and 1973 and Yield Books for 74-16/R/PG/5.

Treatments: Combinations of:

Whole plots

1.	Manure	Fert	Fertilizers and organic manures:				
	N1	Plo	ot 1	N1			
	К	Plo	ot 2/1	K since 1996 (as 2/2 before)			
	None (FYM)	Plo	ot 2/2	None (FYM until 1863)			
	None	Plo	ot 3	None			
	Р	Plo	ot 4/1	Р			
	N2P	Plo	ot 4/2	N2 P			
	N1PKNaMg	Plo	ot 6	N1 P K Na Mg			
	(P)KNaMg	Plo	ot 7/1	K Na Mg (+P until 2012)			
	PKNaMg	Plo	ot 7/2	P K Na Mg			
	PNaMg	Plo	ot 8	P Na Mg			
	PKNaMg(N2)	Plo	ot 9/1	P K Na Mg (+ N2 until 1989)			
	N2PKNaMg	Plo	ot 9/2	N2 P K Na Mg			
	N2PNaMg	Plo	ot 10	N2 P Na Mg			
	N3PKNaMg	Plo	ot 11/1	N3 P K Na Mg			
	N3PKNaMgSi	Plo	ot 11/2	N3 P K Na Mg Si			
	None	Plo	ot 12	None			
	(FYM/F)	Plo	ot 13/1	None (FYM/F until 1993/1995)			
	FYM/PM	Plo	ot 13/2	FYM/PM (FYM/F until 1999)			
	PKNaMg (N2*)	Plo	ot 14/1	P K Na Mg (+ N2* until 1989)			
	N2*PKNaMg	Plo	ot 14/2	N2* P K Na Mg			
	N3*PKNaMg (N2	2*) Plo	ot 15	N3*P K Na Mg (N2* until 1875; P K Na Mg			
				1876-2012)			
	N1*PKNaMg	Plo	ot 16	N1* P K Na Mg			
	N1*	Plo	ot 17	N1*			
	N2KNaMg	Plo	ot 18	N2 K Na Mg			
	FYM	Plo	ot 19	FYM			
	FYM/N*PK	Plo	ot 20	FYM/N*P K			
	N1, N2, N3:	48, 96, 144	96, 144 kg N as sulphate of ammonia				
	N1*, N2*,		8, 96, 144 kg N as nitrate of soda (30 kg N to plot 20 in				
	N3*:		ears with no farmyard manure). In 2013 plot 15				
		started to	tarted to receive 144 kg N/ha as nitrate of soda to				
		provide a	provide a comparison with plot 11/1, which receives				
		144 kg N/ł	na as sulphate of	ammonia.			

Ρ:	17 kg P/ha applied as triple superphosphate since 2017, except for plot 20 which receives 15 kg P/ha in years with no farmyard manure. Prior to this, 35 kg P (15 kg P to plot 20 in years with no farmyard manure) was applied as triple superphosphate in 1974 and since 1987, single superphosphate in other years.
(P):	In 2013 plot 7 was split into 7/1 & 7/2. P was withheld from plot 7/1 but 7/2 continues to receive P as above.
К:	225 kg K (45 kg K to plot 20 in years with no farmyard manure) as sulphate of potash
Na:	15 kg Na as sulphate of soda
Mg:	10 kg Mg as sulphate of magnesia
Si:	Silicate of soda at 450 kg
FYM:	Farmyard manure at 35 t every fourth year
F:	Fishmeal every fourth year to supply 63 kg N (stopped 1999; replaced by PM)
PM	Pelleted poultry manure at 2 t, every fourth year to supply 63 kg N (started 2003)

Sub-plots

2.	Lime	Liming plots 1-18 (excluding 18/2):
	а	Ground chalk applied as necessary to achieve pH7
	b	Ground chalk applied as necessary to achieve pH6
	С	Ground chalk applied as necessary to achieve pH5
	d	None

NOTE: A small amount of chalk was applied to all plots during tests in the 1880s and 1890s. A regular test of liming was started in 1903 when most plots were divided in two and 4 t ha⁻¹ CaCO₃ was applied every four years to the southern half. In 1965, most plots were divided into four: sub-plots "a" and "b" on the previously limed halves and subplots "c" and "d" on the unlimed halves. Sub-plots "a", "b" and "c" now receive different amounts of chalk, when necessary, to achieve and/or maintain soil (0-23cm) at pH 7, 6 and 5, respectively. Sub-plot "d" receives no lime and its pH reflects inputs from the various treatments and the atmosphere. Lime was last applied in 2014-2015; the eighth application in a triennial scheme of soil pH analysis and remedial chalk applications.

[This note was incorrect in earlier Yield book entries.]

NOTE: A separate scheme of liming was introduced on plots 18, 19 & 20 in 1920; subplot /1, /2 and /3 receive no lime, "high" lime and "light" lime respectively every 4 years. Since 1965 plot 18-1 has been split into two for treatments 'c' and 'd' as above and plot 18-3 split into two for treatments 'a' and 'b. Plots 19 and 20 received no further chalk after 1968; plot 18/2 no further chalk after 1972.

[This note was incorrect in earlier Yield book entries. See further details on the e-RA website at http://www.era.rothamsted.ac.uk]

Experimental Diary

Date		Application	Rate	Units
17/10/2016	а	Cut Paths - in and around trial	-	-
20/10/2016	а	Mowed All Grass	-	-
20/10/2016	а	Rowed up All Grass	-	-
25/11/2016	f	Applied TSP - plots 4/1, 4/2, 6, 7/2, 8, 9/1, 9/2, 10, 11/1, 11/2, 14/1, 14/2, 15, 16	73	kg/ha
28/11/2016	f	Applied Sulphate of Potash - plots 2/1, 6, 7/1, 7/2, 9/1, 9/2, 11/1, 11/2, 14/1, 14/2, 15, 16, 18	542	kg/ha
28/11/2016	f	Applied Sulphate of Soda - plots 6, 7/1, 7/2, 8, 9/1, 9/2, 10, 11/1, 11/2, 14/1, 14/2, 15, 16, 18	43	kg/ha
28/11/2016	f	Applied Sulphate of Magnesia (Epsom Salts) - plots 6, 7/1, 7/2, 8, 9/1, 9/2, 10, 11/1, 11/2, 14/1, 14/2, 15, 16, 18	111	kg/ha
28/11/2016	f	Applied Silicate of Soda - plot 11/2	450	kg/ha
30/11/2016	f	Applied FYM - plots 13/2, 19, 20	35	t/ha
18/04/2017	f	Applied Sulphate of Ammonia (21% N) - plot 1, 6a, 6b	229	kg/ha
18/04/2017	f	Applied Sulphate of Ammonia (21% N) - plots 4/2, 9/2, 10, 18	457	kg/ha
18/04/2017	f	Applied Sulphate of Ammonia (21% N) - plots 11/1, 11/2	686	kg/ha
19/04/2017	f	Applied Sodium Nitrate (16% N) - plots 16, 17	300	kg/ha
19/04/2017	f	Applied Sodium Nitrate (16% N) - plot 14/2	600	kg/ha
19/04/2017	f	Applied Sodium Nitrate (16% N) - plot 15	900	kg/ha
10/05/2017	а	Cut All Paths	-	-
23/05/2017	а	Cut paths	-	-
07/06/2017	а	Cut Paths	-	-
19/06/2017	а	Cut Paths and surrounds	-	-
20/06/2017	а	Started harvesting grass yields - 1st Cut	-	-

21/06/2017	а	Completed grass yield - 1st Cut	-		-
21/06/2017	а	Mowed all grass	-		-
23/06/2017	а	Turned all hay	-		-
26/06/2017	а	Rowed up all grass for baling	-		-
19/10/2017	а	Cut Paths	-		-
24/10/2017	а	Started harvesting plot yields - 2nd Cut	-		-
25/10/2017	а	Completed harvesting yield plots - 2nd Cut	-	-	
01/11/2017	а	Baled leftover grass and removed	-	-	

NOTE: Samples of herbage (1st and 2nd Cut) were taken for chemical analysis. Unground herbage samples from all plots were archived.

Yields

1ST CUT (20-21 JUN 2017) DRY MATTER, TONNES/HECTARE

Tables of means

Grand mean 3.00 Lime b d Mean Manure С а N1 1 1.42 1.10 0.83 0.31 0.91 K 2/1 1.44 0.85 0.52 0.91 0.85 None(FYM) 2/2 1.72 0.93 0.88 1.29 1.63 None 3 1.44 1.64 0.83 0.82 1.18 P 4/1 2.34 2.69 1.91 1.51 2.11 N2P 4/2 1.87 2.53 2.92 1.28 2.15 N1PKNaMg 5.23 5.04 5.14 6 _ (P)KNaMg 7/1 3.84 4.28 3.88 1.78 3.44 PKNaMg 7/2 4.54 5.28 4.90 2.76 4.37 PNaMg 8 2.13 2.62 2.13 2.51 2.35 PKNaMg(N2) 9/1 3.96 4.46 3.97 0.37 3.19 N2PKNaMg 9/2 5.21 5.53 4.11 1.91 4.19 N2PNaMg 10 2.40 2.86 3.09 1.27 2.40 N3PKNaMg 11/1 4.75 4.15 4.44 2.32 3.91 4.49 4.04 N3PKNaMgSi 11/2 4.96 4.21 2.49 None 12 1.71 1.44 1.06 1.03 1.31 (FYM/F) 13/1 2.16 2.45 2.26 1.88 2.19 FYM/PM 13/2 4.41 3.81 4.71 4.50 4.63 PKNaMg(N2*) 14/1 5.20 6.25 5.29 5.32 5.52 N2*PKNaMg 14/2 4.58 4.42 4.08 4.46 4.55 N3*PKNaMg(N2*) 15 4.78 4.45 5.45 3.82 3.74 N1*PKNaMg 16 4.91 5.35 3.90 3.74 4.48 N1* 17 0.87 1.64 1.24 1.38 1.77 N2KNaMg 18 1.33 1.97 1.69 0.08 1.27

N2KNaMg 18/2	2.81
FYM 19/1	4.32
FYM 19/2	4.75
FYM 19/3	4.29
FYM/N*PK 20/1	4.63
FYM/N*PK 20/2	4.48
FYM/N*PK 20/3	4.40

1st cut mean DM% 30.8

2ND CUT (24-25 OCT 2017) DRY MATTER, TONNES/HECTARE

Tables of means								
Grand mean	2.01							
Manure	Lime	а	b	С	d	Mean		
N1 1		1.67	1.56	1.74	0.61	1.39		
К 2/1		1.41	1.28	0.99	0.94	1.15		
None(FYM) 2/2		1.24	1.23	1.47	1.32	1.32		
None 3		1.03	1.27	1.40	1.31	1.25		
P 4/1		1.85	1.86	2.02	2.10	1.96		
N2P 4/2		1.51	1.92	1.98	1.24	1.66		
N1PKNaMg 6		2.08	2.23	-	-	2.15		
(P)KNaMg 7/1		2.05	2.40	2.01	1.37	1.96		
PKNaMg 7/2		2.01	2.43	2.47	1.60	2.13		
PNaMg 8		1.79	1.83	2.07	2.89	2.15		
PKNaMg(N2) 9/1		2.27	2.49	2.18	0.43	1.84		
N2PKNaMg 9/2		2.56	2.52	2.03	1.55	2.17		
N2PNaMg 10		1.74	1.86	2.61	1.12	1.83		
N3PKNaMg 11/1		2.00	2.23	2.39	2.23	2.21		
N3PKNaMgSi 11/2		2.90	2.48	2.15	1.85	2.35		
None 12		1.96	1.63	1.20	1.20	1.50		
(FYM/F) 13/1		2.48	2.69	2.10	1.73	2.25		
FYM/PM 13/2		2.32	3.11	2.72	2.80	2.74		
PKNaMg(N2*) 14/1		2.14	2.92	3.22	3.11	2.85		
N2*PKNaMg 14/2		1.82	2.37	2.62	2.71	2.38		
N3*PKNaMg(N2*) 15		2.26	2.31	2.83	2.65	2.51		
N1*PKNaMg 16		2.37	2.90	2.64	2.24	2.54		
N1* 17		1.59	1.82	1.64	1.91	1.74		
N2KNaMg 18		1.48	1.34	1.27	0.46	1.14		
N2KNaMg 18/2						1.68		
FYM 19/1						3.04		
FYM 19/2						3.22		
FYM 19/3						2.90		
FYM/N*PK 20/1						2.85		
FYM/N*PK 20/2						2.94		
FYM/N*PK 20/3						2.92		

2nd cut mean DM% 25.13

TOTAL OF 2 CUTS DRY MATTER, TONNES/HECTARE

Tables of means

Grand mean 5.00

Ma	nure	Lime	а	b	с	d	Mean
N1	1		3.09	2.65	2.56	0.92	2.31
К	2/1		2.25	2.73	1.84	1.46	2.07
None(FYM)	2/2		2.88	2.95	2.40	2.20	2.61
None	, 3		2.48	2.91	2.23	2.13	2.44
Р	4/1		4.18	4.55	3.93	3.61	4.07
N2P	4/2		3.38	4.45	4.91	2.52	3.81
N1PKNaMg	6		7.31	7.27	-	-	7.29
(P)KNaMg	7/1		5.89	6.68	5.88	3.15	5.40
PKNaMg	7/2		6.55	7.72	7.37	4.36	6.50
PNaMg	8		3.92	4.45	4.20	5.40	4.49
PKNaMg(N2)	9/1		6.73	6.45	6.15	0.80	5.03
N2PKNaMg	9/2		7.77	8.05	6.14	3.47	6.36
N2PNaMg	10		4.14	4.73	5.70	2.38	4.24
N3PKNaMg	11/1		6.74	6.38	6.83	4.55	6.13
N3PKNaMgSi	11/2		7.86	6.96	6.36	4.34	6.38
None	12		3.67	3.07	2.26	2.22	2.81
(FYM/F)	13/1		4.64	5.14	4.36	3.61	4.44
FYM/PM	13/2		6.13	7.82	7.22	7.43	7.15
PKNaMg(N2*)	14/1		7.34	9.17	8.51	8.44	8.36
N2*PKNaMg	14/2		5.90	6.95	7.08	7.26	6.80
N3*PKNaMg(N2*)	15		7.71	7.09	6.65	6.38	6.96
N1*PKNaMg	16		7.28	8.25	6.54	5.99	7.01
N1*	17		2.46	3.46	2.88	3.68	3.12
N2KNaMg			2.81	3.31	2.96	0.54	2.41
N2KNaMg							4.49
FYM							7.36
FYM							7.97
FYM							7.19
FYM/N*PK							7.48
FYM/N*PK	•						7.42
FYM/N*PK	20/3						7.33
TOTAL OF 2 (
TOTAL OF Z							

Mean DM% 27.98