Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments



Full Table of Content

19/R/HB/2 - Hoos Barley (Hoosfield)

Rothamsted Research

Rothamsted Research (2021) 19/R/HB/2 - Hoos Barley (Hoosfield); Yields Of The Field Experiments, pp 12 - 18

19/R/HB/2

19/R/HB/2 HOOS BARLEY (Hoosfield)

Object: To study the effects of organic manures and inorganic fertilizers on continuous spring barley. From 1968 to 1978 a rotation of potatoes, beans and spring barley was practised on parts of the experiment. The rotation was discontinued in 1979 and the whole experiment reverted to continuous spring barley. The experiment was modified for 2003. The main plots continue as previously. The Silicate Test plots continue but are not split to test rates of N (basal N is applied). The remaining plots are to be used to study the effect on yield of P residues, (basal N applied).

The 168th year, spring barley.

For previous years see 'Details' 1967 and 1973, Station Report for 1966 and Yield Books for 74-18/R/HB/2.

Main plots

Treatments:

Whole plots

| MANURE | | Fertilizers and Organic Manures:- | | | | | |
|-----------|-------------------|-----------------------------------|-------------|------------|--|--|--|
| | | | Additional | | | | |
| | | Form of N | treatments | Treatments | | | |
| | Plot | 1852-1966 | 1852-2002 | since 2003 | | | |
| | 11 | None | 32 2 | - | | | |
| -P- | 21 | None | Р | (P) | | | |
| K | 31 | None | K (Na) Mg | K(Mg) | | | |
| -PK | 41 | None | PK (Na) Mg | (P) K (Mg) | | | |
| A | 12 | Α | | ¥ | | | |
| AP- | 22 | Α | Р | (P) | | | |
| A-K | 32 | Α | K (Na) Mg | K(Mg) | | | |
| APK | 42 | Α | PK (Na) Mg | (P) K (Mg) | | | |
| D1852 | 72 | None | D | D | | | |
| (D) | 71 | None | (D) | (D) | | | |
| (A) | 62 | None | (Ashes) | (Ashes) | | | |
| | 61 | None | 15 | ā | | | |
| D2001 (a) | 73 ^(a) | 22 | D | D | | | |
| P2KMg (a) | 63 ^(a) | | P2KMg | P2KMg | | | |

⁽a) Plots 63 and 73 started in 2001

Form of N: A, sulphate of ammonia to supply 48kg N

- P: 35 kg P as triple superphosphate in 1974 and from 1988 to 2002, single superphosphate in other years
- (P): (none), P application to be reviewed for 2018
- P2: 44kg P as triple superphosphate
- K: 90 kg K as sulphate of potash
- (Na): (none), 16 kg Na as sulphate of soda until 1973
- Mg: 35kg Mg as kieserite every third year since 1974 (applied at 30 kg in 1992, 1995 and 1998) (sulphate of magnesia annually until 1973). Annually to new plot 63.
- (Mg): (none), Mg application to be reviewed for 2021

19/R/HB/2

D1852: Farmyard manure at 35t since 1852
D2001: Farmyard manure at 35t since 2001
(D): Farmyard manure 1852 – 1871 only

(Ashes): Weed ash 1852-1916, furnace ash 1917-1932, none since

Sub-plots

(2) N Nitrogen fertilizer (kg N), as 'Nitro-Chalk', since 1968 (cumulative N applications until 1973, on a cyclic system since 1974):

0
48
96
144

Silicate Test plots

Treatments:

Whole plots

| MANURE | Plot | Fertilize | ers:- | | |
|--------|------|-----------|-------|--------------------|-----------------------|
| | | Addition | nal | Changes since 1980 | Treatments since 2003 |
| | | treatme | ent | | |
| | | 1852-19 | 979 | | |
| N | 131 | - | | - | N3 |
| NP | 231 | Р | | | N3 (P) |
| N-K | 331 | K(Na)I | Mg | - | N3 K(Mg) |
| NPK | 431 | PK(Na)N | Иg | | N3(P)K(Mg) |
| N-S- | 134 | | Si | Si omitted | N3 (Si) |
| NP-S- | 234 | Р | Si | Si omitted | N3(P) (Si) |
| N-KS- | 334 | K(Na)I | MgSi | Si omitted | N3 K(Mg)(Si) |
| NPKS- | 434 | PK(Na)N | ∕lgSi | Si omitted | N3(P)K(Mg)(Si) |
| NS | 132 | *** | | Si added | N3 Si |
| NPS | 232 | Р | | Si added | N3(P) Si |
| N-K-S | 332 | K(Na)I | Mg | Si added | N3 K(Mg) Si |
| NPK-S | 432 | PK(Na)N | Иg | Si added | N3(P)K(Mg) Si |
| NSS | 133 | | Si | ≅. | N3 Si |
| NP-SS | 233 | Р | Si | | N3(P) Si |
| N-KSS | 333 | K(Na)I | MgSi | ¥ | N3 K(Mg) Si |
| NPKSS | 433 | PK(Na)N | ИgSi | | N3(P)K(Mg) Si |
| | | | | | |

N: From 1852-1966 whole plots received 48kg N as nitrate of soda. Between 1968-2002 whole plots were split to test 4 rates of N as "Nitro-chalk" (cumulative applications until 1973, on a cyclic system from 1974).

N3: Basal N, 144kg as "Nitro-chalk" since 2003

Si: Silicate of soda at 450kg (Note: S also refers to silicate of soda)

⁽Si): Silicate of soda omitted since 1980

P, (P), K, Mg, (Mg), (Na): as above

19/R/HB/2

Phosphorus Test plots

Treatments:

Since 2003 the remaining plots [ex-Castor meal (plots 14, 24, 34 & 44) and those testing combinations of NPK with and without Mg (plots 55, 56, 57 & 58)] have been used to study the effect of P residues on yield. Previous treatments have resulted in different levels of available P in the soil. Large dressings of K were applied to some plots to increase levels of exchangeable K in the soil such that K should not limit yield; plots 141 and 241 were sacrificed and used as discard areas so that the K application did not encroach on adjacent no K plots on the Silicate Test. Other plots received the normal rate of K. The level of exchangeable Mg in the soil is such that Mg should not limit yield; the need to apply Mg was reviewed for 2019.

Whole plots

Manure

| Plot | Treatment since |
|------|-----------------|
| | 2003 |
| 142 | N3K* |
| 143 | N3K* |
| 144 | N3K* |
| 242 | N3K* |
| 243 | N3K* |
| 244 | N3K* |
| 341 | N3K |
| 342 | N3K |
| 343 | N3K |
| 344 | N3K |
| 441 | N3K |
| 442 | N3K |
| 443 | N3K |
| 444 | N3K |
| 551 | N3K |
| 552 | N3K |
| 561 | N3K |
| 562 | N3K |
| 571 | N3K* |
| 572 | N3K* |
| 581 | N3K* |
| 582 | N3K* |
| | |

N3: Basal N, 144kg as "Nitro-chalk"

K: 90kg K as sulphate of potash

K*: 450kg K as sulphate of potash

In 2005 the extra dressings of K (i.e. K^*) was stopped and all of the P test plots reverted to K

19/R/HB/2

Experimental Diary

| Date | | Application | Rate | Units |
|------------|---|--|----------------|-------------------|
| 06/09/2018 | f | Applied Chalk to plots 711-734, 411-551, 611-634, 311-561, 241-571, 141-581 | 4 | t/ha |
| 08/11/2018 | f | Applied TSP - to plots 631-634 | 215 | kg/ha |
| 08/11/2018 | f | Applied SOP - to plots 631-634, 411-551, 311-561, 241-571, 141-581 | 217 | kg/ha |
| 08/11/2018 | f | Applied Kieserite - to plots 631-636 | 233 | kg/ha |
| 12/11/2018 | р | Sprayed Buffalo Elite | 1 | lt/ha |
| 12/11/2018 | р | Sprayed Samurai | 4 | t/ha |
| 15/11/2018 | f | Applied Silicate of Soda - to plots 433-133, 432-132 | 450 | kg/ha |
| 16/11/2018 | f | Applied FYM - to plots 721-734 | 35 | t/ha |
| 20/11/2018 | а | Ploughed, thrown North | - | • |
| 21/02/2019 | а | Cousin cultivated trial and surrounds | 1 | ; = 7(|
| 22/02/2019 | а | Ring Rolled All New Drilling | 18 | |
| 22/02/2019 | S | Drilled KWS Irina,trt Redigo Pro | 350 | seed/m² |
| 11/04/2019 | а | Rotavated Paths | 8.73 | 5 0 |
| 18/04/2019 | f | Applied N as Nitro-Chalk by hand - to plots 114,122, 213, 224, 312, 323, 411, 424, 612, 622, 632, 714, 723, 733 | 48 | kg/ha |
| 18/04/2019 | f | Applied N as Nitro-Chalk by hand - to plots 111, 121, 214, 221, 311, 322, 413, 423, 614, 623, 633, 713, 724, 734 | 96 | kg/ha |
| 18/04/2019 | f | Applied N as Nitro-Chalk by hand - to plots 113, 124, 211, 222, 313, 321, 412, 421, 611, 621, 631, 712, 721, 732 | 144 | kg/ha |
| 30/04/2019 | р | Sprayed Hallmark | 50 | ml/ha |
| 15/05/2019 | р | Sprayed Refine Max | 75 | g/ha |
| 15/05/2019 | р | Sprayed Kingdom | 1.5 | lt/ha |
| 15/05/2019 | р | Sprayed Claw 500 | 1 | lt/ha |
| 15/05/2019 | р | Sprayed Starane Hi | 400 | ml/ha |
| 16/05/2019 | f | Applied Nitram - to old series 5, c, aa - not plots 6, 7 | 417 | kg/ha |
| 02/07/2019 | р | Sprayed Mobius | 400 | ml/ha |
| 15/07/2019 | а | Pulled Oats | 13 | plants |
| 06/09/2019 | а | Harvested All Plots | 1. | #0 |
| 09/09/2019 | а | Bale surplus straw from plots | := | 2 23 |
| 10/09/2019 | а | Straw Weights recorded in grams for all main plots | 15 | 康治 |

19/R/HB/2

Yields

Main Plots

Grain Yield, tonnes/hectare

Table of means

| N MANURE | 0 | 48 | 96 | 144 | Mean |
|----------------|-------|------|------|------|------|
| | 1.86 | 2.26 | 3.35 | 2.21 | 2.42 |
| -P- | 2.31 | 3.33 | 4.23 | 4.94 | 3.70 |
| K | 3.08 | 4.00 | 4.62 | 3.88 | 3.90 |
| -PK | 2.99 | 5.19 | 6.15 | 5.94 | 5.07 |
| A | 1.70 | 2.34 | 2.42 | 2.86 | 2.33 |
| AP- | 2.71 | 4.10 | 4.32 | 4.23 | 3.84 |
| A-K | 2.78 | 3.36 | 3.78 | 3.62 | 3.38 |
| APK | 3.34 | 4.99 | 6.14 | 5.86 | 5.08 |
| FYM1852onwards | 8.35 | 9.15 | 8.54 | 8.76 | 8.70 |
| FYM1852-1871 | 2.95 | 5.78 | 5.81 | 5.06 | 4.90 |
| (A) | 3.37 | 3.58 | 4.18 | 3.82 | 3.74 |
| - | 2.45 | 3.54 | 4.04 | 3.78 | 3.45 |
| FYM2001onwards | 7.39 | 7.84 | 7.16 | 7.85 | 7.56 |
| P2K | 2.81 | 5.35 | 5.76 | 6.56 | 5.12 |
| | | | | | |
| Mean | 3.43 | 4.63 | 5.04 | 4.96 | 4.51 |
| Grain mean DM% | 87.00 | | | | |

19/R/HB/2

Table of means

| N MANURE | 0 | 48 | 96 | 144 | Mean |
|-------------------|------------------|---------|------|------|------|
| | 0.09 | 0.54 | 0.71 | 0.66 | 0.50 |
| -P- | 0.46 | 0.66 | 1.19 | 1.37 | 0.92 |
| K | 0.56 | 0.97 | 1.42 | 1.10 | 1.01 |
| -PK | 0.70 | 1.65 | 2.14 | 2.21 | 1.68 |
| A | 0.36 | 0.59 | 0.58 | 0.55 | 0.52 |
| AP- | 0.53 | 1.00 | 1.28 | 1.10 | 0.98 |
| A-K | 0.69 | 0.69 | 0.90 | 1.01 | 0.82 |
| APK | 0.84 | 1.86 | 2.13 | 1.60 | 1.61 |
| FYM1852onwards | 3.28 | 3.80 | 3.80 | 3.18 | 3.51 |
| FYM1852-1871 | 0.93 | 1.77 | 0.99 | 1.35 | 1.26 |
| (A) | 0.95 | 0.96 | 1.15 | 0.68 | 0.93 |
| 20 | 0.20 | 0.98 | 1.04 | 0.47 | 0.67 |
| FYM2001 onwards | 2.54 | 2.61 | 2.94 | 3.40 | 2.87 |
| P2K | 0.84 | 1.82 | 1.62 | 2.28 | 1.64 |
| | | | | | |
| Mean | 0.93 | 1.42 | 1.56 | 1.50 | 1.35 |
| 82-000 Z-00000000 | 221(2)(4.478.00) | | | | |
| Straw mean DM% | 84.70 | 0.00193 | | | |
| Plot Area | 0.00244 | 0.00183 | | | |

PHOSPHATE PLOTS

Grain Yield, tonnes/hectare

Tables of means

PLOTS

| 142 | 4.07 |
|-----|------|
| 143 | 4.43 |
| 144 | 3.81 |
| 242 | 4.56 |
| 243 | 4.62 |
| 244 | 4.85 |
| 341 | 4.33 |

19/R/HB/2

| 342 | 4.50 |
|----------------|------|
| 343 | 5.38 |
| 344 | 5.44 |
| 441 | 3.71 |
| 442 | 4.96 |
| 443 | 5.55 |
| 444 | 5.60 |
| 551 | 6.09 |
| 552 | 6.15 |
| 561 | 5.80 |
| 562 | 6.09 |
| 571 | 5.25 |
| 572 | 5.68 |
| 581 | 2.56 |
| 582 | 2.46 |
| Mean | 4.81 |
| Grain Mean DM% | 85.2 |
| | |

Plot area Harvested 0.00244

SILICATE PLOTS

Grain Yield, tonnes/hectare

Tables of means

| PK Silicate | N3 | N3P- | N3-K | N3PK | Mean |
|----------------|------|------|------|------|------|
| (-)- | 3.21 | 3.96 | 3.18 | 5.32 | 3.92 |
| Si)- | 3.57 | 4.41 | 4.78 | 4.93 | 4.42 |
| (-)Si | 4.36 | 4.87 | 4.89 | 5.38 | 4.88 |
| (Si)Si | 4.28 | 5.03 | 5.05 | 6.03 | 5.10 |
| Mean | 3.85 | 4.57 | 4.48 | 5.41 | 4.58 |

Grain Mean DM% 85.2

Plot area harvested 0.00244