

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



**ROTHAMSTED
RESEARCH**

Yields of the Field Experiments



[Full Table of Content](#)

19/R/PG/5 - Park Grass

Rothamsted Research

Rothamsted Research (2021) *19/R/PG/5 - Park Grass* ; Yields Of The Field Experiments , pp 24 - 30

19/R/PG/5 PARK GRASS

Object: To study the effects of organic manures and inorganic fertilisers and lime on old grass for hay.

The 164th year, hay.

For previous years see 'Details' 1977 and 1973 and Yield Books for 74-18/R/PG/5.

Treatments: Combinations of:

Whole plots

1.	Manure	Fertilizers and organic manures:
	N1	Plot 1
	K	Plot 2/1
	None (FYM)	Plot 2/2
	None	Plot 3
	P	Plot 4/1
	N2P	Plot 4/2
	N1PKNaMg	Plot 6
	(P)KNaMg	Plot 7/1
	PKNaMg	Plot 7/2
	PNaMg	Plot 8
	PKNaMg(N2)	Plot 9/1
	N2PKNaMg	Plot 9/2
	N2PNaMg	Plot 10
	N3PKNaMg	Plot 11/1
	N3PKNaMgSi	Plot 11/2
	None	Plot 12
	(FYM/F)	Plot 13/1
	FYM/PM	Plot 13/2
	PKNaMg (N2*)	Plot 14/1
	N2*PKNaMg	Plot 14/2
	N3*PKNaMg (N2*)	Plot 15
	N1*PKNaMg	Plot 16
	N1*	Plot 17
	N2KNaMg	Plot 18
	FYM	Plot 19
	FYM/N*PK	Plot 20
	N1, N2, N3:	48, 96, 144 kg N as sulphate of ammonia
	N1*, N2*,	48, 96, 144 kg N as nitrate of soda (30 kg N to plot 20 in
	N3*:	years with no farmyard manure). In 2013 plot 15
		started to receive 144 kg N/ha as nitrate of soda to
		provide a comparison with plot 11/1, which receives
		144 kg N/ha as sulphate of ammonia.

P:	17 kg P/ha applied as triple superphosphate since 2017, except for plot 20 which receives 15 kg P/ha in years with no farmyard manure. Prior to this, 35 kg P (15 kg P to plot 20 in years with no farmyard manure) was applied as triple superphosphate in 1974 and since 1987, single superphosphate in other years.
(P):	In 2013 plot 7 was split into 7/1 & 7/2. P was withheld from plot 7/1 but 7/2 continued to receive P as above.
K:	225 kg K (45 kg K to plot 20 in years with no farmyard manure) as sulphate of potash
Na:	15 kg Na as sulphate of soda
Mg:	10 kg Mg as sulphate of magnesia
Si:	Silicate of soda at 450 kg
FYM:	Farmyard manure at 35 t every fourth year
F:	Fishmeal every fourth year to supply 63 kg N (stopped 1999; replaced by PM)
PM	Pelleted poultry manure at 2 t, every fourth year to supply 63 kg N (started 2003)

Sub-plots

2.	Lime	Liming plots 1-18 (excluding 18/2):
	a	Ground chalk applied as necessary to achieve pH7
	b	Ground chalk applied as necessary to achieve pH6
	c	Ground chalk applied as necessary to achieve pH5
	d	None

NOTE: A small amount of chalk was applied to all plots during tests in the 1880s and 1890s. A regular test of liming was started in 1903 when most plots were divided in two and 4 t ha⁻¹ CaCO₃ was applied every four years to the southern half. In 1965, most plots were divided into four: sub-plots "a" and "b" on the previously limed halves and sub-plots "c" and "d" on the unlimed halves. Sub-plots "a", "b" and "c" now receive different amounts of chalk, when necessary, to achieve and/or maintain soil (0-23cm) at pH 7, 6 and 5, respectively. Sub-plot "d" receives no lime and its pH reflects inputs from the various treatments and the atmosphere. Lime was last applied in 2017-2018; the ninth application in a triennial scheme of soil pH analysis and remedial chalk applications.

[This note was incorrect in earlier Yield book entries.]

NOTE: A separate scheme of liming was introduced on plots 18, 19 & 20 in 1920; subplot /1, /2 and /3 receive no lime, "high" lime and "light" lime respectively every 4 years. Since 1965 plot 18-1 has been split into two for treatments 'c' and 'd' as above and plot 18-3 split into two for treatments 'a' and 'b'. Plots 19 and 20 received no further chalk after 1968; plot 18/2 no further chalk after 1972.

[This note was incorrect in earlier Yield book entries. See further details on the e-RA website at <http://www.era.rothamsted.ac.uk>]

Results of the Classical and other Long-term Experiments 2019

19/R/PG/5

Experimental Diary

Date	Application	Rate	Units
06/11/2018	f Applied TSP Treatments - plots 4/1,4/2, 6a, 6b, 7/2, 8, 9/1, 9/2, 10, 11/1, 11/2, 14/2, 14/1, 15, 16	83	kg/ha
06/11/2018	f Applied TSP Treatments - plot 20	73	kg/ha
28/01/2019	f Sulphate of Potash (50% K ₂ O) - plots 2/1, 6a, 6b, 7/1, 7/2, 9/1, 9/2, 11/1, 11/2, 14/2, 14/1, 15, 16, 18	542	kg/ha
28/01/2019	f Sulphate of Potash (50% K ₂ O) - plots 20	108	kg/ha
28/01/2019	f Sulphate of Soda (35% Na) - plots 6a, 6b, 7/1, 7/2, 8, 9/1, 9/2, 10, 11/1, 11/2, 14/2, 14/1, 15, 16, 18	43	kg/ha
28/01/2019	f Sulphate of Magnesia as Epson Salts (9% Mg) - plots 6a, 6b, 7/1, 7/2, 8, 9/1, 9/2, 10, 11/1, 11/2, 14/2, 14/1, 15, 16, 18	111.1	kg/ha
28/01/2019	f Silicate of Soda - plots 11/2	450	kg/ha
31/01/2019	f Poultry Manure - plot 13/2	2	t /ha
01/04/2019	a Topped paths	-	-
10/04/2019	f Applied Sulphate of Ammonia (21% N) - plots 1 and 6a, 6b	229	kg/ha
10/04/2019	f Applied Sulphate of Ammonia (21% N) - plots 4/2, 9/2, 10, 18	457	kg/ha
10/04/2019	f Applied Sulphate of Ammonia (21% N) - plots 11/1, 11/2	686	kg/ha
10/04/2019	f Applied Sodium Nitrate (16% N) - plot 20	188	kg/ha
10/04/2019	f Applied Sodium Nitrate (16% N) - plots 16, 17	300	kg/ha
10/04/2019	f Applied Sodium Nitrate (16% N) - plots 14/2	600	kg/ha
11/04/2019	f Applied Sodium Nitrate (16% N) - plots 15	900	kg/ha
11/04/2019	a Cut Paths	-	-
29/04/2019	a Cut Paths	-	-
15/05/2019	a Cut Paths	-	-
24/05/2019	a topped surrounds and paths	-	-
20/06/2019	a Cut Paths	-	-
26/06/2019	a Harvest - 1st Cut for grass yields	-	-
28/06/2019	a Test Cut plots for yield with Haldrup	-	-
01/07/2019	a Mowed all grass plots	-	-
02/07/2019	a turned grass plots	-	-
15/08/2019	a Path Cutting - Kilworth Topper - Izeki tractor	-	-
17/10/2019	a Path Cutting - Kilworth Topper - Izeki tractor	-	-
22/10/2019	a Harvested - 2nd Cut for grass yields - plot 18d second pass moved to south after accident with mower - still full length of plot	-	-
29/10/2019	a Row up	-	-
30/10/2019	a Baling	7	bales

NOTE: Samples of herbage (1st and 2nd Cut) were taken for chemical analysis. Unground herbage samples from all plots were archived.

Yields

1ST CUT (26-27 JUN 2019) DRY MATTER, TONNES/HECTARE

Tables of means

Grand mean		4.16					
Manure	Lime	a	b	c	d	Mean	
N1	1	2.66	2.49	2.03	1.24	2.10	
K	2/1	2.46	2.63	1.87	2.49	2.36	
None(FYM)	2/2	2.92	2.98	2.41	2.24	2.64	
None	3	2.50	2.90	2.51	2.02	2.48	
P	4/1	2.72	3.50	3.30	2.83	3.09	
N2P	4/2	4.26	4.83	4.81	2.62	4.13	
N1PKNaMg	6	5.26	5.81	-	-	5.54	
(P)KNaMg	7/1	4.10	5.47	4.97	2.69	4.31	
PKNaMg	7/2	3.92	5.49	5.05	4.20	4.66	
PNaMg	8	3.13	3.75	4.07	4.75	3.93	
PKNaMg(N2)	9/1	4.08	4.99	4.28	1.00	3.59	
N2PKNaMg	9/2	5.34	6.07	5.19	4.56	5.29	
N2PNaMg	10	4.31	4.60	5.08	3.31	4.32	
N3PKNaMg	11/1	5.27	5.78	5.66	4.91	5.41	
N3PKNaMgSi	11/2	6.86	6.52	5.66	5.67	6.18	
None	12	2.90	2.35	2.68	2.45	2.59	
(FYM/F)	13/1	4.00	4.34	4.17	3.75	4.06	
FYM/PM	13/2	4.01	5.25	5.22	6.05	5.13	
PKNaMg(N2*)	14/1	3.45	4.96	4.79	4.84	4.51	
N2*PKNaMg	14/2	5.32	6.06	4.85	6.03	5.56	
N3*PKNaMg(N2*)	15	5.45	6.48	5.60	6.15	5.92	
N1*PKNaMg	16	4.79	5.27	4.37	4.48	4.73	
N1*	17	3.03	3.58	2.71	3.28	3.15	

Results of the Classical and other Long-term Experiments 2019

19/R/PG/5

N2KNaMg 18	3.49	3.73	3.26	1.11	2.90
N2KNaMg 18/2	-	-	-	-	3.93
FYM 19/1	-	-	-	-	5.50
FYM 19/2	-	-	-	-	5.62
FYM 19/3	-	-	-	-	5.84
FYM/N*PK 20/1	-	-	-	-	5.75
FYM/N*PK 20/2	-	-	-	-	5.84
FYM/N*PK 20/3	-	-	-	-	5.39

1st cut mean DM% 26.80

2ND CUT (22 OCT 2019) DRY MATTER, TONNES/HECTARE

Tables of means

Grand mean 0.93

Manure	Lime	a	b	c	d	Mean
N1 1		0.90	0.83	0.91	0.42	0.77
K 2/1		0.70	0.60	0.62	0.81	0.68
None(FYM) 2/2		0.80	0.80	1.04	0.91	0.89
None 3		0.86	0.87	1.13	1.01	0.97
P 4/1		1.09	1.18	1.75	1.18	1.30
N2P 4/2		0.57	0.76	0.60	0.60	0.63
N1PKNaMg 6		0.64	0.63	-	-	0.63
(P)KNaMg 7/1		0.79	0.90	0.74	0.72	0.79
PKNaMg 7/2		0.67	0.76	0.82	0.55	0.70
PNaMg 8		0.73	0.57	0.54	0.69	0.63
PKNaMg(N2) 9/1		0.55	0.65	0.48	0.08	0.44
N2PKNaMg 9/2		0.65	0.78	0.38	0.71	0.63
N2PNaMg 10		0.27	0.46	0.67	0.65	0.51
N3PKNaMg 11/1		1.35	1.20	0.83	1.44	1.21
N3PKNaMgSi 11/2		2.20	1.55	1.12	1.58	1.61
None 12		0.52	0.38	0.55	0.43	0.47
(FYM/F) 13/1		0.88	0.81	0.59	0.54	0.70

Results of the Classical and other Long-term Experiments 2019

19/R/PG/5

FYM/PM 13/2	1.27	2.00	1.45	1.21	1.48
PKNaMg(N2*) 14/1	0.57	1.05	1.40	1.35	1.09
N2*PKNaMg 14/2	1.46	1.84	1.54	1.60	1.61
N3*PKNaMg(N2*) 15	1.68	1.81	1.76	1.29	1.63
N1*PKNaMg 16	1.31	1.61	1.17	0.94	1.26
N1* 17	0.89	0.90	0.81	0.81	0.85
N2KNaMg 18	0.46	0.63	0.68	0.24	0.50
N2KNaMg 18/2	-	-	-	-	0.96
FYM 19/1	-	-	-	-	0.95
FYM 19/2	-	-	-	-	1.28
FYM 19/3	-	-	-	-	1.09
FYM/N*PK 20/1	-	-	-	-	1.05
FYM/N*PK 20/2	-	-	-	-	1.22
FYM/N*PK 20/3	-	-	-	-	0.87

2nd cut mean DM% 24.05

TOTAL OF 2 CUTS DRY MATTER, TONNES/HECTARE

Tables of means

Grand mean		5.09					
Manure	Lime	a	b	c	d	Mean	
N1 1		3.56	3.32	2.94	1.66	2.87	
K 2/1		3.16	3.23	2.48	3.30	3.04	
None(FYM) 2/2		3.72	3.78	3.46	3.14	3.52	
None 3		3.36	3.77	3.64	3.03	3.45	
P 4/1		3.81	4.69	5.04	4.01	4.39	
N2P 4/2		4.83	5.59	5.42	3.21	4.76	
N1PKNaMg 6		5.91	6.43	-	-	6.17	
(P)KNaMg 7/1		4.89	6.38	5.70	3.41	5.10	
PKNaMg 7/2		4.59	6.25	5.87	4.75	5.36	
PNaMg 8		3.86	4.31	4.61	5.44	4.56	
PKNaMg(N2) 9/1		4.63	5.64	4.76	1.08	4.03	

Results of the Classical and other Long-term Experiments 2019

19/R/PG/5

N2PKNaMg	9/2	5.99	6.85	5.57	5.26	5.92
N2PNaMg	10	4.58	5.05	5.75	3.96	4.83
N3PKNaMg	11/1	6.62	6.98	6.49	6.35	6.61
N3PKNaMgSi	11/2	9.06	8.07	6.78	7.25	7.79
None	12	3.42	2.72	3.23	2.88	3.06
(FYM/F)	13/1	4.88	5.15	4.75	4.29	4.77
FYM/PM	13/2	5.28	7.24	6.66	7.27	6.61
PKNaMg(N2*)	14/1	4.02	6.01	6.19	6.19	5.60
N2*PKNaMg	14/2	6.78	7.89	6.39	7.63	7.17
N3*PKNaMg(N2*)	15	7.13	8.29	7.36	7.44	7.55
N1*PKNaMg	16	6.1	6.87	5.53	5.42	5.98
N1*	17	3.93	4.49	3.52	4.10	4.01
N2KNaMg	18	3.95	4.36	3.94	1.35	3.40
N2KNaMg	18/2	-	-	-	-	4.89
FYM	19/1	-	-	-	-	6.45
FYM	19/2	-	-	-	-	6.90
FYM	19/3	-	-	-	-	6.94
FYM/N*PK	20/1	-	-	-	-	6.80
FYM/N*PK	20/2	-	-	-	-	7.06
FYM/N*PK	20/3	-	-	-	-	6.26
TOTAL OF 2 CUTS						
Mean DM%	25.38					