


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1957

[Full Table of Content](#)



57/R/CF/4 and 57/W/CF/4 Potatoes - Control of Skin Spot (Fungicides)

Rothamsted Research

Rothamsted Research (1958) *57/R/CF/4 and 57/W/CF/4 Potatoes - Control of Skin Spot (Fungicides)* ;
Yields Of The Field Experiments 1957, pp 102 - 103 - DOI:

<https://doi.org/10.23637/ERADOC-1-177>

57/Cf/4.1

POTATOES

Control of skin spot (*Oospora pustulans*) by fungicides - Rothamsted (R)
Highfield IV and Woburn (W) Great Hill 1957.

Design: 3 randomized blocks of 12 plots each.

Area of each plot: 0.0141 acres. Area harvested: 0.0071 acres.

Treatments:

None (3 plots per block).

Dung (3 plots per block): 15 tons per acre applied in the furrows.

PCNB, 20% Pentachloronitrobenzene dust applied: To seed at $1\frac{1}{2}$ oz.
per cwt (S); to furrows before planting at 300 lb per acre (F);
(S) and (F).

Nomersan, 10% Thiram dust applied: To seed at 3 oz. per cwt (S);
to furrows before planting at 150 lb per acre (F); (S) and (F).

Basal dressing: 15 cwt compound fertilizer (9% N, 9% P₂O₅, 15% K₂O) per
acre.

Cultivations, etc.:

Highfield IV (R). Ploughed twice: Nov 22, 1956 and Feb 22 - Mar 1,
1957. Ridged, basal fertilizer, dung and fungicides applied,
potatoes hand planted: May 1. Earthed up: July 8. Sprayed
with copper fungicide, 5 lb in 40 gallons per acre: July 31 and
Aug 21. Sprayed with sulphuric acid, 20% BOV, 100 gallons per
acre: Sept 28. Lifted: Oct 4. Variety: Majestic. Previous
crop: Wheat.

Great Hill (W). Ploughed: Feb 6, 1957. Ridged: Apr 24. Dung,
basal fertilizers and fungicides applied, potatoes hand planted:
Apr 25. Earthed up: July 6. Sprayed with copper fungicide at
5 lb in 40 gallons per acre: Aug 3, Aug 23, Sept 3. Sprayed with
sodium arsenite at 1 gallon in 40 gallons per acre: Sept 20.
Lifted: Oct 2. Variety: Majestic. Previous crop: Barley.

Note: Emergence counts and records of incidence of root browning and
skin spot (*Oospora pustulans*), black scurf and stem canker (*Corticium
solani*) and common scab (*Actinomyces scabies*) were made.

In 1956 the Pentachloronitrobenzene dust, PCNB, was recorded in
error as Parachloronitrobenzene

Standard errors per plot. Total tubers.

Highfield IV (R): 1.21 tons per acre or 8.6% (26 d.f.)

Great Hill (W): 1.52 tons per acre or 11.5% (26 d.f.)

57/Cf/4.2

Summary of Results

	0	D	Treatment					NSF	Mean
			PS	PF	FSF	NS	NF		
Total tubers: tons per acre									
Highfield IV, Rothamsted									
		(±0.40)			(±0.70)				
Mean	14.08	14.54	13.81	13.79	14.48	12.65	14.67	13.64	14.08
Increase (±0.81)	-	+0.46 ⁽¹⁾	-0.27	-0.29	+0.40	-1.43	+0.59	-0.44	
(1) ±0.57									
Great Hill, Woburn									
		(±0.51)			(±0.88)				
Mean	12.96	13.08	14.25	14.12	13.17	12.86	12.42	13.85	13.23
Increase (±1.02)	-	+0.12 ⁽¹⁾	+1.29	+1.16	+0.21	-0.10	-0.54	+0.89	
(1) ±0.72									
Percentage Ware									
Highfield IV, Rothamsted ⁽¹⁾									
Mean	70.4	76.3	72.8	74.3	67.8	70.2	66.0	67.7	71.6
Increase	-	+5.9	+2.4	+3.9	-2.6	-0.2	-4.4	-2.7	
Great Hill, Woburn ⁽²⁾									
Mean	87.0	86.4	87.1	91.7	87.4	83.0	86.4	85.7	86.8
Increase	-	-0.6	+0.1	+4.7	+0.4	-4.0	-0.6	-1.3	

Fungicide

D Dung at 15 tons per acre.
 P FCNB (Pentachloronitrobenzene) 20% dust.
 N Nomersan, 10% Thiram dust.

Method of application

S Seed treated.
 F Applied to furrows before planting.

Riddle: (1) 1½" (2) 1⅝"