

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1957

[Full Table of Content](#)



---

### 57/R/CB/3 Barley - Varieties and N

#### Rothamsted Research

Rothamsted Research (1958) *57/R/CB/3 Barley - Varieties and N* ; Yields Of The Field Experiments 1957, pp 81 - 81 - DOI: <https://doi.org/10.23637/ERADOC-1-177>

57/Cb/3

BARLEY

Varieties and levels of nitrogen - Great Field II 1957.

Design: 3 randomized blocks of 5 plots each, plots being split into 2 for the application of nitrogen.

Area of each sub plot: 0.0106 acres. Area harvested: 0.0071 acres.

Treatments: All combinations of:-

Whole plots. Varieties: Domen (1); Herta (2); Ingrid (3);

Plumage Archer (4); Proctor (5).

Sub plots. Nitrogen: 0.3; 0.6 cwt N per acre applied as sulphate of ammonia.

Basal dressing: None.

Cultivations, etc.: Ploughed: Nov 26, 1956. Seed drilled at 2 bushels per acre: Mar 26, 1957. Sulphate of ammonia applied: Mar 28. Sprayed with CMPP at 6 pints in 40 gallons per acre: May 23. Combine harvested: Aug 26. Previous crop: Potatoes.

Standard errors per plot, Grain (at 85% dry matter):

Whole plot: 1.26 cwt per acre or 3.3% (8 d.f.)

Sub plot: 1.73 cwt per acre or 4.6% (10 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

N: cwt per acre	Variety					Mean
	1	2	3	4	5	
	(±1.01)*					
0.3	37.7	40.1	37.9	34.6	39.9	38.1
0.6	36.4	40.1	37.7	32.8	41.1	37.6
Mean (±0.72)	37.0	40.1	37.8	33.7	40.5	37.8
Diff. (±1.41)	-1.3	0.0	-0.2	-1.8	+1.2	-0.5 (±0.63)

\*For use in comparisons other than vertical.

Mean dry matter % as harvested: 84.2