


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1957

[Full Table of Content](#)



57/W/CB/6 Barley - N and Residual Dung N P K

Rothamsted Research

Rothamsted Research (1958) *57/W/CB/6 Barley - N and Residual Dung N P K ; Yields Of The Field Experiments 1957*, pp 85 - 86 - DOI: <https://doi.org/10.23637/ERADOC-1-177>

57/Cb/6.1

BARLEY

Residual effects of dung, N, P and K applied to potatoes 1956 and direct effect of N - Butt Close Woburn 1957.

Design: 4 randomized blocks of 16 plots each, certain high order interactions being confounded with block differences.

Area of each plot: 0.0210 acres. Area harvested: 0.0141 acres.

Treatments: All combinations of:-

To potatoes 1956

Dung: None; 5; 10; 20 tons per acre ploughed in.

N: None; 0.9 cwt N per acre as sulphate of ammonia.

P: None; 0.75 cwt P_2O_5 per acre as superphosphate.

K: None; 1.5 cwt K_2O per acre as muriate of potash.

To barley 1957

N: None; 0.4 cwt N per acre as 'Nitro-Chalk'.

Basal dressing: 0.2 cwt P_2O_5 , 0.4 cwt K_2O as compound fertilizer (10% P_2O_5 , 20% K_2O) per acre.

Cultivations, etc.: Ploughed: Dec 10, 1956. Nitrogen fertilizer applied: Mar 18, 1957. Seed combine drilled with basal fertilizer at $2\frac{1}{2}$ bushels per acre: Mar 19. Sprayed with MCPA at 2 pints in 20 gallons per acre: May 23. Combine harvested: Aug 19. Variety: Herta. Previous crop: Potatoes.

Standard error per plot.

Grain (at 85% dry matter): 4.12 cwt per acre or 18.9% (35 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

	Dung to potatoes 1956:				Mean
	None	tons per acre			
		5	10	20	
Mean (± 1.03)	19.6	20.0	22.4	25.2	21.8
<u>Applied to potatoes 1956</u>					
N: cwt per acre					
None (± 1.46)	19.4	18.6	21.0	25.2	21.0
0.9 (± 1.46)	19.8	21.4	23.8	25.3	22.6
Difference (± 2.06)	+0.4	+2.8	+2.8	+0.1	+1.6 (± 1.03)
P ₂ O ₅ : cwt per acre					
None (± 1.46)	19.6	18.5	23.8	26.0	22.0
0.75 (± 1.46)	19.6	21.4	21.0	24.4	21.6
Difference (± 2.06)	0.0	+2.9	-2.8	-1.6	-0.4 (± 1.03)
K ₂ O: cwt per acre					
None (± 1.46)	19.4	20.3	23.3	26.3	22.3
1.5 (± 1.46)	19.9	19.7	21.5	24.1	21.3
Difference (± 2.06)	+0.5	-0.6	-1.8	-2.2	-1.0 (± 1.03)
<u>Applied to barley 1957</u>					
N: cwt per acre					
None (± 1.46)	16.4	16.7	19.7	23.9	19.2
0.4 (± 1.46)	22.9	23.3	25.1	26.6	24.4
Difference (± 2.06)	+6.5	+6.6	+5.4	+2.7	+5.2 (± 1.03)

Responses to treatments
cwt per acre

Response to	Applied to potatoes 1956						Applied to barley 1957	
	None	N 0.9	P ₂ O ₅ None	P ₂ O ₅ 0.75	K ₂ O None	K ₂ O 1.5	None	N 0.4
<u>Applied to potatoes 1956</u>								
			(± 1.46)					
N	-	-	+1.2	+1.8	+1.2	+1.8	+1.9	+1.1
P ₂ O ₅	-0.7	-0.1	-	-	+1.2	-2.0	-1.2	+0.4
K ₂ O	-1.3	-0.7	+0.6	-2.6	-	-	+0.4	-2.4
<u>Applied to barley 1957</u>								
N	+5.7	+4.9	+4.5	+6.1	+6.7	+3.9	-	-

Mean dry matter % as harvested: 82.1