

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1960

[Full Table of Content](#)



60/R/CD/1 Cereals and Beans - Rotations

Rothamsted Research

Rothamsted Research (1961) *60/R/CD/1 Cereals and Beans - Rotations* ; Yields Of The Field Experiments 1960, pp 105 - 107 - DOI: <https://doi.org/10.23637/ERADOC-1-180>

60/Cd/1.1

CEREALS AND BEANS ROTATIONS

The effect of crop sequences on the incidence of cereal foot and root rot diseases - Great Field I 1960 - the 4th year.

Design: Three series each of 3 randomised blocks of 6 plots, starting in each of the years 1957, 1958 and 1959.

Area of each plot: 0.0305 acres. Area harvested (acres): Winter wheat, series starting 1957 - 0.0096; series starting 1958, Spring wheat, Oats, Barley and Beans - 0.0200.

Treatments:

Crop sequences for each series:

1st year:	WW	WW	WW	SW	O	B
2nd year:	WW	O	O	WW	WW	WW
3rd year:	SW	SW	Be	SW	SW	B

WW = Winter wheat, SW = Spring wheat, O = Oats, B = Barley, Be = Beans.

In the 4th year the plots are split for N and all cropped with winter wheat, the series starting in 1957 falling due for this treatment this year, and receiving N at 0.5, 1.0 cwt per acre in 2 doses on Mar 7 and May 2, 1960 as 'Nitro-Chalk'.

Basal dressing: 2 cwt compound fertiliser (16% P_2O_5 , 16% K_2O) per acre combine drilled with seed (placed in sideband for beans); all blocks received 23 cwt ground chalk per acre in Nov 1956.

Nitrogen for cereals: 0.46 cwt N as 'Nitro-Chalk' 21 per acre to spring wheat and 0.31 cwt N as 'Nitro-Chalk' 21 per acre to oats and barley, all in seedbed. 0.93 cwt N as 'Nitro-Chalk' 21 per acre to winter wheat in the series started in 1959 as spring top dressing, half applied in March and half in May.

Cultivations, etc.: Ploughed: Sept 10, 1959. Winter wheat combine drilled at $2\frac{1}{2}$ bushels per acre; beans placement drilled at 275 lb per acre: Oct 14. 'Nitro-Chalk' applied to oats: Mar 4, 1960. Oats combine drilled at 4 bushels per acre: Mar 5. 'Nitro-Chalk' applied to spring wheat, barley and winter wheat, barley combine drilled at 2 bushels per acre: Mar 7. Spring wheat combine drilled at 3 bushels per acre: Mar 8. Winter wheat sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: Apr 21. 2nd application of 'Nitro-Chalk' to winter wheat: May 2. Spring wheat, barley and oats sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: May 6. Combine harvested: Oats and barley - Aug 16; beans - Aug 19; winter wheat - Aug 31; spring wheat - Sept 12. Varieties: Beans - S.Q; winter wheat - Cappelle; spring wheat - Koga II; barley - Proctor; oats - Sun II.

-60/Cd/1.2

Note. Estimates of plant height, % area lodged, incidence of Eyespot (*Cercospora herpotrichoides*) and Take-all (*Ophiobolus graminis*) and counts of plant shoot and ear number were made.

For details of the previous years' results etc. see 'Results of the Field Experiments' 57/Cd/1, 58/Cd/1 and 59/Cd/1.

Standard errors per plot, Grain (at 85% dry matter):

Series starting:

1957	Winter wheat	
	Whole plot	2.38 cwt per acre or 7.8% (10 d.f.)
	Sub plot	2.83 cwt per acre or 9.2% (12 d.f.)
1958	Spring wheat	1.71 cwt per acre or 6.2% (6 d.f.)
1959	Winter wheat	2.11 cwt per acre or 6.8% (6 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

Series starting in 1957

Winter wheat

Crop in 1957	WW	SW	O	WW	B	WW	Mean
1958	WW	WW	WW	O	WW	O	
1959	SW	SW	SW	SW	B	Be	
N cwt per acre	(±1.64) ⁽¹⁾		(±1.79) ⁽²⁾				
0.5	27.7	20.8	21.7	18.8	33.8	48.4	28.5
1.0	27.3	32.9	28.3	21.3	34.6	51.4	32.6
Mean (±1.36)	27.5	26.8	25.0	20.0	34.2	49.9	30.5
Diff. (±2.31)	-0.4	+12.1	+6.6	+2.5	+0.8	+3.0	+4.1 (±0.94)

Mean dry matter
% as harvested:

79.1

(1) for use in vertical and interaction comparisons

(2) for use in horizontal and diagonal comparisons

60/ca/1.3

Grain (at 85% dry matter): cwt per acre

Crop in 1958 1959	Series starting in 1958					Mean	Barley	Winter beans
	Spring wheat				B		W	
	W	SW	O	W	W		O	
	28.7	26.4	24.7	31.3	27.8	43.5	20.9	
	(±0.99)							
Mean dry matter % as harvested:		81.4				82.1	74.8	

Crop in 1959	Series starting in 1959					Mean	Oats
	Winter wheat				WW		
	W	SW	B	O	WW		
	25.2	21.2	30.2	47.5	31.0	38.5	
	(±1.21)						
Mean dry matter % as harvested:		78.3				83.7	