


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1960

[Full Table of Content](#)



60/R/CF/1 Potatoes - Control of Blight (Copper and Zinc Sprays)

Rothamsted Research

Rothamsted Research (1961) *60/R/CF/1 Potatoes - Control of Blight (Copper and Zinc Sprays)* ;
Yields Of The Field Experiments 1960, pp 119 - 120 - DOI:

<https://doi.org/10.23637/ERADOC-1-180>

60/Cf/1.1

POTATOES

Control of blight (*Phytophthora infestans*) by copper and zinc fungicide sprays and times of spraying - Delharding 1960.

Design: 4 × 4 Latin square.

Area of each plot: 0.1270 acres. Area harvested: 0.0035 acres.

Treatments: 0, 1, 2, 3 as follows:-

Treatment	Stage			
	I	II	III	IV
0	-	-	-	-
1	-	C	C	-
2	-	Z	C	C
3	Z	C	C	-

where stage I = sprayed at closure of leaf canopy, July 14, 1960.

II = sprayed on issue of blight forecast, July 25, 1960.

III = sprayed on blight outbreak, Aug 13, 1960.

IV = sprayed when the deposit from the previous spray had gone, Aug 27, 1960.

C = sprayed with copper oxychloride (15% copper) at 5 lb in 40 gallons per acre.

Z = sprayed with zineb (zinc ethylene bis dithiocarbamate - 65 % active ingredient) at 2 lb in 40 gallons per acre.

Basal dressing: 12 cwt compound fertiliser (10% N, 10% P₂O₅, 18% K₂O) per acre.

Cultivations, etc.: Ground chalk applied at 63 cwt per acre: Sept 14 - 18, 1959. Ploughed: Nov 3 - 12. Ridged, basal dressing applied: Apr 20, 1960. Potatoes machine planted: Apr 22. Earthed up: June 27. Sprayed with undiluted BOV at 15 gallons per acre: Sept 13. Haulm destroyed mechanically: Oct 18. Lifted*: Oct 30. Variety: Majestic. Previous crop: Wheat.

*Hand dug. Harvested area very much reduced, owing to wet conditions.

Note: Fortnightly estimates of yield by sampling, and estimates of foliage destroyed by blight were made commencing early August; also estimates of blight on the tubers at the times of sampling and at harvest.

Standard error per plot.

Total tubers: 0.770 tons per acre or 4.6% (6 d.f.)

60/Cf/1.2

Summary of Results

Spray

	0	1	2	3	Mean
<u>Total tubers: tons per acre</u>					
Mean (± 0.385)	14.43	16.66	16.76	18.46	16.58
Increase (± 0.544)		+2.23	+2.33	+4.03	
<u>Percentage ware (1$\frac{1}{2}$" riddle)</u>					
Mean	92.8	95.2	94.6	96.0	94.6
Increase		+2.4	+1.8	+3.2	

Sprays

- 0 = None.
- 1 = Copper oxychloride (15% copper) at 5 lb in 40 gallons per acre after issue of blight forecast.
- 2 = Zineb (zinc ethylene bis dithiocarbamate - 65% active ingredient) at 2 lb in 40 gallons per acre after issue of blight forecast plus copper oxychloride when the previous deposit had been removed.
- 3 = Zineb after closure of leaf canopy plus copper oxychloride after issue of blight forecast.