


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1960

[Full Table of Content](#)



60/R/CH/1 Kale - Control of Weeds (Thiotriazine)

Rothamsted Research

Rothamsted Research (1961) *60/R/CH/1 Kale - Control of Weeds (Thiotriazine)* ; Yields Of The Field Experiments 1960, pp 128 - 128 - DOI: <https://doi.org/10.23637/ERADOC-1-180>

60/Ch/1

KALE

The control of weeds by thiotriazine sprays - Dell Piece 1960.

Design: 2 randomised blocks of 6 plots each.

Area of each plot: 0.0084 acres. Area harvested: 0.0042 acres.

Treatments: Thiotriazine sprays:-

Sprayed on May 25, 1960:

3 oz active ingredient in 40 gallons per acre (3 plots per block) (E1);

6 oz active ingredient in 40 gallons per acre (2 plots per block) (E2).

Sprayed on June 13, 1960:

3 oz active ingredient in 40 gallons per acre (1 plot per block) (L1).

Basal dressing: 9 cwt compound fertiliser (16% N, 9% P₂O₅, 9% K₂O) per acre.

Cultivations, etc.: Ploughed: Nov 30 - Dec 12, 1959. Basal fertiliser applied: Apr 8, 1960. Seed drilled at 3 lb per acre: Apr 11. 2 rows per plot cut and weighed: Dec 6. Variety: Thousand Head. Previous crop: Spring wheat.

Standard error per plot.

Fresh weight: 2.415 tons per acre or 13.6% (8 d.f.)

Summary of Results

Fresh weight: tons per acre

E1	Spray		Mean
	E2	L1	
19.04 (±0.986)	18.08 (±1.207)	13.26 (±1.707)	17.76

Note: Unsprayed strips within the experimental area gave a mean yield of 8.51 tons per acre.