

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1960

[Full Table of Content](#)



---

### 60/R/CI/1 Grass - Levels of K and N

#### Rothamsted Research

Rothamsted Research (1961) *60/R/CI/1 Grass - Levels of K and N* ; Yields Of The Field Experiments 1960, pp 129 - 130 - DOI: <https://doi.org/10.23637/ERADOC-1-180>

60/Ci/1.1

GRASS

Levels of N and K - Harwoods Piece 1960 - the 3rd year.

Design: 4 randomised blocks of 12 plots each.

Area of each plot: 0.0087 acres. Area harvested: 0.0059 acres.

Treatments: None and all combinations of:-

Nitrogen: 0.3; 0.6; 0.9 cwt N per acre as 'Nitro-Chalk'.

Potash: None; 0.3; 0.6 cwt  $K_2O$  per acre as muriate of potash.

All treatments in the presence of 0.6 cwt  $P_2O_5$  per acre as superphosphate.

In addition 2 plots per block, receiving 0.9 cwt N and 0.6 cwt  $K_2O$  per acre, also received phosphate at either None or 1.2 cwt  $P_2O_5$  per acre as superphosphate.

Note: (1) N and K dressings are applied for each cut. All P dressings are applied once annually.

(2) All treatments were applied to the same plots as in the previous seasons.

Basal dressing: None.

Cultivations, etc.: Ploughed: Dec 2, 1959. 1st dressing of fertilisers applied, seed drilled at 30 lb per acre: Mar 28, 1960. Sprayed with CMPP at 6 pints in 40 gallons per acre: May 24. Cut 3 times: July 21, Sept 13 and Nov 7. Variety: S22 Italian Ryegrass.

Standard errors per plot. Dry matter:

1st cut: 4.00 cwt per acre or 14.9% (33 d.f.)

2nd cut: 1.51 cwt per acre or 7.2% (33 d.f.)

3rd cut: 0.95 cwt per acre or 8.5% (33 d.f.)

Total of 3 cuts: 5.29 cwt per acre or 9.0% (33 d.f.)

Note: (3) For details for the previous years results see "Results of the Field Experiments" 58/Cg/2 and 59/Cg/2.

Errata to "Results of the Field Experiments" 1958 and 1959.

Pages 58/Cg/2.1 and 59/Cg/2.1.

Under 'Treatments' alter the last paragraph to read

"In addition 2 plots per block receiving 0.9 cwt N and 0.6 cwt  $K_2O$  per acre ....."

60/Gi/1.2

Summary of Results

Dry matter: cwt per acre

cwt per acre													
N*	0.0	0.3	0.3	0.3	0.6	0.6	0.6	0.9	0.9	0.9	0.9	0.9	
P <sub>2</sub> O <sub>5</sub>	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.0	1.2	
K <sub>2</sub> O*	0.0	0.0	0.3	0.6	0.0	0.3	0.6	0.0	0.3	0.6	0.6	0.6	Mean
1st cut (±2.00)	14.9	21.3	24.1	27.6	26.8	27.7	31.3	24.9	27.8	33.2	29.7	32.9	26.9
2nd cut (±0.75)	4.9	16.1	15.7	14.4	22.6	23.0	22.3	26.1	26.5	27.1	27.7	25.8	21.0
3rd cut (±0.48)	2.3	9.3	9.8	9.0	11.9	13.3	13.3	12.9	12.9	13.3	12.7	13.3	11.2
Total of 3 cuts (±2.64)	22.1	46.7	49.5	51.0	61.3	64.1	66.9	63.9	67.1	73.7	70.1	72.0	59.0

\*For each cut.

Mean dry matter % as cut:

1st cut:	16.7	3rd cut:	13.4
2nd cut:	18.3	Total of 3 cuts:	16.1