


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1960

[Full Table of Content](#)



60/R/A/1 Broadbalk - Wheat

Rothamsted Research

Rothamsted Research (1961) *60/R/A/1 Broadbalk - Wheat ; Yields Of The Field Experiments 1960*, pp 5 - 6 - DOI: <https://doi.org/10.23637/ERADOC-1-180>

60/A/1.1

WHEAT - BROADBALK 1960

The 117th year

For history, treatments, etc., see "Details of the Classical and Long Term Experiments" 1956.

Cultivations, etc.:

Cropped sections. Ground chalk applied: Sept 4, 1959. Section IA sprayed with 2,4-D ester at $1\frac{3}{4}$ pints in 40 gallons per acre: Sept 8. Dung applied: Sept 21. Ploughed: Sept 21 - 28. Autumn fertilisers applied: Oct 7. Seed drilled at $2\frac{3}{4}$ bushels per acre: Oct 19. Spring fertilisers applied: Apr 4, 1960. Second dressing of nitrate of soda applied to plot 16; Section IA sprayed with CMFP at 6 pints in 40 gallons per acre: Apr 28. Combined: Aug 31 - Sept 9. Variety: Squarehead's Master 13/4.

Fallow section. (III) Ploughed: Sept 21 - 28, 1959; Apr 26, 1960; July 2.

Broadbalk Wilderness. N.

In 1960 grazing of the mown portion of Broadbalk Wilderness was introduced.

Cultivations, etc.: Shrubs grubbed out: Nov 18 - 27, 1959. Part grazed (originally mown): Mar 18 - 21, 1960, Apr 19 - 22, May 17 - 21, June 14 - 18, July 11 - 15, Aug 4 - 8, Sept 2 - 7. The grass was topped after each grazing except the first two.

Summary of Results

Grain (at 85% dry matter): cwt per acre

Section Years after fallow	IV	VA	VB	II	IB	IA	Mean
	1	unlimed 2	limed 2	3	4	9	
2A	24.8	23.0	20.5	23.3	16.8	21.6	22.3
2B	21.8	19.6	18.1	21.2	16.9	19.7	20.0
3	17.3	14.4	12.6	10.1	11.1	11.0	13.1
5	17.4	14.6	11.4	11.5	11.8	11.8	13.5
6	19.1	16.8	16.0	13.2	14.5	13.4	15.8
7	22.5	21.2	19.5	18.7	18.1	18.6	20.1
8	25.7	24.4	17.8	23.5	23.1	21.6	23.2
9	20.3	20.4	18.7	14.9	16.9	18.3	18.1
10	21.5	23.9	22.1	17.3	19.0	20.8	20.4
11	17.7	23.5	21.5	19.1	18.4	19.2	19.6
12	18.8	23.3	19.6	19.8	18.9	20.7	19.9
13	17.4	15.4	14.7	17.5	17.1	16.8	16.7
14	20.1	19.8	16.4	18.0	16.8	20.8	18.6
15	22.2	21.3	12.8	17.9	16.2	18.6	18.6
16	22.9	22.5	19.7	21.9	20.1	23.5	21.8
17	22.3	19.7	17.8	16.3	17.1	15.8	18.6
18	17.0	8.3	9.9	6.7	8.7	7.7	10.4
19	23.5	18.1	14.2	17.4	20.0	18.8	19.1
20				14.5	16.3	14.6	15.2

60/A/1.2

Straw (at 85% dry matter): cwt per acre

Section Years after fallow	IV	VA	VB	II	IB	IA	Mean
		unlimed	limed				
	1	2	2	3	4	9	
2A	45.9	41.0	37.4	34.9	29.9	25.2	37.6
2B	41.9	41.3	37.2	37.8	37.9	18.6	38.1
3	23.4	18.2	14.5	13.2	14.5	11.2	16.8
5	28.0	20.9	19.4	19.1	20.5	14.6	21.6
6	35.0	26.0	29.1	20.7	25.7	16.6	26.7
7	37.4	34.2	32.5	29.4	29.9	23.4	32.3
8	37.9	35.6	35.3	33.1	38.8	29.5	35.6
9	35.8	27.7	27.1	22.0	28.5	26.8	28.3
10	25.0	26.9	27.5	19.1	23.2	23.8	23.7
11	28.9	33.1	25.6	22.1	24.3	22.7	26.2
12	29.1	35.6	28.5	24.6	24.4	27.6	28.0
13	32.8	27.7	29.4	29.0	20.6	24.0	28.4
14	31.2	28.5	26.3	19.9	19.0	21.5	24.9
15	33.1	32.1	24.1	24.7	29.1	27.8	28.6
16	42.0	37.7	36.7	27.1	26.0	27.4	33.7
17	35.8	30.3	30.8	23.7	26.9	18.7	28.9
18	29.1	22.6	24.7	11.8	17.6	8.4	20.1
19	40.5	32.0	29.9	26.3	30.3	24.2	31.7
20				22.6	24.6	18.6	22.6

Mean dry matter % as harvested: Grain 77.3
Straw 90.0