


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1960

[Full Table of Content](#)



60/R/B/10 N Levels and Residues Rotation

Rothamsted Research

Rothamsted Research (1961) *60/R/B/10 N Levels and Residues Rotation* ; Yields Of The Field Experiments 1960, pp 76 - 77 - DOI: <https://doi.org/10.23637/ERADOC-1-180>

60/B/10.1

N LEVELS AND RESIDUES ROTATION

Direct and residual effects of sulphate of ammonia - Long Hoos III
1960 (preliminary year).

Rotation: Wheat, potatoes.

Design (each crop): 3 randomised blocks of 9 plots each.

Area of each plot (acres): 0.0212. Area harvested: Wheat - 0.0141;
Potatoes - 0.0035.

Treatments:

Nitrogen (applied as sulphate of ammonia).
To wheat: None; 0.5; 1.0 cwt per acre.
To potatoes: None; 0.75; 1.5 cwt per acre.
(Three plots per block for each treatment in 1960.)

Basal dressing (per acre):

To wheat: $2\frac{1}{2}$ cwt compound fertiliser, 12% P_2O_5 , 24% K_2O combine
drilled.
To potatoes: 6 cwt compound fertiliser, 12% P_2O_5 , 24% K_2O ,
broadcast on the flat.

Cultivations, etc.: All land ploughed: Nov 3, 1959.

Wheat: Combine drilled with basal fertiliser, sulphate of ammonia
broadcast by hand: Mar 19, 1960. Sprayed with TCB/MCPA at
4 pints in 40 gallons per acre: May 10. Combine harvested:
Sept 13. Variety: Jufy I. Previous crop: Spring wheat.

Potatoes: Basal fertiliser and sulphate of ammonia broadcast on
flat: Apr 13, 1960. Ridged, potatoes planted: Apr 14.
Earthed up: June 21. Sprayed with copper fungicide at 5 lb
in 40 gallons per acre: July 15 and Aug 9. Sprayed with
undiluted BOV at 15 gallons per acre: Aug 31. Haulm destroyed
mechanically: Sept 21. Lifted*: Dec 1. Variety: Ulster
Supreme. Previous crop: Spring wheat.

*Hand dug. Harvested area very much reduced, because of wet
conditions.

Standard errors per plot.

Wheat, grain (at 85% dry matter): 2.48 cwt per acre or 10.4%
(22 d.f.)
Potatoes, total tubers: 0.942 tons per acre or 6.1% (22 d.f.)

60/B/10.2

Summary of Results

Wheat, (grain at 85% dry matter): cwt per acre

	N: cwt per acre		Mean
None	0.5	1.0	
16.4	24.8	30.3	23.8
	(±0.83)		
Mean dry matter % as harvested: 82.3			

Potatoes

	N: cwt per acre		Mean
None	0.75	1.50	
<u>Total tubers: tons per acre</u>			
11.99	16.33	17.71	15.34
	(±0.314)		
<u>Percentage ware (1½" riddle)</u>			
96.1	97.5	98.0	97.2