


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1960

[Full Table of Content](#)



60/R/B/11 and 60/W/B/11 Triazine Weedkiller Rotations

Rothamsted Research

Rothamsted Research (1961) *60/R/B/11 and 60/W/B/11 Triazine Weedkiller Rotations* ; Yields Of The Field Experiments 1960, pp 78 - 82 - DOI: <https://doi.org/10.23637/ERADOC-1-180>

60/B/11.1

TRIAZINE WEEDKILLER ROTATIONS

The direct and residual effects of triazine weedkillers - Rothamsted (R) Great Knott II and Woburn (W) Great Hill I and II 1960.

Owing to the unsuitability of the Rothamsted site this experiment is discontinued. The Woburn experiment will be continued in 1961 in an altered form.

Rotations: Great Knott II (R): Winter beans, winter wheat, potatoes, barley.

Great Hill I and II (W): Potatoes, barley.

Design (each field): 2 randomised blocks of 9 plots each per crop.

Area of each plot (acres):	Area harvested (acres):
Great Knott II (R): 0.0636	Winter beans - 0.0139, winter wheat - 0.0150, potatoes - 0.0035, barley - 0.0139.
Great Hill I and II (W): 0.0482	Potatoes - 0.0107, barley - 0.0115.

Treatments.

Great Knott II (R): All plots were ploughed and received normal cultivations before planting. The potato plots were also rotary cultivated. Cultivations described below were carried out after planting.

To potatoes and beans:

No cultivations	(0)
Normal weed control cultivations	(N)
No cultivations, 1 lb simazine ⁺	(1)
No cultivations, 2 lb simazine ⁺	(2)

To potatoes only:

2 lb simazine ⁺ , then potatoes grubbed and earthed up	(2E)
Potatoes grubbed and earthed up and later 2 lb simazine ⁺ applied before crop emergence	(2L)

To beans only:

Normal weed control cultivations, 1 lb simazine ⁺ in autumn and 1 lb simazine ⁺ in spring	(2D)
---	------

The barley and wheat plots were split* for 0 v. hormone spray for weed control.

Great Hill I and II (W):

To potatoes only:

As above (excluding treatment 2D) except that the plots were not rotary cultivated.

The barley plots were split* for 0 v. hormone spray for weed control.

*1 plot per block was not split, but received hormone spray only. It was intended that it should be followed by a "No weed control" treatment. The yields from these plots are not presented.

⁺In 40 gallons per acre

60/B/11.2

Note: 2 plots for each of treatments 1, 2, N, 2D were included in each block to accommodate a comparison between ploughing and no ploughing in later seasons.

Basal dressings per acre:

Great Knott II (R):

Beans: $4\frac{1}{2}$ cwt compound fertiliser (12% P_2O_5 , 24% K_2O) placement drilled.

Wheat: $2\frac{1}{2}$ cwt compound fertiliser (6% N, 15% P_2O_5 , 15% K_2O) combine drilled: $3\frac{1}{2}$ cwt sulphate of ammonia top dressed.

Potatoes: 10 tons dung: 8 cwt compound fertiliser (10% N, 10% P_2O_5 , 18% K_2O).

Barley: $3\frac{1}{2}$ cwt compound fertiliser (16% N, 9% P_2O_5 , 9% K_2O) combine drilled.

Great Hill I and II (W):

Potatoes: 14 tons dung: 12 cwt compound fertiliser (10% N, 10% P_2O_5 , 18% K_2O).

Barley: 4 cwt compound fertiliser (16% N, 9% P_2O_5 , 9% K_2O) combine drilled.

Cultivations, etc.:

Great Knott II (R):

Beans: Ground chalk applied at 2 tons per acre: Oct 1, 1959.

Ploughed: Oct 3. Seed placement drilled at 275 lb per acre with basal fertiliser: Oct 16. Simazine applied to appropriate plots: Oct 30 and Mar 22, 1960. Treatment N harrowed: Apr 4. Treatment N horse-hoed: Apr 12, Apr 29, May 2. Combine harvested: Aug 20. Variety S.Q. The crop was poor and on one block certain plots were discarded.

Wheat: Ground chalk applied at 2 tons per acre: Oct 1, 1959.

Ploughed: Oct 3. Seed combine drilled at $2\frac{3}{4}$ bushels per acre with basal fertiliser: Oct 16. Top dressed with sulphate of ammonia: Apr 14, 1960. Appropriate sub plots sprayed with CMPP at 6 pints in 40 gallons per acre: Apr 22. Combine harvested: Aug 30. Variety: Cappelle.

Potatoes: Ploughed: Oct 3, 1959. Dung applied: Jan 15 - Feb 9, 1960. Ploughed 2nd time: Feb 10. Basal fertiliser applied:

Mar 28. Rotary cultivated: Apr 14. Potatoes planted: Apr 19. Simazine applied (excluding treatment 2L): Apr 30. Treatments N and 2L tractor weeded: May 16 and May 25.

Treatments N and 2L grubbed: May 26. Treatment 2L earthed up and sprayed with simazine: May 27. Treatments N and 2E grubbed: June 17. Treatments N and 2E earthed up: June 20.

Sprayed with copper fungicide at 5 lb in 40 gallons per acre: July 16 and Aug 10. Sprayed with undiluted BOV at 15 gallons per acre: Aug 31. Haulm destroyed mechanically:

Sept 22. Lifted: Nov 30. Variety: Ulster Supreme.

* Hand dug. Harvested area much reduced owing to wet condition.

60/B/11.3

Barley: Ploughed: Oct 3, 1959. Seed combine drilled at 2 bushels per acre with basal fertiliser: Mar 7, 1960. Appropriate sub plots sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: May 10. Combine harvested: Aug 16. Variety: Proctor. Previous crop (whole area): Spring wheat Great Hill I and II (W):

Potatoes: Dung applied at 14 tons per acre; ploughed: Feb 24, 1960. Basal fertiliser applied: Apr 19. Potatoes planted: Apr 20. Simazine applied (excluding 2L plots): May 2. Treatments N, 2L tractor weeded: May 7. Treatment 2L grubbed and earthed up and simazine applied: May 23. Treatment N tractor weeded: May 31. Treatment N grubbed: June 14. Treatment 2E grubbed, treatments N, 2E earthed up: June 18. Sprayed with zineb at 2 lb in 40 gallons per acre: July 15. Sprayed with copper fungicide at 5 lb in 40 gallons: July 26. Sprayed with undiluted BOV at 15 gallons per acre: Sept 8. Haulm destroyed mechanically: Sept 15. Lifted: Sept 26. Variety: Ulster Supreme. Previous crop: Barley

Barley: Ploughed: Jan 4 - 5, 1960. Seed combine drilled at $2\frac{1}{4}$ bushels per acre with basal fertiliser: Mar 25. Appropriate sub plots sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: May 7. Combine harvested: Aug 22. Variety: Proctor. Previous crop: Potatoes.

Standard errors per plot.

Great Knott II (R).

Winter wheat, grain (at 85% dry matter): 3.23 cwt per acre or 8.3% (13 d.f.)

Potatoes, total tubers: 0.615 tons per acre or 5.3% (11 d.f.)

Barley, grain (at 85% dry matter): 0.87 cwt per acre or 2.5% (15 d.f.)

Great Hill I and II (W).

Potatoes, total tubers: 2.142 tons per acre or 10.0% (11 d.f.)

Barley, grain (at 85% dry matter): 1.33 cwt per acre or 6.0% (15 d.f.)

60/B/11.4

Summary of Results

Winter beans Great Knott II (R)

0	Treatment				Mean
	N	1	2	2D	
<u>Grain (at 85% dry matter): cwt per acre</u>					
24.4	26.8	28.1	23.0	25.2	25.6

Mean dry matter % as harvested: 79.1

Winter wheat Great Knott II (R)

Hormone		Mean	Difference
None	Sprayed		
<u>Grain (at 85% dry matter): cwt per acre</u>			
38.8	39.3	39.0	+0.5(±1.22)

Mean dry matter % as harvested: 79.7

Potatoes

	Treatment						Mean
	0	N	1	2	2E	2L	
<u>Total tubers: tons per acre Great Knott II (R)</u>							
Mean	8.71 (±0.435)	14.16	9.86 (±0.307)	11.86	11.84 (±0.435)	12.14	11.61
Increase		+5.45	+1.15 (±0.533)	+3.15	+3.13 (±0.615)	+3.43	

Total tubers: tons per acre Great Hill I and II (W)

Mean	21.61 (±1.514)	24.24	19.56 (±1.071)	21.01	22.38 (±1.514)	19.15	21.42
Increase		+2.63	-2.05 (±1.906)	-0.60	+0.77 (±2.201)	-2.46	

60/B/11.5

Potatoes

	Treatment						Mean
	0	N	1	2	2E	2L	
<u>Percentage ware (1$\frac{1}{2}$" riddle) Great Knott II (R)</u>							
Mean	96.0	97.8	96.4	97.0	96.8	98.0	97.0
Increase		+1.8	+0.4	+1.0	+0.8	+2.0	

<u>Percentage ware (1$\frac{3}{8}$" riddle) Great Hill I and II (W)</u>							
Mean	99.5	99.4	99.4	99.6	99.6	99.1	99.4
Increase		-0.1	-0.1	+0.1	+0.1	-0.4	

Barley

Grain (at 85% dry matter): cwt per acre

None	Hormone		Mean	Difference
	Sprayed			
<u>Great Knott II (R)</u>				
34.8	33.9		34.3	-0.9 (± 0.31)
<u>Great Hill I and II (W)</u>				
21.9	22.6		22.3	+0.7 (± 0.47)

Mean dry matter % as harvested: Great Knott II (R) 81.2
 Great Hill I and II (W) 79.0