


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1960

[Full Table of Content](#)



60/R/CA/6 Spring Wheat - Forms and Application of N

Rothamsted Research

Rothamsted Research (1961) *60/R/CA/6 Spring Wheat - Forms and Application of N* ; Yields Of The Field Experiments 1960, pp 92 - 93 - DOI: <https://doi.org/10.23637/ERADOC-1-180>

60/Ca/6.1

SPRING WHEAT

Forms of N and methods of application - Little Knott I 1960.

Design: $4 \times 2 \times 4$ in 6 blocks of 16, with certain high order interactions partially confounded with block differences, plus 2 control plots per block.

Area of each plot: 0.0097 acres.

Treatments: No nitrogen and all combinations of:-

<u>Forms of N:</u> Ammonium sulphate	21% N	(S)
Calcium nitrate	15.5% N	(C)
Ammonium nitrate	23% N	(A)
or Urea	45.6% N	(U)

Levels of N: 0.4; 0.8 cwt N per acre.

Methods of application: Broadcast (B), combine drilled (D), side band placed (P), top dressed (T).

Basal dressing: 2 cwt granular compound fertiliser (14% P_2O_5 , 28% K_2O) per acre cross drilled.

Cultivations, etc.: Ploughed: Nov 11, 1959. Ground chalk applied at 23 cwt per acre: Feb 24 - Mar 2, 1960. Seed drilled at 3 bushels per acre and seedbed fertilisers applied: Mar 22. Nitrogen top dressings applied: May 2. Sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: May 10. Combine harvested: Sept 13. Variety: July II Previous crop: Oats.

Standard error per plot.

Grain (at 85% dry matter): 2.01 cwt per acre or 5.8% (67 d.f.)

60/Ca/6.2

Summary of Results

Grain (at 85% dry matter): cwt per acre

	Form of N				Mean
	S	C	A	U	
Mean (± 0.41)	34.6	36.0	35.4	35.4	35.4 (± 0.21)
N: cwt per acre					
0.4 (± 0.58)	33.8	35.3	34.4	34.9	34.6
0.8	35.3	36.8	36.4	35.9	36.1
Diff. (± 0.82)	+1.5	+1.5	+2.0	+1.0	+1.5 (± 0.41)
Method of application					
B	34.0	34.2	34.1	35.2	34.4
D (± 0.82)	34.5	36.4	36.2	35.9	35.7 (± 0.41)
P	36.3	37.5	35.4	34.9	36.0
T	33.4	36.0	36.0	35.8	35.3

	Method of application			
	B	D	P	T
N: cwt per acre				
0.4 (± 0.58)	33.1	35.3	35.4	34.6
0.8	35.6	36.2	36.7	36.0
Diff. (± 0.82)	+2.5	+0.9	+1.3	+1.4

Control: 29.9 (± 0.58)

General Mean 34.8

Mean dry matter % as harvested (all plots): 84.4

Form of N

S = Ammonium sulphate 21% N

C = Calcium nitrate 15.5% N

A = Ammonium nitrate 23% N

U = Urea 45.6% N

Method of application

B = Broadcast

D = Combine drilled

P = Side band placed

T = Top dressed