


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1960

[Full Table of Content](#)



### 60/R/CA/7 and 60/W/CA/7 Spring Wheat - Combine Drilling of N

#### Rothamsted Research

Rothamsted Research (1961) *60/R/CA/7 and 60/W/CA/7 Spring Wheat - Combine Drilling of N* ;  
Yields Of The Field Experiments 1960, pp 94 - 95 - DOI: <https://doi.org/10.23637/ERADOC-1-180>

60/Ca/7.1

SPRING WHEAT

Combine drilling of nitrogen - Rothamsted (R) Little Knott I and Woburn (W) Lansome Field 1960.

Design (each field): 4 randomised blocks of 7 plots each.

Area of each plot: 0.0212 acres. Area harvested: 0.0141 acres.

Treatments:

No nitrogen. 0.41 cwt N per acre ( $N_1$ ); 0.82\* cwt N per acre ( $N_2$ ) either broadcast as sulphate of ammonia or combine drilled as part of a compound fertiliser.

0.41 cwt N per acre as above plus 0.35 cwt N per acre as 'Nitro-Chalk' top dressing.

Compound fertilisers used:

$N_1$ : 8% N, 8%  $P_2O_5$ , 8%  $K_2O$ .

$N_2$ : 16% N, 9%  $P_2O_5$ , 9%  $K_2O$ .

\*Note: 0.88 on Lansome Field, Woburn.

Basal dressing per acre: combine drilled

(a) on the plots receiving drilled nitrogen, as compounds  $N_1$ ,  $N_2$ :-

Little Knott I (R): 0.46 cwt  $P_2O_5$ ; 0.46 cwt  $K_2O$ .

Lansome Field (W): 0.49 cwt  $P_2O_5$ ; 0.49 cwt  $K_2O$ .

(b) on the no nitrogen and broadcast nitrogen plots: as compound 16%  $P_2O_5$ , 16%  $K_2O$ :-

Little Knott I (R): 0.46 cwt  $P_2O_5$ ; 0.46 cwt  $K_2O$ .

Lansome Field (W): 0.60 cwt  $P_2O_5$ ; 0.60 cwt  $K_2O$ .

Note: The rates of application aimed at were  $N_1$ , 0.45;  $N_2$ , 0.8; N top-dressed, 0.35 cwt per acre; basal  $P_2O_5$  and  $K_2O$ , 0.45 cwt per acre. The discrepancies were due to machine application.

Cultivations, etc.:

Little Knott I (R): Ploughed: Nov 10, 1959. Ground chalk applied at 23 cwt per acre: Feb 24 - Mar 2, 1960. Seed combine drilled at  $2\frac{3}{4}$  bushels per acre, sulphate of ammonia applied: Mar 21. 'Nitro-Chalk' top dressings applied: Apr 22. Sprayed with TCB/MCPA at 4 pints in 40 gallons per acre: May 10. Combine harvested: Sept 13. Variety: Jufy I. Previous crop: Oats.

60/Ca/7.2

Lansome Field (W): Sprayed twice with sodium trichloroacetate at 15 lb in 40 gallons per acre: Nov 17 and Dec 29, 1959. Ground chalk applied at 46 cwt per acre: Feb 16, 1960. Seed combine drilled at  $2\frac{3}{4}$  bushels per acre, sulphate of ammonia applied: Mar 22. Sprayed with 2,4-D butoxyethyl ester at  $\frac{1}{2}$  pint in 40 gallons per acre: May 7. Combine harvested: Sept 10. Variety: Jufy I. Previous crop: Potatoes.

Note: Plant counts at germination were made.

Standard errors per plot, Grain (at 85% dry matter):

Little Knott (R): 2.06 cwt per acre or 6.5% (18 d.f.)

Lansome Field (W): 1.58 cwt per acre or 8.4% (18 d.f.)

Summary of Results

Grain (at 85% dry matter): cwt per acre

N: cwt per acre

None	Broadcast			Drilled			Mean
	0.41	0.82*	0.41 & 0.35 <sup>+</sup>	0.41	0.82	0.41 & 0.35 <sup>+</sup>	

Little Knott I (R)

26.9	31.5	33.3	32.5	31.4	33.1	34.8	31.9
			(±1.03)				

Mean dry matter % as harvested: 83.2

Lansome Field (W)

14.2	17.9	19.9	20.8	19.5	20.7	18.5	18.8
			(±0.79)				

Mean dry matter % as harvested: 80.3

\*0.88 at Woburn

<sup>+</sup>Top dressing.