


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1907

[Full Table of Content](#)



Barley. Hoos Field

Barley. Hoos Field, Rothamsted Research (1908) *Yields Of The Field Experiments 1907*, pp 4 - 5 -
DOI: <https://doi.org/10.23637/ERADOC-1-144>

WHEAT. BROADBALK FIELD, 1907.

(See "Guide," page 26, Table XIV.)

Plot.	Manuring.	Dressed Grain.		Straw.
		Yield.	Weight per Bushel.	
		Bushels.	lbs.	Cwt.
2	Farmyard Manure	33.7	60.5	56.6
3	Unmanured	9.1	60.7	9.8
5	Complete Mineral Manure	11.5	60.6	15.1
6	As 5, and single Amm.-salts	23.9	60.4	31.4
7	As 5, and double do.	33.6	60.5	55.8
8	As 5, and treble do.	34.7	59.4	71.6
9	As 5, and single Nitrate Soda	30.2	60.5	46.0
10	Double Amm.-salts alone	27.6	60.2	33.1
11	As 10, and Superphosphate	32.1	60.7	40.5
12	" and Super and Sulph. Soda	41.6	60.9	51.3
13	" and Super and Sulph. Potash	34.7	60.2	57.3
14	" and Super and Sulph. Mag.	36.6	60.9	46.9
15	Double Amm.-salts in Autumn, and Minerals	33.1	61.6	48.6
16	Double Nitrate and Minerals	34.7	60.4	65.7
17	} Minerals alone, or Double Amm.-salts alone, in alternate years }	*10.9	61.3	13.0
18		†31.0	60.7	44.3
19	Rape Cake alone	29.0	60.3	39.3

* Produce by Minerals. † Produce by Ammonium-salts.

BARLEY. HOOS FIELD, 1907.

(See "Guide," page 33, Table XVI.)

Plot.	Manuring.	Dressed Grain.		Straw.
		Yield.	Weight per Bushel.	
		Bushels.	lbs.	Cwt.
1 O	No Minerals, and no Nitrogen	7.7	53.5	6.8
2 O	Superphosphate only	13.0	56.5	9.2
3 O	Alkali salts only	8.1	54.3	14.8
4 O	Complete Minerals	13.2	55.0	24.3
1 A	Amm.-salts only	19.9	53.6	14.2
2 A	Superphos. and Amm.-salts	28.4	55.2	20.0
3 A	Alkali salts and Amm.-salts	20.1	54.4	15.8
4 A	Complete Minerals and Amm.-salts	32.7	55.4	25.5
1 N	Nitrate of Soda alone	22.3	54.0	17.2
2 N	Superphos. and Nitrate Soda	29.0	56.1	25.4
3 N	Alkali salts and Nitrate Soda	21.2	55.0	15.9
4 N	Complete Minerals and Nitrate Soda	31.7	56.1	27.2
1 C	Rape Cake alone	29.1	55.3	21.2
2 C	Superphos. and Rape Cake	28.8	56.1	21.6
3 C	Alkali salts and Rape Cake	26.3	55.9	19.9
4 C	Complete Minerals and Rape Cake	31.1	56.7	23.4
7-1	Unmanured (after Dung, 1852-71)... ..	15.5	55.3	18.4
7-2	Farmyard Dung	42.1	57.1	38.0

BARLEY. HOOS FIELD, 1907.

(Previous cropping: Potatoes, 1876-1901; Barley, 1902 and 1903; Oats, 1904; Barley, 1905 and 1906).

(See "Guide," page 40, Table XIX.)

Plot.	Manures applied to the Potatoes, 1876-1901. Unmanured since.	Dressed Grain.		Straw.	Total Produce.
		Yield.	Weight per Bushel.		
		Bushels.	lbs.	Cwt.	lbs.
1	Unmanured	6.2	56.0	4.2	838
2	Unmanured 1882 and since, previously Dung alone ...	10.3	57.3	7.0	1400
3	Dung 1883-1901	18.9	56.7	15.4	2855
4	Dung 1883-1901	19.1	56.8	14.9	2818

WHEAT AFTER FALLOW (without manure 1851 and since).

HOOS FIELD, 1907.

(See "Guide," page 41, Table 20).

Dressed Grain	{ Yield—14.3 bushels. Weight per bushel—58.6 lbs.
Straw	
Total Produce	19.5 cwt.
		3094 lbs.
