


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1907

[Full Table of Content](#)



Wheat After Fallow

Wheat After Fallow, Rothamsted Research (1908) *Yields Of The Field Experiments 1907*, pp 5 - 5 -
DOI: <https://doi.org/10.23637/ERADOC-1-144>

BARLEY. HOOS FIELD, 1907.

(Previous cropping: Potatoes, 1876-1901; Barley, 1902 and 1903; Oats, 1904; Barley, 1905 and 1906).

(See "Guide," page 40, Table XIX.)

Plot.	Manures applied to the Potatoes, 1876-1901. Unmanured since.	Dressed Grain.		Straw.	Total Produce.
		Yield.	Weight per Bushel.		
		Bushels.	lbs.	Cwt.	lbs.
1	Unmanured	6.2	56.0	4.2	838
2	Unmanured 1882 and since, previously Dung alone ...	10.3	57.3	7.0	1400
3	Dung 1883-1901	18.9	56.7	15.4	2855
4	Dung 1883-1901	19.1	56.8	14.9	2818

WHEAT AFTER FALLOW (without manure 1851 and since).

HOOS FIELD, 1907.

(See "Guide," page 41, Table 20).

Dressed Grain	}	Yield—14.3 bushels.
			Weight per bushel—58.6 lbs.
Straw		19.5 cwt.
Total Produce		3094 lbs.
