


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1949

[Full Table of Content](#)



---

## 49/CE/3 Potatoes - Method of Planting

*49/CE/3 Potatoes - Method of Planting*, Rothamsted Research (1950) Yields Of The Field Experiments 1949, pp 87 - 88

49/Ce/3.1

## POTATOES

The effects of four methods of planting and of three levels of a compound fertilizer.

RP - Great Knott III 1949

System of replication: 4 randomized blocks of 12 plots each.

Area of each plot: 0.014 acres.

Area Harvested: 0.007 acres.

### Treatments:

Methods of planting: Broadcast fertilizer on the flat, ridge, plant in ridges by planting machine (A); broadcast fertilizer on the flat, plant on flat by planting machine (B); ridge, broadcast fertilizer over ridge, plant in furrow by hand (standard method) (C); plant seed and fertilizer with combined seed/fertilizer planting machine (D).

Fertilizer: 0, 8, 16 cwt per acre granular fertilizer containing 7% N, 7% P<sub>2</sub>O<sub>5</sub>, 10½% K<sub>2</sub>O.

Basal Manuring: None.

Cultivations, etc.: Ploughed: Oct 4-7 and Jan 13-15.

Cultivated: Mar 30-31. Harrowed and ring rolled: Apr 1-2.

Harrowed: Apr 20. Applied artificials to A and B, ridged

A and C plots: Apr 29. Applied artificials to C, planted

A, B and C plots: Apr 30. Planted and applied artificials

to D plots: May 2. Re-ridged: May 5. Grubbed: June 3,

7 and 29. Hood: July 5-6. Earthed up: July 14.

Sprayed with 20% B.O.V. solution to kill off haulm: Sept 23.

Lifted: Sept 29. Variety: Majestic. Previous crop: Wheat.

Standard errors per plot:

Total tubers: 0.868 tons per acre or 16.4% {33 d.f.}

Percentage ware: 1.12 {33 d.f.}

49/Ce/3.2

Granular fertilizer cwt per acre	Method of Planting				Mean
	A	B	C*	D	
Total tubers: tons per acre					
			(±0.434)		(±0.217)
0	5.10	5.64	4.23	4.79	4.94
8	5.53	6.05	4.33	6.13	5.51
16	5.81	5.88	5.35	4.72	5.44
Mean (±0.250)	5.48	5.86	4.64	5.21	5.30
Percentage ware					
			(±0.56)		(±0.28)
0	94.2	94.1	94.7	94.5	94.4
8	95.0	95.4	96.4	96.4	95.8
16	94.8	95.8	97.1	97.2	96.2
Mean (±0.32)	94.6	95.1	96.1	96.0	95.5

- A Broadcast fertilizer on the flat, ridge, plant in ridges by planting machines.
- B Broadcast fertilizer on the flat, plant on flat by planting machine.
- C Ridge, broadcast fertilizer over ridges, plant in furrows by hand (present standard method).
- D Plant seed and fertilizer with combined seed/fertilizer planting machine.

\* About half of each C plot suffered mechanical damage during cultivation. No correction has been made for this.