

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1949

[Full Table of Content](#)



49/DA/1 Irrigation - Kasgravo - Sugar Beet

49/DA/1 Irrigation - Kasgravo - Sugar Beet, Rothamsted Research (1950) Yields Of The Field Experiments 1949, pp 91 - 93

49/Da/i.1

SUGAR BEET

The comparison of different rates and methods of irrigation and the effects of three levels of nitrogen.

Milford, Surrey - 1949

System of replication: 6 x 6 Latin Square, plots being split into 4 for the application of nitrochalk.

Area of each sub-plot: 0.0156 acre. Area harvested: 0.00663 acre.

Treatments.

Whole plots. Irrigation: None (duplicate plots); restricted, based on weather data ($6\frac{1}{2}$ ""); restricted at higher level ($8\frac{1}{2}$ ""); full, as decided by farmer (13""); full, with Chilean potash nitrate (16% N, 16% K₂O) in solution 1 in 10,000 (12"").

Sub-plots. Nitrochalk: None, 0.3, 0.6, 0.9 cwt N per acre.

Basal manuring: Compound fertilizer (7% N, 7% P₂O₅, 10.5% K₂O): 6 cwt per acre.

Compound fertilizer (13% P₂O₅, 13% K₂O): 2 cwt per acre.

Salt: $5\frac{1}{2}$ cwt per acre.

Cultivations, etc.: Ploughed and subsoiled: early winter.

Salt applied: Mar 16. Both compound fertilizers and nitrochalk applied: Apr 4. Seed drilled: Apr 11. Singled: May 3. Harvested: Oct 31. Variety: Klein E. Previous crop: Early potatoes, followed by autumn lettuce.

Standard errors per plot:

Total sugar	whole plot, 3.27 cwt per acre or 8.2%
	sub-plot, 3.61 cwt per acre or 9.1%
Roots (washed),	whole plot, 1.02 tons per acre or 6.3%
	sub-plot, 1.34 tons per acre or 8.2%
Sugar percentage,	whole plot, 0.324
	sub-plot, 0.385
Tops,	whole plot, 1.08 tons per acre or 5.7%
	sub-plot, 1.61 tons per acre or 8.5%
Plant number,	whole plot, 0.808 thousands per acre or 2.2%
	sub-plot, 1.37 thousands per acre or 3.6%
Noxious nitrogen,	whole plot, 7.98
	sub-plot, 10.3

All whole plot standard errors estimated from 21 d.f.

All sub-plot standard errors estimated from 93 d.f.

49/Da/1.2

Cwt N per acre including basal	Irrigation: inches					Mean
	0	6½	8½	13	12 plus pot.nitrate	
Total Sugar: cwt per acre						
	(1) and (2)		(3) and (4)			(±0.60)
0.4	36.0	44.3	44.2	44.1	45.1	41.6
0.7	34.6	43.9	44.5	42.7	41.0	40.2
1.0	35.6	42.7	46.2	40.4	40.5	40.2
1.3	32.1	40.8	43.0	39.2	37.5	37.5
Mean	34.6 (±0.94)	42.9	44.5 (±1.34)	41.6	41.0	39.9
Roots (washed): tons per acre						
	(1) and (2)		(3) and (4)			(±0.223)
0.4	14.49	18.06	17.41	17.82	18.21	16.75
0.7	14.30	17.37	17.71	17.33	16.76	16.30
1.0	14.97	18.07	18.34	16.49	17.01	16.64
1.3	13.58	16.63	17.55	16.08	16.10	15.59
Mean	14.34 (±0.296)	17.53	17.75 (±0.418)	16.93	17.02	16.32

Standard errors:

	Total Sugar	Roots (washed)
1 for vertical comparisons	±1.04	±0.387
2 for all others	±1.31	±0.447
3 for vertical comparisons	±1.47	±0.547
4 for all others	±1.85	±0.632

49/Da/1.3

Cwt N per acre including basal	Irrigation: inches					Mean
	0	6½	8½	13	12 plus pot. nitrate	
Sugar Percentage						
	(1) and (2)	(3) and (4)				(±0.064)
0.4	12.42	12.30	12.73	12.40	12.43	12.45
0.7	12.06	12.65	12.57	12.37	12.23	12.32
1.0	11.90	11.87	12.58	12.27	11.95	12.08
1.3	11.81	12.27	12.25	12.20	11.68	12.00
Mean	12.05 (±0.094)	12.27	12.53	12.31 (±0.132)	12.08	12.21
Tops: tons per acre						
	(1) and (2)	(3) and (4)				(±0.268)
0.4	13.45	20.08	18.14	20.43	22.49	18.01
0.7	14.18	19.61	20.85	20.08	23.04	18.66
1.0	14.87	20.09	20.95	21.82	22.51	19.18
1.3	15.19	22.28	21.35	21.45	24.28	19.96
Mean	14.42 (±0.312)	20.52	20.32	20.95 (±0.441)	23.08	18.95

Standard errors:

	Sugar Percentage	Tops
1 for vertical comparisons	±0.111	±0.463
2 for all others	±0.134	±0.508
3 for vertical comparisons	±0.157	±0.655
4 for all others	±0.190	±0.719