

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1974

[Full Table of Content](#)



### 74/R/RN/11 Irrigation - Wheat , Kale

74/R/RN/11 Irrigation - Wheat , Kale , Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 106 - 112 - DOI: <https://doi.org/10.23637/ERADOC-1-119>

74/R/RN/11

IRRIGATION

Object: To study the effects of irrigation on a rotation of crops. Other agronomic factors are included from time to time - Great Field I and II.

Sponsors: B.J. Legg, B.K. French.

The eleventh year, wheat (Gt. Field I), kale (Gt. Field II).

For previous years see 64/C/15(t), 65/C/14(t), 66/C/9(t), 67/C/7(t), 68/C/6(t), 69/R/RN/11(t), 70/R/RN/11(t), 71/R/RN/11(t), 72/R/RN/11(t) and 73/R/RN/11.

Design: 4 randomised blocks of 4 plots, split into half and quarter plots (Gt. Field I)  
2 randomised blocks of 2 plots, split into half and quarter plots (Gt. Field II)

Whole plot dimensions: Wheat - 15.2 x 32.0, kale - 15.2 x 30.5. Sub plot area harvested: Wheat - 0.00356, kale - 0.00098.

Treatments to wheat: All combinations of:-

Whole plots: 1. Irrigation:

None  
Full

IRRIGTN

None  
Full

2. Plant population:

Normal, 18 cm (7 inch) between rows, seed rate 224 kg  
Quarter normal, 36 cm (14 inch) between rows, seed rate 56 kg.

POPULATN

Normal  
Quarter

Half plots: 3. Sowing date:

Autumn, 12 Oct, 1973  
Spring, 27 Mar, 1974

SWING

Autumn  
Spring

Quarter plots: 4. Nitrogen fertiliser (kg N):

45  
90

N

45  
90

Treatments to kale: All combinations of:-

Whole plots: 1. Irrigation:

None  
Full

IRRIGTN

None  
Full

74/R/RN/11

Half plots: 2. Rates of compound fertiliser (20:14:14) kg: COMPFERT

750	750
1130	1130

Quarter plots: 3. Residues of N fertiliser to potatoes in 1973 (kg N): NRES(73)

163	163
326	326

Standard applications:

Wheat: Manures: (0:20:20) at 280 kg, combine drilled. Weedkiller: Dicamba with mecoprop and MCPA (Autumn sown wheat: 'Banlene Plus' at 5.6 l in 220 l, spring sown wheat: 'Tetralex Plus' at 7.0 l in 220 l).

Kale: Weedkiller: Desmetryne ('Semeron 25 WP' at 1.7 kg in 220 l).

Seed: Wheat: Maris Ranger, dressed with dieldrin.

Kale: Maris Kestrel, dressed with BHC and captan, sown at 2.2 kg.

Cultivations, etc.:-

Wheat: Deep-tine cultivated: 28 July, 1973. Rotary cultivated: 29 Aug.

Spring-tine cultivated: 9 Oct. Autumn seed sown: 12 Oct. Plots for spring sowing power harrowed and seed sown: 27 Mar, 1974. N applied: to spring sowing: 8 Mar, to autumn sowing: 19 Apr.

Weedkiller applied: to autumn sowing: 18 Apr, to spring sowing: 28 May. Combine harvested: 18 Sept.

Kale: Ploughed: 17 Dec, 1973. Spring-tine cultivated and NPK applied: 9 Apr, 1974. Rotary harrowed and seed sown: 10 Apr. Weedkiller applied: 3 June. Cut: 19 Nov.

74/R/RN/11

RAINFALL AND IRRIGATION: MM

Week- ending	Rainfall	IRRIGATION	
		WHEAT (Winter & Spring)	KALE
May 4	9.2		
May 11	6.7	15.0	
May 18	4.6	15.0	
May 25	7.0		
June 1	1.9		
June 8	14.1	25.0	
June 15	1.6	20.0	
June 22	23.9		
June 29	36.7		
July 6	8.6		
July 13	10.3		
July 20	8.7		
July 27	1.9	25.0	25.0
Aug 3	2.6		
Aug 10	32.7		
Aug 17	20.7		
Aug 24	2.6		
Aug 31	21.8		
Sept 7	52.4		
Sept 14	5.0		
Sept 21	3.9		
Sept 28	54.1		
Total	331.0	100.0	25.0

Standard errors per plot. Wheat. Grain: tonnes/hectare.

Whole plot: 0.319 or 6.0% (9 d.f.)

Half plot: 0.441 or 8.3% (12 d.f.)

Quarter plot: 0.505 or 9.5% (24 d.f.)

Kale. Total weight: tonnes/hectare.

Quarter plot: 3.17 or 4.0% (4 d.f.)

74/R/RN/11

TABLES OF MEANS

WHEAT

GRAIN: TONNES/HECTARE

	POPULATN		SOWING		N		Mean
	Normal Quarter		Autumn	Spring	45	90	
IRRIGTN							
None	6.05	5.05	7.01	4.09	5.45	5.65	5.55
Full	5.57	4.51	6.26	3.82	4.85	5.23	5.04
	POPULATN						
	Normal Quarter		6.70	4.92	5.65	5.96	5.81
			6.58	2.99	4.64	4.92	4.78
			SOWING				
			Autumn		6.52	6.76	6.64
			Spring		3.78	4.13	3.95
Mean					5.15	5.44	5.29

IRRIGTN	POPULATN	SOWING		SOWING		
		N	Autumn	Spring	Spring	
			45	90	45	90
None	Normal	6.72	7.35	5.06	5.07	
None	Quarter	7.11	6.88	2.92	3.30	
Full	Normal	6.28	6.44	4.56	4.99	
Full	Quarter	5.96	6.36	2.59	3.14	

7<sup>4</sup>/R/RN/11

STANDARD ERRORS OF DIFFERENCES

IRRIGTN	POPULATN	SOWING	N	IRRIGTN POPULATN	IRRIGTN SOWING	IRRIGTN N
0.159	0.159	0.156	0.126	0.226	0.223	0.203
Except when comparing means with same levels of IRRIGTN					0.221	0.179
		POPULATN SOWING	POPULATN N	SOWING N	IRRIGTN POPULATN SOWING N	
Except when comparing means with same levels of POPULATN SOWING IRRIGTN.POPULATN IRRIGTN.POPULATN.SOWING IRRIGTN.POPULATN.N				0.223 0.221	0.203 0.179	0.201 0.179
					0.404	0.401
					0.357	0.401

Mean D.M. % 78.0



74/R/RN/11

WHEAT

STRAW: TONNES/HECTARE

	POPULATN		SOWING		N		Mean
	Normal	Quarter	Autumn	Spring	45	90	
IRRIGTN							
None	4.07	3.27	5.22	2.12	3.75	3.59	3.67
Full	5.20	4.07	5.82	3.45	4.63	4.64	4.64
		POPULATN					
		Normal	6.01	3.27	4.63	4.64	4.64
		Quarter	5.04	2.30	3.75	3.59	3.67
				SOWING			
				Autumn	5.60	5.44	5.52
				Spring	2.78	2.79	2.79
Mean					4.19	4.12	4.15

IRRIGTN	POPULATN	SOWING		Autumn		Spring	
		N		45	90	45	90
None	Normal			5.90	5.54	2.42	2.44
None	Quarter			4.72	4.73	1.97	1.66
Full	Normal			6.25	6.33	3.96	4.27
Full	Quarter			5.53	5.18	2.79	2.77

Mean D.M. % 74.0

74/R/RN/11

KALE

TOTAL WEIGHT: TONNES/HECTARE

	COMPFERT		NRES(73)		Mean
	750	1130	163	326	
IRRIGTN					
None	77.9	85.1	78.4	84.6	81.5
Full	74.1	77.8	74.1	77.8	75.9
	COMPFERT				
		750	71.4	80.6	76.0
		1130	81.0	81.8	81.4
Mean			76.2	81.2	78.7
COMPFERT		750		1130	
NRES(73)	163	326	163	326	
IRRIGTN					
None	73.9	81.8	82.8	87.4	
Full	68.8	79.3	79.3	76.3	

STANDARD ERRORS OF DIFFERENCES

NRES(73)	IRRIGTN(1) NRES(73)	COMPFERT(2) NRES(73)	IRRIGTN(3) COMPFERT NRES(73)
1.58	2.24	2.24	3.17

- (1) Within the same level of IRRIGTN only
- (2) Within the same level of COMPFERT only
- (3) Within the same level of IRRIGTN\*COMPFERT only