

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1974

[Full Table of Content](#)



74/W/RN/13 Intensive Cereals - Ley, Potatoes, Wheat, Barley

74/W/RN/13 *Intensive Cereals - Ley, Potatoes, Wheat, Barley*, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 121 - 127 - DOI: <https://doi.org/10.23637/ERADOC-1-119>

74/W/RN/13

INTENSIVE CEREALS

Object: To study the effects of intensive cereal cropping on yield, incidence of soil-borne diseases and organic matter in the soil-Woburn Stackyard I.

Sponsors: G.W. Cooke, D.B. Slope.

The ninth year, ley, potatoes, winter wheat, barley.

For previous years see 66/B/9(t), 67/B/9, 68/B/7(t), 69/W/RN/13(t), 70/W/RN/13(t), 71/W/RN/13(t) and 72-73/W/RN/13.

Design: For each experiment: 2 randomised blocks of 6 plots, split into 4.

Whole plot dimensions: 8.53 x 20.4. Sub plot area harvested:
Ley - 0.00089. Potatoes - 0.00139. Wheat - 0.00277. Barley - 0.00273.

Treatments:-

One experiment on winter wheat on part of the site of the classical wheat experiment 1877-1954

One experiment on barley on part of the site of the classical barley experiment 1877-1954

Factors tested on both experiments are the same but crop and nitrogen rates differ. All combinations of:-

Whole plots: 1. Previous crops:							PREVCROP
1967	1968	1969	1970	1971	1972	1973	
L	P	C	C	C	L	P	C/C/L/P
P	C	C	C	L	P	C	C/L/P/C
C	C	C	L	P	C	C	L/P/C/C
C	C	L	P	C	C	C	P/C/C/C
C	L	P	C	C	C	L	C/C/C/L
C	C	C	C	C	C	C	C/C/C/C

Ley = 1 year ley P = Potatoes C = Cereal: wheat or barley.

Sub plots: 2. Nitrogen fertiliser (kg N): N

To wheat	To barley	Wheat	Barley
63	50	63	50
126	100	126	100
189	150	189	150
252	200	252	200

NOTE: Ley and potatoes receive basal N only, residues of dressings to cereals are tested (NRESID).

74/W/RN/13

Basal applications: All crops: Manures: Magnesian limestone applied at 3.5 tonnes. P205 at 130 kg, K20 at 260 kg as (0:14:28), half ploughed in, half applied to the plough-farrow. Weedkillers: Paraquat at 0.56 kg ion in 370 l.

Standard applications:

Leys: N at 60 kg, as 'Nitro-Chalk', in seedbed and repeated after sowing and after each cut except the last.

Potatoes: N at 150 kg as 'Nitro-Chalk'. Weedkiller: Linuron at 1.2 kg plus paraquat at 0.28 kg ion in 280 l. Fungicide with insecticide: Mancozeb at 1.3 kg plus demeton-s-methyl at 0.25 kg in 450 l.

Fungicide: Mancozeb at 1.3 kg in 450 l.

Wheat: Weedkiller: Ioxynil at 0.63 kg with mecoprop at 1.9 kg in 280 l on the first occasion and ioxynil at 0.52 kg with mecoprop at 1.6 kg in 280 l on the second occasion.

Barley: Weedkiller: Ioxynil at 0.52 kg and mecoprop at 1.6 kg in 280 l.

Seed: Leys: Italian ryegrass sown at 40 kg.

Potatoes: Pentland Crown.

Wheat: Cappelle, sown at 190 kg.

Barley: Julia, dressed with ethirimol, sown at 160 kg.

Cultivations, etc.: - All plots: Paraquat applied: 12 Sept, 1973. Magnesian limestone applied: 5 Oct. Half PK applied: 10 Oct. Ploughed: 11 Oct. Remaining PK applied: 12 Oct. Spring-tine cultivated: 13 Oct.

Leys: Spring-tine cultivated: 30 Mar, 1974. Power harrowed, N applied, seeds sown: 5 Apr. N applied: 31 May, 2 July, 19 Aug. Cut three times: 1 July, 15 Aug, 24 Sept.

Potatoes: Spring-tine cultivated: 30 Mar, 1974. N applied, rotary cultivated: 9 Apr. Potatoes planted: 10 Apr. Weedkiller applied: 15 May. Rotary ridged: 11 June. Fungicide with insecticide applied: 18 July. Fungicide applied: 7 Aug. Haulm mechanically destroyed: 12 Sept. Sprayed with undiluted BOV at 170 l: 18 Sept. Lifted: 30 Sept.

Wheat: Seed sown: 15 Oct, 1973. Weedkiller applied: 5 Apr, 1974. N applied: 9 Apr. Weedkiller applied on second occasion: 14 May. Combine harvested: 30 Aug.

Barley: Spring-tine cultivated: 27 Mar, 1974. Spring-tine cultivated with crumbler: 28 Mar. Seed sown: 29 Mar. N applied: 1 Apr. Weedkiller applied: 14 May. Combine harvested: 22 Aug.

NOTE: Estimates of eyespot (*Cercosporaella herpotrichoides*) and take-all (*Gaeumannomyces graminis*) were made on both cereal crops in early July.

Standard errors per sub plot.

Wheat, grain: tonnes/hectare: 0.346 or 9.7% (12 d.f.)

Barley, grain: tonnes/hectare: 0.559 or 10.8% (12 d.f.)

74/W/RN/13

TABLES OF MEANS

LEY

WHEAT SITE

DRY MATTER: TONNES/HECTARE

NRESID

63	126	189	252	Mean
1ST CUT				
2.82	2.82	3.16	3.05	2.96
2ND CUT				
3.01	3.09	2.69	3.08	2.97
3RD CUT				
2.54	2.64	2.45	2.34	2.50
TOTAL OF 3 CUTS				
8.37	8.55	8.31	8.48	8.43
Mean D.M. % 1st cut: 17.3				
2nd cut: 18.4				
3rd cut: 16.7				
Total of 3 cuts: 17.5				

74/W/RN/13

LEY

BARLEY SITE

DRY MATTER: TONNES/HECTARE

NRESID

50	100	150	200	Mean
1ST CUT				
2.49	2.88	2.75	2.76	2.72
2ND CUT				
3.09	3.22	3.27	3.12	3.17
3RD CUT				
2.47	2.42	2.69	2.70	2.57
TOTAL OF 3 CUTS				
8.05	8.52	8.71	8.58	8.46
Mean D.M.	%	1st cut:	16.0	
		2nd cut:	16.4	
		3rd cut:	14.6	
		Total of 3 cuts:	15.7	

74/W/RN/13

POTATOES

WHEAT SITE

NRESID

63	126	189	252	Mean
----	-----	-----	-----	------

TOTAL TUBERS: TONNES/HECTARE

48.8	58.4	53.7	54.3	53.8
------	------	------	------	------

PERCENTAGE WARE: 3.81cm (1.5 INCH) RIDDLE

98.2	98.2	98.6	98.9	98.5
------	------	------	------	------

BARLEY SITE

NRESID

50	100	150	200	Mean
----	-----	-----	-----	------

TOTAL TUBERS: TONNES/HECTARE

54.6	55.1	59.0	58.9	56.9
------	------	------	------	------

PERCENTAGE WARE: 3.81 CM (1.5 INCH) RIDDLE

99.1	98.6	99.0	99.0	98.9
------	------	------	------	------

74/W/RN/13

WINTER WHEAT

	N				
	63	126	189	252	Mean
GRAIN: TONNES/HECTARE					
PREVCROP					
C/C/L/P	3.98	5.34	5.53	4.95	4.95
C/L/P/C	2.08	3.60	3.46	2.89	3.01
L/P/C/C	1.78	2.89	3.33	3.41	2.85
C/C/C/C	2.33	3.75	3.97	3.75	3.45
Mean	2.54	3.89	4.07	3.75	3.56

STANDARD ERRORS OF DIFFERENCES

N	PREVCROP*
N	

0.173 0.393

* Within the same level of PREVCROP only

	STRAW: TONNES/HECTARE				
	PREVCROP				
C/C/L/P	4.19	4.96	5.68	5.80	5.16
C/L/P/C	2.87	3.56	4.10	3.78	3.58
L/P/C/C	2.44	2.93	3.16	3.45	2.99
C/C/C/C	3.36	3.98	3.97	4.10	3.85
Mean	3.22	3.86	4.23	4.28	3.89

Mean D.M. % Grain: 82.2
Straw: 83.7

74/W/RN/13

BARLEY

N

	50	100	150	200	Mean
GRAIN: TONNES/HECTARE					
PREVCROP					
C/C/L/P	4.36	5.85	6.10	5.38	5.42
C/L/P/C	3.70	5.70	5.83	5.74	5.24
L/P/C/C	3.66	5.32	5.73	5.25	4.99
C/C/C/C	3.63	5.36	5.39	5.68	5.01
Mean	3.83	5.56	5.76	5.51	5.17

STANDARD ERRORS OF DIFFERENCES

N	PREVCROP*
0.279	0.533

* Within the same level of PREVCROP only

STRAW: TONNES/HECTARE

PREVCROP					
C/C/L/P	2.72	4.13	4.73	4.57	4.04
C/L/P/C	2.13	3.56	4.07	4.40	3.54
L/P/C/C	2.18	3.69	3.72	4.38	3.49
C/C/C/C	1.98	3.08	3.31	3.74	3.03
Mean	2.25	3.62	3.96	4.27	3.52

Mean D.M. % Grain: 85.5
Straw: 87.8