

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1974

[Full Table of Content](#)



### 74/W/CS/99 Effects of Breaks on Take-all - Barley, Beans

74/W/CS/99 *Effects of Breaks on Take-all - Barley, Beans*, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 214 - 215 - DOI: <https://doi.org/10.23637/ERADOC-1-119>

74/W/CS/99

EFFECTS OF BREAKS ON TAKE-ALL

Object: To study the phenomenon of 'take-all' (*Gaeumannomyces graminis*) decline in barley - Woburn Butt Furlong.

Sponsor: D. Hornby.

The third year, barley, spring beans.

For previous years see 72/W/CS/99(t) and 73/W/CS/99.

Design: 2 blocks of 9 plots.

Whole plot dimensions: 5.33 x 15.2. Area harvested: Barley - 0.00434, spring beans - 0.00406.

Treatments: Previous crops:-

PREVCROP

1968-71	1972	1973	
B	B	B (5 plots per block)	B/B/B
B	F	B	B/F/B
B	F	BE	B/F/BE

B = Barley, BE = Beans, F = Fallow

Standard applications:

All plots: Weedkiller: Paraquat at 0.56 kg ion in 370 l.

Barley: Manures: (20:15:15) at 410 kg combine drilled. Weedkiller:

Ioxynil at 0.53 kg and mecoprop at 1.6 kg in 280 l.

Spring beans: Manures: (0:14:28) at 400 kg placement drilled.

Weedkiller: Simazine at 1.1 kg in 390 l. Insecticide: Demeton-s-methyl at 0.25 kg in 390 l.

Seed: Barley: Julia, dressed with ethirimol, sown at 160 kg.

Spring beans, Minor sown at 200 kg.

Cultivations, etc.:-

All plots: Paraquat applied: 12 Sept, 1973. Ploughed: 25 Oct - 5 Nov.

Spring-tine cultivated with crumbler: 7 Mar, 1974,

Barley: Spring-tine cultivated with crumbler second time: 22 Mar, 1974.

Seed sown: 26 Mar. Weedkiller applied: 8 May. Combine harvested: 21 Aug.

Spring beans: Seed sown: 7 Mar, 1974. Weedkiller applied: 23 Mar.

Insecticide applied: 21 June. Combine harvested: 17 Sept.

Fallow: Spring-tine cultivated with crumbler second time: 22 Mar, 1974.

Power harrowed: 7 May.

NOTE: Soil samples were taken before sowing and after harvest and plant samples in July for incidence of 'take-all' (*Gaeumannomyces graminis*).

Standard error per plot.

Barley. Grain: tonnes/hectare: 0.328 or 8.1% (10 d.f.)

74/W/CS/99

TABLES OF MEANS

BARLEY. GRAIN: TONNES/HECTARE

PREVCROP

B/B/B	B/F/B	B/F/BE	Mean
3.98	4.34	4.20	4.06

STANDARD ERRORS OF DIFFERENCES

PREVCROP

B/B/B v B/F/B or B/F/BE	0.254
B/F/B v B/F/BE	0.328

SPRING BEANS. GRAIN: TONNES/HECTARE

Mean 3.84

Mean D.M.  $\frac{1}{2}$  77.4