

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1974

[Full Table of Content](#)



### 74/R/CS/116 Effects of Stem Eelworm - Beans

74/R/CS/116 *Effects of Stem Eelworm - Beans* , Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 235 - 236 - DOI: <https://doi.org/10.23637/ERADOC-1-119>

74/R/CS/116

EFFECTS OF STEM EELWORM

Object: To study the effects of applying field bean straw infested with two races of stem eelworm (*Ditylenchus dipsaci*) on the yield and subsequent infestation of field beans - Highfield 0 and E III.

Sponsor: D.J. Hooper.

The second year, spring beans.

For previous year see 73/R/CS/116.

Design: 3 randomised blocks of 6 plots.

Whole plot dimensions: 4.27 x 9.14. Area harvested: 0.00293.

Treatments: All combinations of:-

- |   |         |
|---|---------|
| 1. Bean straw infested with stem eelworm ( <i>Ditylenchus dipsaci</i> ) worked into soil, autumn 1972:- | EELWORM |
| Oat race  | Oat     |
| Giant race  | Giant   |
| 2. Rate of application of straw (tonnes) to give populations of eelworms:-                              | RATE    |
| None  | 0       |
| Single, 3.1   | Single  |
| Double, 6.2   | Double  |

Basal applications: Manures: (0:14:28) at 400 kg. Weedkillers: Simazine at 0.84 kg in 340 l, paraquat at 0.56 kg ion in 220 l. Insecticide: Demeton-s-methyl at 0.25 kg in 220 l.

Seed: Minor, sown at 220 kg.

Cultivations, etc.:- Paraquat applied: 4 Oct, 1973. Discd and ploughed: 18 Dec. Spring-tine cultivated: 8 Mar, 1974. Power harrowed, seed sown and spring-tine cultivated: 26 Mar. Simazine applied: 5 Apr. Insecticide applied: 13 June. Combine harvested: 30 Sept.

NOTE: Stems showing symptoms of attack by stem eelworm were counted in mid-August. Samples of seed were taken at maturity to assess seed infestation.

Standard error per plot.

Grain, tonnes/hectare: 0.461 or 23.0% (11 d.f.)

74/R/CS/116

TABLES OF MEANS

GRAIN: TONNES/HECTARE

	RATE			Mean
	0	Single	Double	
EELWORM				
Oat		2.39	1.84	2.12
Giant		1.83	1.61	1.72
Mean	2.17	2.11	1.72	2.00

STANDARD ERRORS OF DIFFERENCES

EELWORM	RATE	EELWORM RATE
0.266	0.266	0.376

Mean D.M. % 74.5