Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1974



Full Table of Content

74/R/CS/123 Effects of Viruses - Ryegrass, Red Clover

74/R/CS/123 Effects of Viruses - Ryegrass, Red Clover, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 237 - 242 - DOI: https://doi.org/10.23637/ERADOC-1-119

EFFECTS OF VIRUSES

Object: To study the effects of virus infection on yield and persistence of ryegrass and red clover in pure and mixed stands - Carden Plot 14.

"Sponsors: A.J. Cockbain, R.T. Plumb.

The second year, red clover and Italian ryegrass.

For previous year see 73/R/CS/123.

Design: 4 randomised blocks of 8 plots split into 2.

Whole plot dimensions: 3.35 x 1.22. Area harvested: 0.00011.

Treatments: All combinations of:-

Whole plots: 1. Crop and inoculation with virus: CROP VIR

Red clover uninoculated	Clover/O
Italian ryegrass uninoculated	Grass/O
Red clover and Italian ryegrass (mixed within rows)	12
uninoculated	CloGr/O
Red clover inoculated with Pea Mosaic Virus (PMV)	Clover/P
Italian ryegrass inoculated with Ryegrass Mosaic	
Virus (RMV)	Grass/R
Red clover and Italian ryegrass (mixed) inoculated	52.6.1 (AEC-4007800009)
with PMV	CloGr/P
Red clover and Italian ryegrass (mixed) inoculated	50000-50-60 3 0-50-
with RMV	CloGr/R
Red clover and Italian ryegrass (mixed) inoculated	100000000000000000000000000000000000000
with PMV and RMV	CloGr/PR
	(6.75)
Sub plots: 2. Aldicarb (kg) after each cut:	ALDICARB

NOTE: Inoculation with PMV was unsuccessful.

None

10

Basal applications: Manures: (0:14:28) at 800 kg, 'Nitro-Chalk' at 300 kg after each cut.

Cultivations, etc.:- PK applied: 30 Oct, 1973. Aldicarb applied: 3 June, 1974, 25 July, 19 Sept. Cut and N applied: 28 May, 11 July, 9 Sept.

0

10

- NOTES: (1) Crop was scored for Ryegrass Mosaic Virus: 2 May.
 - (2) Mites (Abacarus hystrix) were counted on 11 July, 9 Sept, 21 Oct.
 - (3) Red clover was sampled for virus incidence during the period 29 Oct to 10 Nov.
 - (4) 3rd cut. Because of a faulty balance the yields from 3 whole plots CROP VIR - Clogr/PR, Clover/O, and Clogr/P were not recorded. Estimated values were used in the analysis.

```
Standard errors per plot. Dry matter: tonnes/hectare:

1st cut.

Whole plot: 0.833 or 10.7% (21 d.f.)

Sub plot: 0.761 or 9.7% (24 d.f.)

2nd cut.

Whole plot: 0.585 or 10.3% (21 d.f.)

Sub plot: 0.419 or 7.4% (24 d.f.)

3rd cut.

Whole plot: 0.769 or 14.3% (18 d.f.)

Sub plot: 0.640 or 11.9% (21 d.f.)

Total of 3 cuts. Whole plot: 1.766 or 9.4% (18 d.f.)

Sub plot: 1.012 or 5.4% (21 d.f.)
```

TABLES OF MEANS

1ST CUT. DRY MATTER: TONNES/HECTARE

ALDICARB

	0	10	Mean
CROP VIR			
Clover/O Grass/O CloGr/O Clover/P Grass/R CloGr/P CloGr/R CloGr/PR	7.27 7.95 8.17 6.11 7.78 8.58 7.87 6.89	7.31 8.98 7.84 6.66 8.46 8.76 8.84 7.47	7.29 8.47 8.00 6.38 8.12 8.67 8.35 7.18
Mean	7.58	8.04	7.81

STANDARD ERRORS OF DIFFERENCES

CROP VIR	ALDICARB	CROP VIR
0.589	0.190 comparing means	0.702
with same		1.00
CROP	VIR	0.538

2ND CUT. DRY MATTER: TONNES/HECTARE

ALDICARB

	o	10	Mean
CROP VIR			
Clover/O Grass/O CloGr/O Clover/P Grass/R CloGr/P CloGr/R CloGr/PR	5.74 4.10 5.61 5.59 4.25 5.33 6.18 5.93	6.75 5.70 5.36 6.16 5.52 6.18 6.10 6.01	6.24 4.90 5.48 5.87 4.89 5.75 6.14 5.97
Mean	5.34	5.97	5.66

STANDARD ERRORS OF DIFFERENCES

CROP VIR ALDICARB CROP VIR
ALDICARB

0.414 0.105 0.464

Except when comparing means with same level of CROP VIR 0.296

3RD CUT. DRY MATTER: TONNES/HECTARE

ALDICARB

///			
	0	10	Mean
CROP VIR			
Clover/O Grass/O CloGr/O Clover/P Grass/R CloGr/P CloGr/R	5.10 2.61 5.73 5.36 3.17 5.97 5.89 6.10	7.34 4.07 6.26 5.85 4.18 6.37 6.12 5.84	6.22 3.34 6.00 5.61 3.67 6.17 6.00 5.97
Mean	4.99	5.75	5.37

STANDARD ERRORS OF DIFFERENCES

CROP VIR	ALDICARB	CROP VIR ALDICARB
	0.160 comparing means	0.631
with same CROP V		0.453

TOTAL OF 3 CUTS. DRY MATTER: TONNES/HECTARE

ALDICARB

	0	10	Mean
CROP VIR			
Clover/O Grass/O CloGr/O Clover/P Grass/R CloGr/P CloGr/R	17.77 14.66 19.51 17.05 15.20 19.80 19.93 18.59	21.78 18.75 19.46 18.67 18.16 21.39 21.06	19.77 16.71 19.48 17.86 16.68 20.59 20.50 18.91
Mean	17.81	19.81	18.81

STANDARD ERRORS OF DIFFERENCES

CROP VIR	ALDICARB	CROP VIR ALDICARB
1.248	0.253	1.347
Except when	comparing means	
with same	level of	
CROP	VIR	0.716
Mean D.M. %	1st cut:	21.1
- 10.000 to 0.000 (0.000 (0.000 (0.000)	2nd cut:	21.2
	3rd cut:	20.8
	Total of 3 cuts:	