

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1974

[Full Table of Content](#)



### 74/R/CS/135 Direct Drilling and Diseases - Winter Barley

74/R/CS/135 *Direct Drilling and Diseases - Winter Barley*, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 255 - 256 - DOI: <https://doi.org/10.23637/ERADOC-1-119>

74/R/CS/135

DIRECT DRILLING AND DISEASES

Object: To study the effects of direct drilling on pathogens and yield of winter barley - Meadow.

Sponsor: R.D. Prew.

The first year, winter barley.

Design: 2 randomised blocks of 3 plots.

Whole plot dimensions: 26.7 x 30.5. Area harvested: 0.00929.

Treatments: Cultivations:	CULTIVATION
Ploughed, spring-tine cultivated, drilled (P)	Plough
Chisel ploughed twice, disced, drilled (C)	Chisel
Paraquat to stubble, direct drilled (D)	Direct

NOTE: Paraquat was applied at 0.56 kg ion in 220 l.

Basal applications: Manures: (10:24:24) at 250 kg combine drilled. 'Nitro-Chalk' at 300 kg. Weedkiller: Mecoprop ('Compitox Extra' at 4.2 l in 220 l).

Seed: Maris Otter sown at 170 kg, dressed fungicide only.

Cultivations, etc.: - P plots ploughed, C plots deep-tine cultivated: 6 Sept, 1973. D plots sprayed with paraquat: 18 Sept. C plots deep-tine cultivated: 10 Oct. P plots spring-tine cultivated, C plots disced and all plots sown: 11 Oct. N applied: 17 Apr, 1974. Mecoprop applied: 1 May. Combine harvested: 30 Aug. Previous crops: Barley 1972 and 1973.

NOTE: Root diseases were assessed in spring and summer and foliar diseases throughout the year.

74/R/CS/135

TABLE OF MEANS

GRAIN: TONNES/HECTARE

CULTIVIN

Plough	Chisel	Direct	Mean
5.05	5.36	4.33	4.91

Mean D.M. % 85.4