Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1974



Full Table of Content

74/R/CS/140 Chemical Reference Plots - Barley

74/R/CS/140 Chemical Reference Plots - Barley, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 262 - 264 - DOI: https://doi.org/10.23637/ERADOC-1-119

74/R/CS/140

CHEMICAL REFERENCE PLOTS

Object: To study the persistence in soil of four agricultural chemicals applied annually, singly and in combination and their effects on soil microflora and on yield of barley - Long Hoos V3.

Sponsors: G.G. Briggs, N. Walker, R. MacDonald.

The first year, barley.

Design: Two replicates of 2 x 2 x 2 x 2 fully randomised.

Whole plot dimensions: 4.06 x 4.57. Area harvested: 0.00075.

Treatments: All combinations of:-

1. Fungicide: FUNGCIDE

None
Benomyl at 4 kg applied 28 March, 1974
Benomyl

2. Insecticide: INSCTCDE

None None Chlorfenvinphos at 2 kg applied 28 March, 1974 Chlorfen

3. Nematicide: NEMACIDE

None
Aldicarb at 6 kg applied 28 March, 1974
Aldicarb

4. Weedkiller: WEEDKILR

None None Chlortoluron at 2 kg applied 2 Apr, 1974 Chlortol

Basal applications: Manures: (0:20:20) at 1000 kg. 'Nitro-Chalk' at 380 kg. Weedkiller: Dicamba with mecoprop and MCPA ('Banlene Plus' at 5.6 l in 340 l).

Seed: Julia - undressed, sown at 160 kg.

74/R/CS/140

Cultivations, etc.:- PK applied: 24 Oct, 1973. Ploughed: 7 Dec. Spring-tine cultivated: 27 Mar, 1974. Power harrowed: 28 Mar. Seed drilled and N applied: 29 Mar. Weedkiller applied: 21 May. Combine harvested: 21 Aug. Previous crops: Maize and sweetcorn 1972, potatoes 1973.

NOTE: The following assessments were made:
Seedling emergence, weed numbers in April, mildew and aphid
numbers in June, barley yellow dwarf virus in July, algae
on soil surface in July and chemical residues throughout
the season.

Standard error per plot.
Grain, tonnes/hectare: 0.343 or 5.6% (16 d.f.)

74/R/CS/140

TABLES OF MEANS

GRAIN: TONNES/HECTARE

		CTCDE Chlorfen		ALDE Aldicarb		EDKLLR Chlortol	Mean
FUNGCIDE							
None Benomyl	5.99 6.19	6.15 6.36	5.94 6.25		6.19 6.58	5•95 5•97	6.07 6.28
	None Chlorfen		5.97 6.22	6.21 6.29	6.24 6.53		6.09 6.26
			None Aldicarb		6.1 9 6. 58	6.00 5.92	6.09 6.25
Mean					6.39	5.96	6.17
	NEMACIDE WEEDKLLE			one Chlortol		dicarb Chlortol	
FUNGCIDE	INSCICDE	3					-
None	None Chlorfen		5.62 6.21	6.04 5.89	6.45	5.84 6.01	
Benomyl	None Chlorfer		6.20 6.72	6.01	6.68 6.72		
STANDARD I	ERRORS OF I	DIFFERENCE	ES .				
FUNGCIDE 1	INSCICDE NI	EMACIDE WI	ŒDKLLR	FUNGCIDE INSCICDE	FUNGCI NEMACI	DE FUNGCII DE WEEDKLI	DE INSCICDE LR NEMACIDE
0.121	0.121	0.121 0.121		0.172	0.1	.72 0.1	72 0.172
INSCICDE I	NEMACIDE F	UNGCIDE					

WEEDKILR WEEDKILR INSCICDE

NEMACIDE

WEEDKLLR

0.343 0.172 0.172

Mean D.M. % 85.3