

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1974

[Full Table of Content](#)



74/R/CS/140 Chemical Reference Plots - Barley

74/R/CS/140 Chemical Reference Plots - Barley, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 262 - 264 - DOI: <https://doi.org/10.23637/ERADOC-1-119>

74/R/CS/140

CHEMICAL REFERENCE PLOTS

Object: To study the persistence in soil of four agricultural chemicals applied annually, singly and in combination and their effects on soil microflora and on yield of barley - Long Hoos V3.

Sponsors: G.G. Briggs, N. Walker, R. MacDonald.

The first year, barley.

Design: Two replicates of 2 x 2 x 2 x 2 fully randomised.

Whole plot dimensions: 4.06 x 4.57. Area harvested: 0.00075.

Treatments: All combinations of:-

	FUNGICIDE
1. Fungicide:	
None	None
Benomyl at 4 kg applied 28 March, 1974	Benomyl
2. Insecticide:	INSECTICIDE
None	None
Chlorfenvinphos at 2 kg applied 28 March, 1974	Chlorfen
3. Nematicide:	NEMACIDE
None	None
Aldicarb at 6 kg applied 28 March, 1974	Aldicarb
4. Weedkiller:	WEEDKILLER
None	None
Chlortoluron at 2 kg applied 2 Apr, 1974	Chlortol

Basal applications: Manures: (0:20:20) at 1000 kg. 'Nitro-Chalk' at 380 kg. Weedkiller: Dicamba with mecoprop and MCPA ('Banlene Plus' at 5.6 l in 340 l).

Seed: Julia - undressed, sown at 160 kg.

74/R/CS/140

Cultivations, etc.: - PK applied: 24 Oct, 1973. Ploughed: 7 Dec.
Spring-tine cultivated: 27 Mar, 1974. Power harrowed: 28 Mar.
Seed drilled and N applied: 29 Mar. Weedkiller applied: 21 May.
Combine harvested: 21 Aug. Previous crops: Maize and sweetcorn
1972, potatoes 1973.

NOTE: The following assessments were made:
Seedling emergence, weed numbers in April, mildew and aphid
numbers in June, barley yellow dwarf virus in July, algae
on soil surface in July and chemical residues throughout
the season.

Standard error per plot.

Grain, tonnes/hectare: 0.343 or 5.6% (16 d.f.)

74/R/CS/140

TABLES OF MEANS

GRAIN: TONNES/HECTARE

	INSCTCDE		NEMACIDE		WEEDKLLR		Mean
	None	Chlorfen	None	Aldicarb	None	Chlortol	
FUNGCIDE							
None	5.99	6.15	5.94	6.20	6.19	5.95	6.07
Benomyl	6.19	6.36	6.25	6.31	6.58	5.97	6.28
	INSCTCDE						
	None	Chlorfen	5.97	6.21	6.24	5.94	6.09
			6.22	6.29	6.53	5.98	6.26
	NEMACIDE						
	None	Aldicarb	6.19	6.00	6.19	6.09	
			6.58	5.92	6.58	6.25	
Mean					6.39	5.96	6.17
	NEMACIDE		None		Aldicarb		
	WEEDKLLR		None	Chlortol	None	Chlortol	
FUNGCIDE	INSCTCDE						
None	None		5.62	6.04	6.45	5.84	
	Chlorfen		6.21	5.89	6.48	6.01	
Benomyl	None		6.20	6.01	6.68	5.88	
	Chlorfen		6.72	6.06	6.72	5.94	

STANDARD ERRORS OF DIFFERENCES

FUNGCIDE INSCTCDE NEMACIDE WEEDKLLR FUNGCIDE FUNGCIDE FUNGCIDE INSCTCDE
INSCTCDE NEMACIDE WEEDKLLR NEMACIDE

0.121 0.121 0.121 0.121 0.172 0.172 0.172 0.172

INSCTCDE NEMACIDE FUNGCIDE
WEEDKLLR WEEDKLLR INSCTCDE
NEMACIDE
WEEDKLLR

0.172 0.172 0.343

Mean D.M. % 85.3