

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1974

[Full Table of Content](#)



74/R/B/5 Systemic Fungicides Study - Barley

74/R/B/5 Systemic Fungicides Study - Barley, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 301 - 302 - DOI: <https://doi.org/10.23637/ERADOC-1-119>

74/R/E/5

BARLEY

SYSTEMIC FUNGICIDE STUDY

Object: To study the effectiveness of different methyl benzimidazol-2-yl-carbamate (MBC) precursors and to relate chemical measurements of persistence, movement and conversion to MBC to field performance - Delafield.

Sponsor: I.J. Graham-Eryce.

Design: 3 randomised blocks of 6 plots.

Whole plot dimensions: 2.41 x 9.14. Area harvested: 0.00151.

Treatments: Fungicidal seed dressings (at 1.0 kg/ha):	FUNGCIDE
None	None
Benomyl	Benomyl
Carbendazim	Carbenda
Cypendazol	Cypendaz
Thiophanate methyl	Thiophan
R 28921 (2-((3-methoxy carbonyl)-thioureido)-0, 0-diethyl-phosphoranilide)	R 28921

NOTE: Seed was naturally infected with loose smut (*Ustilago nuda*).

Basal applications: Manures: (0:20:20) at 1060 kg. (20:14:14) at 440 kg.
Weedkiller: Dicamba with mecoprop and MCPA ('Tetralex Plus' at 7.0 l in 220 l).

Seed: Sultan, sown at 160 kg.

Cultivations, etc.:- PK applied: 10 Sept, 1973. Ploughed: 28 Nov. NPK applied, spring-tine cultivated twice: 1 Apr, 1974. Seed sown: 3 Apr. Weedkiller applied: 17 Apr. Combine harvested: 22 Aug. Previous crops: Potatoes 1972, winter wheat 1973.

NOTES: Plant counts were made shortly after germination. Loose smut (*Ustilago nuda*) was assessed at end of June, eyespot (*Cercospora herpotrichoides*) in early July and mildew (*Erysiphe graminis*) on three occasions.

Standard error per plot.

Grain, tonnes/hectare: 0.200 or 4.4% (10 d.f.).

74/R/B/5

TABLES OF MEANS

GRAIN: TONNES/HECTARE

FUNGICIDE						Mean
None	Benomyl	Carbenda	Cypendaz	Thiophan	R 28921	
3.81	4.86	4.53	4.86	5.02	4.04	4.52

STANDARD ERROR OF DIFFERENCES

FUNGICIDE

0.163

Mean D.M.% 87.9