

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1974

[Full Table of Content](#)



74/R/B/14 Slow-release N - Barley

74/R/B/14 *Slow-release N - Barley*, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 312 - 315 - DOI: <https://doi.org/10.23637/ERADOC-1-119>

74/R/B/14

BARLEY

SLOW-RELEASE N

Object: To compare the effects of slow-release nitrogen fertiliser ('Gold-N', sulphur-coated urea) with a conventional form ('Nitro-Chalk', ammonium nitrate/calcium carbonate) on the yield of barley - Long Hoos VI/VII.

Sponsors: D. Cox, T.M. Addiscott.

Design: 2 randomised blocks of 26 plots.

Whole plot dimensions: 4.27 x 9.14. Area harvested: 0.00195.

Treatments: All combinations of:-

1. Form and time of applying nitrogen fertiliser:	N FORM(1)
'Gold-N' (sulphur-coated urea), to seedbed	Gold-N/E
'Nitro-Chalk' (ammonium nitrate/calcium carbonate) to seedbed	Nitro/E
2. Rate of nitrogen fertiliser (kg N):	N RATE(1)
None	0
15	15
30	30
45	45
60	60
75	75
90	90
105	105
120	120

plus all combinations of:-

1. Form and time of applying nitrogen fertiliser:	N FORM(2)
'Nitro-Chalk', half to seedbed (3 Apr) and half in mid-May (23 May)	Nitro/EL
'Nitro-Chalk', all in mid-May (23 May)	Nitro/L
2. Rate of nitrogen fertiliser (kg N):	N RATE(2)
30	30
60	60
90	90
120	120

74/R/E/14

Basal applications: Manures: (0:20:20) at 190 kg combine drilled.
Weedkiller: Dicamba with mecoprop and MCPA ('Tetralex Plus' at
7.0 l in 220 l).

Seed: Julia, dressed with ethirimol, sown at 160 kg.

Cultivations, etc.: - Ploughed: 20 Nov, 1973. Spring-tine cultivated:
23 Nov. Power harrowed and seed sown: 4 Apr, 1974. Weedkiller
applied: 28 May. Combine harvested: 21 Aug. Previous crops:
Winter wheat 1972, fallow 1973.

NOTE: The percentage of N in the crop was determined at growth stages
3, 7 and 10.5.

Standard error per plot.

Grain, tonnes/hectare: 0.179 or 2.8% (26 d.f.)

74/R/B/14

TABLES OF MEANS

GRAIN: TONNES/HECTARE

N RATE(1)

	0	15	30	45	60	75	90	105	120	Mean
N FORM(1)										
Gold-N/E		6.37	6.57	6.44	6.44	6.49	6.24	6.52	6.39	6.43
Nitro/E		6.63	6.54	6.22	6.48	6.18	6.34	5.97	6.24	6.32
Mean	6.54	6.50	6.55	6.33	6.46	6.34	6.29	6.24	6.32	6.40

N RATE(2)

	30	60	90	120	Mean
N FORM(2)					
Nitro/EL	6.44	6.44	5.98	5.95	6.20
Nitro/L	6.48	6.49	6.06	5.83	6.22
Mean	6.46	6.46	6.02	5.89	6.21

STANDARD ERRORS OF DIFFERENCES

N FORM(1)	N RATE(1)	N FORM(1) N RATE(1)	N FORM(2)	N RATE(2)	N FORM(2) N RATE(2)
0.063	0.127	0.179	0.090	0.127	0.179
Between any of N RATE(1)	v	any of N RATE(2)		0.127	
Between any of N FORM(1)	x	any of N FORM(2)		0.179	
	v				
	N RATE(1)		N RATE(2)		

Grand mean 6.34

Mean D.M. % 83.8

74/R/B/14

STRAW: TONNES/HECTARE

N RATE(1)

	0	15	30	45	60	75	90	105	120	Mean
N FORM(1)										
Gold-N/E		4.20	4.41	3.79	3.90	4.11	3.97	4.05	4.00	4.05
Nitro/E		3.80	4.13	4.09	4.40	4.72	4.52	4.12	4.63	4.30
Mean	3.81	4.00	4.27	3.94	4.15	4.42	4.25	4.08	4.32	4.14

N RATE(2)

	30	60	90	120	Mean
N FORM(2)					
Nitro/EL	4.01	4.01	3.90	4.25	4.04
Nitro/L	3.86	3.93	4.30	4.15	4.06
Mean	3.94	3.97	4.10	4.20	4.05

GRAND MEAN 4.11

Mean D.M. % 89.2