

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1974

[Full Table of Content](#)



Beans

Beans, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 323 - 334 - **DOI:**
<https://doi.org/10.23637/ERADOC-1-119>

74/R/BE/1

SPRING BEANS

VIRUS CONTROL

Object: To study the effects of heat treatment of seed, roguing and control of weevil vectors on yield and incidence of broad bean stain (BBSV) and broad bean true mosaic (EBTMV) viruses in field beans - Whittlocks.

Sponsor: A.J. Cockbain.

Design: 4 randomised blocks of 5 plots.

Whole plot dimensions: 4.27 x 15.2. Area harvested: 0.00488.

Treatments: Virus control:	VRUSCONT
None	O
Seed heat treated (1 hour at 80°C), crop rogued	HR
Seed not heat treated, crop rogued	R
Seed heat treated, crop rogued and sprayed with methomyl	HRS
Seed not heat treated, crop rogued and sprayed with methomyl	RS

NOTES: (1) Plants with symptoms of BBSV or EBTMV were removed from R plots on 1, 8, 16, 21 and 31 May, 7, 14 and 20 June.
(2) Methomyl was applied on 15, 30 May, 20 June at 1.0 kg in 500 l.

Basal applications: Manures: (0:14:28) at 400 kg placement drilled.
Weedkiller: Simazine at 1.1 kg in 220 l. Insecticide: Demeton-s-methyl at 0.25 kg in 220 l.

Seed: Maris Bead, sown at 220 kg.

Cultivations, etc.: - Deep-tine cultivated: 19 and 20 Sept, 1973. Deep-tine cultivated: 26 Sept. Power harrowed: 27 Mar, 1974. Seed sown and spring-tine cultivated: 28 Mar. Simazine applied: 5 Apr. Insecticide applied: 13 June. Combine harvested: 26 Sept. Previous crops: Winter wheat 1972, barley 1973.

NOTE: Plant counts were made on 9 May. Damage by weevils (*Sitona* and *Apion*) was recorded on 9 May. Estimates of virus incidence were made on 7 and 20 June and 5 July. Seed samples were taken at harvest for assessment of virus infection.

Standard error per plot.

Grain, tonnes/hectare: 0.312 or 11.2% (12 d.f.)

74/R/BE/1

TABLES OF MEANS

GRAIN: TONNES/HECTARE

VRUSCOMP					Mean
O	HR	R	HRS	RS	
2.59	2.48	2.60	2.91	3.37	2.79

STANDARD ERROR OF DIFFERENCES

VRUSCOMP

0.221

Mean D.M. % 67.1

74/R/BE/2

SPRING BEANS

INSECTICIDE AND BENEFICIAL INSECTS

Object: To study the effects of a range of rates of dimethoate on beneficial insects, particularly predators and parasites of aphids and the yield of field beans - Whittlocks.

Sponsor: J.H. Stevenson.

Design: 5 randomised blocks of 6 plots.

Whole plot dimensions: 21.3 x 22.9. Area harvested: 0.00439.

Treatments: Rates of dimethoate (g a.i.)	DIMETH
None	0
26	26
52	52
105	105
210	210
420	420

- NOTES: (1) Treatments were applied on 8 July in 390 l.
(2) The application of treatments was deliberately delayed to allow aphid infestations and associated parasites and predators to establish. Weather conditions then delayed spraying until 8 July, by which time aphids had caused appreciable damage to all plots.

Basal applications: Manures: Chalk at 7.5 tonnes. (0:14:28) at 400 kg placement drilled. Weedkillers: Paraquat at 0.56 kg ion in 220 l, and simazine at 1.1 kg in 220 l.

Seed: Minor, sown at 220 kg.

Cultivations, etc.: Chalk applied: 6 Sept, 1973. Paraquat applied: 24 Oct. Chisel ploughed twice: 6 Dec and 10 Dec. Spring-tine cultivated: 23 Mar, 1974. Rotary harrowed, seed sown and spring-tine cultivated: 27 Mar. Simazine applied: 5 Apr. Combine harvested: 30 Sept. Previous crops: Winter wheat 1972, barley 1973.

74/R/EE/2

- NOTES: (1) Insect populations were sampled by sweep nets and water traps, aphid (*Aphis fabae*) counts were made.
 (2) There was evidence of a linear fertility trend across the site and yields adjusted for trend are presented.

Standard error per plot.

Grain, tonnes/hectare: 0.277 or 15.3% (19 d.f.)

TABLES OF MEANS

GRAIN: TONNES/HECTARE

DIMETH						Mean
0	26	52	105	210	420	
1.36	1.39	1.64	1.89	2.49	2.09	1.81

STANDARD ERROR OF DIFFERENCES

DIMETH

0.176

Mean D.M.% 69.7

74/R/BE/3

SPRING BEANS

VARIETIES AND VIRUSES

Object: To study the spread and effects on yield of broad bean stain and broad bean true mosaic viruses in different varieties of field bean - Long Hoos I/II.

Sponsor: A.J. Cockbain.

Design: 4 randomised blocks of 6 plots.

Whole plot dimensions: 6.40 x 15.2. Area harvested: 0.00488.

Treatments: Varieties (all healthy seed except where stated):	VARIETY
Herz Freya	Freya H
Maris Bead, seed infected with both viruses, two plots per block	Bead I
Maris Bead	Bead H
Minden	Minden H
Minor	Minor H

NOTE: Examination of seedlings showed that Herz Freya had 0.01% of plants with virus infection from the seed and Maris Bead, Healthy 0.04%. Maris Bead, Infected had 3.2%. Minden and Minor had none.

Basal applications: Manures: (0:14:28) at 400 kg placement drilled.
Weedkiller: Simazine at 1.1 kg in 220 l. Insecticide: D-meton-s-methyl at 0.25 kg in 450 l.

Seed: Sown at 220 kg.

Cultivations, etc.: - Ploughed: 19 Nov, 1973. Spring-tine cultivated: 8 and 9 Mar, 1974. Spring-tine cultivated: 26 Mar. Seed sown and spring-tine cultivated: 28 Mar. Weedkiller applied: 10 Apr. Insecticide applied: 19 June. Combine harvested: 26 Sept.

NOTE: Plant counts were made on 15 May. Seed-borne virus infection was recorded on 9 May and damage by Sitona and Apion weevils on 15 May. Incidence of viruses was assessed on 15 May, 13 June, 2 and 7 July. Estimates of the numbers of adult weevils were made on 13 June. Seed samples were taken at harvest to assess virus infection.

Standard error per plot.

Grain, tonnes/hectare: 0.447 or 12.7% (16 d.f.)

74/R/BE/3

TABLE OF MEANS

GRAIN: TONNES/HECTARE

VARIETY

Freya H	Bead I	Bead E	Minden H	Minor H	Mean
3.03	3.32	3.14	4.64	3.68	3.52

STANDARD ERRORS OF DIFFERENCES

VARIETY

Bead I v any of remainder
Between any of remainder

0.274
0.316

Mean D.M. % 72.7

74/R/BE/4

SPRING BEANS

SYNTHETIC PYRETHROID

Object: To study the effects of the synthetic pyrethroid NRDC 143 on the control of bean weevils and the yield of field beans - Long Hoos I and II.

Sponsor: J.H. Stevenson.

Design: 3 randomised blocks of 3 plots.

Whole plot dimensions: 8.53 x 9.14. Area harvested: 0.00293.

Treatments: Insecticides:	INSECTICIDE
None	None
Methomyl at 1.0 kg	Methomyl
NRDC 143 at 0.56 kg	NRDC 143

NOTE: Treatments were applied on 12 June in 390 l.

Basal applications: Manures: (0:14:26) at 400 kg placement drilled.
Weedkiller: Simazine at 1.1 kg in 220 l.

Seed: Minor, sown at 220 kg.

Cultivations, etc.: - Ploughed: 19 Nov, 1973. Spring-tine cultivated twice: 8 and 9 Mar, 1974. Spring-tine cultivated: 26 Mar. Seed sown and spring-tine cultivated: 27 Mar. Weedkiller applied: 10 Apr. Combine harvested: 26 Sept. Previous crops: Winter wheat 1972, barley 1973.

NOTE: Counts were made of weevils (Sitona and Apion) just before spraying and daily for eight days after.

Standard error per plot.

Grain, tonnes/hectare: 0.489 or 15.4% (4 d.f.)

74/R/BE/4

TABLE OF MEANS

GRAIN: TONNES/HECTARE

INSECTICIDE			Mean
None	Methomyl	NRIC 143	
1.89	3.74	3.89	3.17

STANDARD ERROR OF DIFFERENCES

INSECTICIDE

0.399

Mean D.M. % 75.9

74/R/BE/6

SPRING BEANS

CONTROL OF WEEVILS

Object: To study the effects of several insecticides on the yield and control of weevils and weevil-transmitted viruses of field beans - Dell Piece.

Sponsors: A.J. Cockbain, P. Etheridge.

Design: 3 randomised blocks of 8 plots.

Whole plot dimensions: 8.53 x 12.2. Area harvested: 0.00390.

Treatments: All combinations of insecticides:

1. Sprays to foliage	SPRAY
None	None
Fenitrothion at 0.75 kg	Fenitro
Malathion at 1.0 kg	Malathio
Methomyl at 1.0 kg	Methomyl
2. Granules to foliage	GRANULE
None	None
Phorate at 1.0 kg	Phorate

NOTE: Sprays, in 500 l, and granules were applied on 15 May, 30 May and 19 June.

Basal applications: Fures: (0:14:28) at 400 kg. Weedkiller: Simazine at 1.1 kg in 220 l. Insecticides: Demeton-s-methyl at 0.25 kg in 220 l.

Seed: Maris Bead, sown at 220 kg.

Cultivations, etc.: - Ploughed: 3 Jan, 1974. Spring-tine cultivated twice: 27 Mar. Rotary harrowed: 1 Apr. Seed sown and spring-tine cultivated: 2 Apr. Simazine applied: 5 Apr. Demeton-s-methyl applied: 13 June. Combine harvested: 26 Sept. Previous crops: Follow 1972, winter wheat 1973.

74/R/EE/6

NOTE: Amounts of damage by weevils (*Agrion* and *Sitona*) were recorded on 9 May and 25 May. Numbers of adults were estimated on 12 June. Incidence of viruses was assessed on 25 April, 12 June, 4 and 23 July and samples of seed were taken at harvest to assess virus infection.

Standard error per plot.

Grain, tonnes/hectare: 0.472 or 12.1% (14 d.f.)

TABLES OF MEANS

GRAIN: TONNES/HECTARE

	SPRAY				Mean
	None	Fenitro	Malathio	Methomyl	
GRANULE:					
None	3.55	3.84	4.41	3.45	3.82
Phorate	4.42	3.58	3.89	4.10	4.00
Mean	3.98	3.71	4.15	3.78	3.91

STANDARD ERRORS OF DIFFERENCES

GRANULE	SPRAY	GRANULE SPRAY
0.193	0.273	0.385

Mean D.M.% 65.2

74/R/BE/7

SPRING BEANS

CONTROL OF STEM EELWORM

Object: To study the effects of a range of rates of aldicarb, applied in the row, on yield and the control of stem eelworm (*Ditylenchus dipsaci*) in a seed-infested stock of field beans - Garden Plot 7.

Sponsor: D.J. Hooper.

Design: 4 randomised blocks of 4 plots.

Whole plot dimensions: 1.52 x 9.14. Area harvested: 0.00139.

Treatments: Rates of aldicarb (kg):	ALDICARB
None	0
1	1
2	2
4	4

NOTE: Aldicarb was applied in bands approx. 7.5 cm wide over the seed in rows spaced 50 cm apart.

Basal applications: Manures: (0:20:20) at 1000 kg. Weedkiller: Simazine at 0.84 kg in 340 l. Insecticide: Demeton-s-methyl at 0.25 kg in 340 l.

Seed: Minor, sown at 220 kg.

Cultivations, etc.: - PK applied: 20 Nov, 1973. Ploughed: 11 Dec. Rotary cultivated: 1 Apr, 1974. Seed sown and treatments applied: 2 Apr. Weedkiller applied: 5 Apr. Insecticide applied: 20 June and 19 July. Combine harvested: 30 Sept. Previous crops: Potatoes 1972, fallow 1973.

NOTES: Number of stems infested with stem eelworm were assessed in mid August. Samples of seed were taken at harvest to assess incidence of stem eelworm.

Standard error per plot.

Grain, tonnes/hectare: 0.534 or 11.7% (9 d.f.)

74/R/BE/7

TABLE OF MEANS

GRAIN: TONNES/HECTARE

ALDICARB

0	1	2	4	Mean
4.52	4.48	4.53	4.68	4.55

STANDARD ERROR OF DIFFERENCES

ALDICARB

0.377

Mean D.M. % 71.3