

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1974

[Full Table of Content](#)



### 74/R/BE/3 Varieties and Viruses - Beans

74/R/BE/3 Varieties and Viruses - Beans, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 327 - 328 - DOI: <https://doi.org/10.23637/ERADOC-1-119>

74/R/BE/3

SPRING BEANS

VARIETIES AND VIRUSES

Object: To study the spread and effects on yield of broad bean stain and broad bean true mosaic viruses in different varieties of field bean - Long Hoos I/II.

Sponsor: A.J. Cockbain.

Design: 4 randomised blocks of 6 plots.

Whole plot dimensions: 6.40 x 15.2. Area harvested: 0.00488.

Treatments: Varieties (all healthy seed except where stated):	VARIETY
Herz Freya	Freya H
Maris Bead, seed infected with both viruses, two plots per block	Bead I
Maris Bead	Bead H
Minden	Minden H
Minor	Minor H

NOTE: Examination of seedlings showed that Herz Freya had 0.01% of plants with virus infection from the seed and Maris Bead, Healthy 0.04%. Maris Bead, Infected had 3.2%. Minden and Minor had none.

Basal applications: Manures: (0:14:28) at 400 kg placement drilled.  
Weedkiller: Simazine at 1.1 kg in 220 l. Insecticide: D-meton-s-methyl at 0.25 kg in 450 l.

Seed: Sown at 220 kg.

Cultivations, etc.: - Ploughed: 19 Nov, 1973. Spring-tine cultivated: 8 and 9 Mar, 1974. Spring-tine cultivated: 26 Mar. Seed sown and spring-tine cultivated: 28 Mar. Weedkiller applied: 10 Apr. Insecticide applied: 19 June. Combine harvested: 26 Sept.

NOTE: Plant counts were made on 15 May. Seed-borne virus infection was recorded on 9 May and damage by Sitona and Apion weevils on 15 May. Incidence of viruses was assessed on 15 May, 13 June, 2 and 7 July. Estimates of the numbers of adult weevils were made on 13 June. Seed samples were taken at harvest to assess virus infection.

Standard error per plot.

Grain, tonnes/hectare: 0.447 or 12.7% (16 d.f.)

74/R/BE/3

TABLE OF MEANS

GRAIN: TONNES/HECTARE

VARIETY

Freya H	Bead I	Bead E	Minden H	Minor H	Mean
3.03	3.32	3.14	4.64	3.68	3.52

STANDARD ERRORS OF DIFFERENCES

VARIETY

Bead I v any of remainder  
Between any of remainder

0.274  
0.316

Mean D.M. % 72.7