

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1974

[Full Table of Content](#)



74/R/P/1 and 74/W/P/1 Varieties - Potatoes

74/R/P/1 and 74/W/P/1 Varieties - Potatoes, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 335 - 338 - DOI: <https://doi.org/10.23637/ERADOC-1-119>

74/R/P/1 and 74/W/P/1

POTATOES

VARIETIES

Object: To study the yield and susceptibility to diseases of newer varieties of potatoes - Rothamsted (R) Pastures and Woburn (W) Lansome III.

Sponsors: J.R. Moffatt, G.A. Hide.

Design: Pastures (R): 3 randomised blocks of 7 plots
Lansome III (W): 3 randomised blocks of 5 plots

Whole plot dimensions: 2.85 x 15.2. Area harvested: 0.00217.

Treatments: Varieties:-

	VARIETY
Desiree, Pastures (R) only	Desiree
King Edward, Pastures (R) only	Edward
Maris Piper	Piper
Pentland Crown	Crown
Pentland Ivory	Ivory
Stormont Enterprise	Enterprise
Ulster Lancer	Lancer

Basal applications:

Pastures (R): Manures: (13:13:20) at 1510 kg. Weedkillers: Linuron at 1.2 kg plus paraquat at 0.42 kg ion in 450 l. Fungicide and insecticide: Mancozeb at 1.3 kg plus demeton-s-methyl at 0.25 kg in 450 l. Fungicide: Mancozeb at 1.3 kg in 450 l.

Lansome III (W): Manures: (13:13:20) at 1880 kg. Weedkillers: Linuron at 1.2 kg plus paraquat at 0.28 kg ion in 280 l. Fungicide with insecticide: Mancozeb at 1.3 kg plus demeton-s-methyl at 0.25 kg in 450 l. Fungicide: Mancozeb at 1.3 kg in 450 l.

Cultivations, etc.:-

Pastures (R): Ploughed: 23 Nov, 1973. Spring-tine cultivated: 17 Apr, 1974. NPK applied: 24 Apr. Spike rotary cultivated, potatoes planted: 29 Apr. Weedkiller applied: 21 May. Grubbed: 19 June. Rotary ridged: 21 June. Fungicide with insecticide applied: 12 July. Fungicide applied: 2 Aug. Haulm mechanically destroyed: 10 Sept. Sprayed with undiluted BOV at 220 l: 16 Sept. Lifted: 25 Oct. Previous crops: Barley 1972, spring beans 1973.

74/R/P/1 and 74/W/P/1

Lansome III (W): Subsoiled, tines 140 cm apart, 50 cm deep:
12-14 Nov, 1973. Ploughed: 14 Dec. Spring-tine cultivated:
10 Apr, 1974. NPK applied: 16 Apr. Rotary cultivated, potatoes
planted: 18 Apr. Weedkiller applied: 16 May. Rotary ridged:
11 June. Fungicide with insecticide applied: 18 July. Fungicide
applied: 7 Aug. Haulm mechanically destroyed: 12 Sept. Sprayed
with undiluted BOV at 170 l: 18 Sept. Lifted: 17 Oct. Previous
crops: Barley 1972, beans 1973.

NOTE: Tubers were graded into six sizes. Incidence of *Rhizoctonia solani*
and common scab on the produce was assessed.

Standard errors per plot. Total tubers, tonnes/hectare:

Pastures (R): 2.94 or 5.5% (12 d.f.)
Lansome III (W): 2.72 or 4.8% (8 d.f.)

74/R/P/1 and 74/W/P/1

TABLES OF MEANS

PASTURES (R)

VARIETY							Mean
Desiree	Edward	Piper	Crown	Ivory	Entprise	Lancer	
TOTAL TUBERS: TONNES/HECTARE							
50.6	56.1	58.0	53.4	48.8	58.4	46.5	53.1

STANDARD ERROR OF DIFFERENCES

VARIETY

2.40

PERCENTAGE WARE: 4.44 CM (1.75 INCH) RIDDLE

86.3	88.1	90.8	95.7	94.2	88.9	92.8	90.9
------	------	------	------	------	------	------	------

74/R/P/1 and 74/W/P/1

LANSIME III (W)

VARIETY					
Piper	Crown	Ivory	Enterprise	Lancer	Mean
TOTAL TUBERS: TONNES/HECTARE					
57.4	59.9	51.5	63.2	51.9	56.8

STANDARD ERRORS OF DIFFERENCES

VARIETY

2.22

PERCENTAGE WARE: 4.44 CM (1.75 INCH) RIDDLE

86.8	93.7	90.6	82.5	88.6	88.4
------	------	------	------	------	------