Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1974



Full Table of Content

74/R/RN/7 Residual Phosphate - Potatoes, Barley, Wheat, Ley

74/R/RN/7 Residual Phosphate - Potatoes, Barley, Wheat, Ley, Rothamsted Research (1975) Yields Of The Field Experiments 1974, pp 95 - 99 - DOI: https://doi.org/10.23637/ERADOC-1-119

RESIDUAL PHOSPHATE

Object: Originally to study the fresh and residual effects of phosphate fertiliser on yields of three arable crops grown in rotation. Since 1974 the effects on ley and on yield and pathogens of continuous wheat are also studied. Great Field IV and Sawyers I.

Sponsors: G.E.G. Mattingly, D.B. Slope.

The fifteenth year, potatoes, barley, ley (Gt. Field IV). Potatoes, barley, wheat (Sawyers I).

For previous years see 'Details' 1967, 68/B/5(t), 69/R/RN/7, 70/R/RN/7(t) and 71-73/R/RN/7.

Design: Great Field IV: 1 randomised block of 12 plots for each crop.

Sawyers I: 2 randomised blocks of 12 plots for each crop.

Whole plot dimensions:-

Great Field IV: 4.27 x 18.3. Area harvested: potatoes and barley - 0.00520, ley - 0.00186.

Sawyers I: 4.27 x 20.1. Area harvested: potatoes and barley - 0.00572, wheat - 0.00615.

Treatments: Rates and frequency of applying phosphate: P205

| None | 0 |
|---|------|
| Annual dressings, kg P205:- | |
| 29 | ANN |
| 57 | ANN |
| 115 | ANN |
| 172 | ANN |
| Triennial dressings, kg P205 (last applied 1972):- | |
| 86 | TRI |
| 172 | TRI |
| Six-yearly dressings, kg P205 (last applied 1973):- | |
| | SIX |
| 688 | SIX |
| 1032 | SIX |
| Single dressing, kg P205 applied in 1959:- | |
| 376 as Gafsa rock phosphate 376 | G(1) |
| 376 as superphosphate 376 | s(1) |

- NOTES: (1) Since 1974 the original rotation of potatoes, barley, swedes on both fields has been changed. Blocks after barley are sown to ley on Gt. Field IV, to continuous wheat on Sawyers I.
 - (2) Since 1960 all phosphate has been applied as superphosphate.
 - (3) The six-yearly dressings were applied half in autumn before ploughing, half in spring.
 - (4) For a fuller record of treatments see 'Details' etc.

Standard applications:

Potatoes: Manures: N at 250 kg as 'Nitro-Chalk'. K20 at 250 kg as sulphate of potash. Weedkillers: Linuron at 1.2 kg and paraquat at 0.42 kg ion in 450 l. Fungicide: Mancozeb at 1.3 kg in 450 l on two occasions. Insecticide: Demeton-s-methyl at 0.25 kg applied with mancozeb on the first occasion.

Barley: Manures: (25:0:16) at 390 kg combine drilled. Weedkillers: Dicamba with mecoprop and MCPA ('Tetralex Plus' at 7.0 l in 220 l).

Winter wheat (Sawyers I only): Manures: Ground chalk at 2.9 tonnes. K20 at 87 kg as muriate of potash. N at 125 kg as 'Nitro-Chalk'. Weedkiller: Dicamba with mecoprop and MCPA ('Banlene Plus' at 5.6 1 in 370 1).

Ley (Gt.Field IV only): Manures: Chalk at 2.9 tonnes. N at 60 kg as 'Nitro-Chalk' and K20 at 120 kg as muriate of potash in seedbed. N at 100 kg for each cut. Weedkiller: Dicamba with mecoprop and MCPA ('Tetralex Plus' at 7.0 l in 220 l).

Seed: Potatoes: Majestic, Foundation Stock.

Barley: Julia, dressed with ethirimol, sown at 160 kg.

Wheat: Cappelle, sown at 200 kg.

Grass Clover mixture: 11.2 kg S215 Meadow Fescue, 3.4 kg S48 Timothy,

2.2 kg S51 Timothy, 3.4 kg S26 Cocksfoot, 1.7 kg N.Z. White Clover,

O.6 kg Wild White clover. Sown at 22.5 kg.

Cultivations, etc.:- (both fields where applicable).

Potatoes: Ploughed: 19 Nov, 1973. N, P and K applied, rotary cultivated, seed planted: 22 Apr, 1974. Weedkiller applied: 16 May. Grubbed: 20 June. Rotary ridged: 24 June. Fungicide and insecticide applied: 11 July. Fungicide applied: 2 Aug. Haulm destroyed mechanically: 10 Sept. Sprayed with undiluted BOV at 220 1: 17 Sept. Lifted: 15 Oct (Sawyers I), 21 Oct (Great Field IV).

Barley: Ploughed: 11 Oct, 1973. Spring-tine cultivated and P applied: 8 Mar, 1974. Power harrowed: 27 Mar. Seed sown and NK applied: 28 Mar. Weedkiller applied: 21 May. Combine harvested: 20 Aug. Wheat: Chalk applied: 11 Sept, 1973. Ploughed: 11 Oct. P and K

applied: 23 Oct. Power harrowed and seed sown: 25 Oct. N applied: 16 Apr, 1974. Weedkiller applied: 7 May. Combine harvested: 29 Aug.

Ley: Chalk applied: 11 Sept, 1973. Ploughed: 13 Sept. Rotary harrowed twice, N and K applied, power harrowed and seed sown: 14 Sept. P applied: 17 Sept. Weedkiller applied: 28 May, 1974. N applied: 5 June, 5 Aug. Cut: 25 July, 23 Dec.

Standard errors per plot. Sawyers I.

Potatoes, total tubers: tonnes/hectare: 2.41 or 7.0% (11 d.f.)

Barley, grain: tonnes/hectare: 0.274 or 5.7% (11 d.f.)

Wheat, grain: tonnes/hectare: 0.484 or 7.2% (11 d.f.)

TABLES OF MEANS

POTATOES

| | | | | (1.5 INCH) RIDDLE Sawyers I |
|---|--|--|--|--|
| P205 0 29 ANN 57 ANN 115 ANN 172 ANN 86 TRI 172 TRI 344 SIX 688 SIX 1032 SIX 376 G(1) 376 S(1) | 16.9 21.7 18.8 35.4 39.1 18.6 25.6 33.9 39.1 40.5 15.9 17.6 | 19.8 30.8 36.7 47.0 47.6 24.8 31.5 41.2 47.3 47.8 19.2 18.8 | 96.8 96.9 95.8 98.3 98.0 95.1 97.1 98.3 98.6 97.8 97.9 | 97.2 97.6 98.2 98.0 98.0 96.2 96.6 98.1 98.1 98.7 96.2 96.1 |
| Mean | 26.9 | 34.4 | 97•3 | 97.4 |

Sawyers I only - TOTAL TUBERS

STANDARD ERROR OF DIFFERENCES

P205

2.41

BARLEY

| | GRAIN: TONNES, Great Field IV | | STRAW: TONNE Great Field IV | SAWYERS I |
|--|--|--|--|--|
| P205 0 29 ANN 57 ANN 115 ANN 172 ANN 172 TRI 344 SIX 688 SIX 1032 SIX 376 G(1) 376 S(1) | 3.18 3.98 4.71 3.47 5.05 3.76 4.09 3.65 4.23 3.58 4.55 3.64 | 4.06 4.75 4.86 5.15 5.38 4.25 5.06 4.97 5.53 4.22 4.47 | 2.04 2.61 4.06 3.29 4.67 3.10 3.96 3.55 2.73 3.10 3.63 4.02 | 2.58 2.55 2.85 3.00 3.38 2.82 2.81 2.68 3.03 3.19 2.49 2.72 |
| Mean | 3.99 | 4.81 | 3.40 | 2.84 |

Sawyers I only - GRAIN

STANDARD ERROR OF DIFFERENCES

P205

0.274

Mean D.M. % 73.7 74.5 68.2 86.3

| | Great Field IV LEY DRY MATTER: TONNES/HECTARE | | | WHI | ers I EAT HECTARE |
|---|--|--|--|--|--|
| | 1st cut | 2nd cut | Total of 2 cuts | Grain | Straw |
| P205 | | | | | |
| 29 ANN 57 ANN 115 ANN 172 ANN 86 TRI 172 TRI 344 SIX 688 SIX 1032 SIX 376 G(1) 376 S(1) | 3.70 3.63 4.07 3.71 3.99 4.10 4.10 4.23 3.91 3.50 4.52 3.83 | 3.80 3.56 3.90 4.27 3.73 3.76 3.45 3.57 3.93 4.18 3.55 3.65 | 7.50 7.19 7.97 7.98 7.72 7.86 7.55 7.80 7.68 8.07 7.48 | 4.80 7.48 7.01 6.91 7.12 6.17 7.11 7.41 6.68 7.39 6.17 6.65 | 3.14 4.84 4.46 3.99 4.55 3.73 4.20 4.32 4.41 4.69 3.73 3.80 |
| Mean | 3.94 | 3.78 | 7.72 | 6.74 | 4.15 |

Sawyers I only - Wheat grain

STANDARD ERROR OF DIFFERENCES

P205

0.484

Mean D.N. % 22.9 23.9 23.4 84.1 91.0