

Rothamsted Experimental Station
Harpden
LAWES AGRICULTURAL TRUST

**RESULTS
OF THE
FIELD
EXPERIMENTS**

1939 - 1947

VOLUME 1

Rothamsted Experimental Station

Harpenden

Lawes Agricultural Trust

RESULTS

of the

FIELD

EXPERIMENTS

1939 - 1947

Vol. I. Classical and Long-Term Experiments

The summaries given in this report are similar to those given in the appendices to the Annual Reports of the Station before the war. Only experiments conducted at Rothamsted and Woburn are included. The design and supervision of these experiments are the responsibility of the Field Plots Committee (Members during the period covered by this report: E.M.Crowther (Chairman), H.V.Garner (Secretary), D.J.Finney, J.R.Moffatt, R.G.Warren, D.J.Watson, F.Yates).

Price: 10/-

INDEX

Classical Experiments*

Broadbalk	Wheat	A/1
Hoosfield	Barley	A/2
Hoosfield	Wheat after fallow	A/3
Agdell	Rotation	A/4
Barnfield	Mangolds and sugar beet	A/5
Park Grass	Hay	A/6
Woburn Stackyard	Wheat and barley	A/7

Long Term Experiments

2-course Rotation	Rothamsted	Ba/1
3-course Rotation	Rothamsted	Ba/2
4-course Rotation	Rothamsted	Ba/3
6-course Rotation	Rothamsted and Woburn	Ba/4
Long Period Cultivation	Rothamsted	Bb/1
Deep Cultivation Rotation	Rothamsted	Bc/1
Highfield Grazing	Rothamsted	Bd/1
Green Manuring	Woburn	Be/1
Ley and Arable Rotations	Woburn	Bf/1
Market Garden	Woburn	Bg/1

Miscellaneous Data

Chemical Analyses of Manures	E/1
------------------------------	-----

*All except A/7 at Rothamsted.

WHEAT - BROADBALK

The first experimental crop of wheat on Broadbalk was sown in the autumn of 1843 for harvesting in 1844, and wheat has been grown each year ever since. The plot treatments varied until 1852 when the present system of manuring was established.

The general purpose of the experiment was to determine over a long period of years the manurial requirements of wheat, testing dung alone and a number of combinations of the ash constituents of crops (compounds of phosphorus, potassium, sodium and magnesium), together with several forms of nitrogen compounds. One plot running the whole length of the field was assigned to each of the treatments tested. Several varieties, some of them now out of cultivation, have been used during the experiment, but since 1899 Squareheads Master or the very similar Red Standard has been sown.

Weeds have always been a problem on Broadbalk, but in spite of this, continuous cropping was maintained until the harvest of 1925, except that half of every plot was fallowed in 1904 and the remainder in 1905, and this process was repeated in 1914 and 1915. In 1926 it was decided that more systematic methods must be tried and a system of regular bare fallowing was introduced. The field was divided into five equal sections so that every year a part of each of the original $\frac{1}{2}$ acre plots could be under fallow and the remainder under crop. From 1926-29 two sections were cropped yearly, leaving three fallow; in 1930 the whole field was cropped and then in 1931 the present arrangement was begun under which one section of the field is fallowed every year. In each year therefore, the yields of every treatment are obtained for the 1st, 2nd, 3rd and 4th years after fallow.

Since 1943 it has been necessary to hand weed wild oats before harvesting.

Treatments
Symbols and amounts per acre

O	Unmanured
N ₁ , N ₂ , N ₃	Sulphate of ammonia, 43 lb.N, 86 lb.N, 129 lb.N
N _{1'} , N _{2'}	Nitrate of soda, 43 lb.N, 86 lb.N
P	Superphosphate, 65 lb.P ₂ O ₅
K	Sulphate of potash, 98 lb.K ₂ O
S	Sulphate of soda, 366 lb.
M	Sulphate of magnesia, 280 lb.
C	Complete mineral manure, consisting of superphosphate (65 lb.P ₂ O ₅), sulphate of potash (98 lb.K ₂ O), sulphate of soda (100 lb.) and sulphate of magnesia (100 lb.)
C'	As C but without the superphosphate
D	Farmyard manure, 14 tons
R	Rape cake (castor bean meal from 1940 onwards), 1889 lb.

Dung is ploughed-in in autumn. Rape cake and minerals are applied in autumn on the seed bed. 21.5 lb. N per acre of the N₁, N₂ and N₃ treatments is applied in autumn and the remainder of the dressing in spring, except that Plot 15 receives its full dressing in autumn. Nitrate of soda is applied in spring, there being two applications at intervals of a month on plot 16. Plot 2A was unmanured before 1885.

The experiment is discussed, and early departures from the present manuring scheme are described by Sir John Russell and D.J. Watson in "The Rothamsted Field Experiments in the growth of wheat", Imp. Bureau Soil Sci. Tech. Comm. No.40.

Investigations by the Soil Microbiology Department are described by B.N. Singh, "The effect of artificial fertilizers and dung on the numbers of amoebae in Rothamsted soils", J. Gen. Microbiol. 3, (1949), 204.

Weed surveys have been made annually by the Botany Department.

Wheat - Broadbalk

The present fallowing cycle and the preceding fallows are shown in the diagram below. (C = crop, F = fallow). The sections (I to V) are numbered in order from the upper (western) end of the field. Preparatory to the first fallow the field was harvested in five separate sections (1925).

Year	I	II	III	IV	V	Year	I	II	III	IV	V
1926	F	F	F	C	C	1931, -36, -41, -46	F	C	C	C	C
1927	F	F	F	C	C	1932, -37, -42, -47	C	F	C	C	C
1928	C	C	F	F	F	1933, -38, -43, -48	C	C	C	C	F
1929	C	C	F	F	F	1934, -39, -44, -49	C	C	C	F	C
1930	C	C	C	C	C	1935, -40, -45, -50	C	C	F	C	C

Plot 20 extends over sections I and II only.

Crop Notes

Crop Year	Date sown	Date harvested	Variety
1939	Oct. 26	Aug. 18	Red Standard
1940	Oct. 25	Aug. 8	Squareheads Master
1941	Oct. 25	Aug. 20	Squareheads Master
1942	Oct. 24	Aug. 15	Red Standard
1943	Oct. 15	Aug. 4	Red Standard
1944	Oct. 16	Aug. 5	Red Standard
1945	Nov. 8	Aug. 12	Red Standard
1946	Oct. 20	Aug. 16	Squareheads Master 13/4
1947	Oct. 31	Aug. 8	Squareheads Master 13/4

In 1941 there was a considerable amount of lodging, particularly on plots 2A, 2B, 8, 12, 14 and 16.

Years after fallow	Total Grain: cwt. per acre					Total Straw: cwt. per acre				
	1	2	3	4	Mean	1	2	3	4	Mean

96th season, 1939

Plot	Section	V	II	I	III		V	II	I	III	
2A	D	24.1	18.5	15.7	19.7	19.5	62.7	40.1	43.0	42.8	47.2
2B	D	26.1	23.5	15.7	18.1	20.8	64.2	51.6	48.9	47.8	53.1
3	O	16.7	7.7	8.7	8.9	10.5	29.7	17.0	16.4	19.1	20.6
5	C	18.6	9.9	9.3	7.8	11.4	36.1	24.9	21.9	19.2	25.5
6	N ₁ C	22.5	13.8	10.9	11.8	14.8	45.3	33.9	29.5	54.5	40.8
7	N ₁ C	26.4	19.3	13.5	15.8	18.8	55.8	43.4	39.8	37.7	44.2
8	N ₂ C	20.8	21.6	11.8	12.5	16.7	59.0	54.7	62.9	52.9	57.4
9	N ₃ 'C	24.1	16.2	11.3	14.0	16.4	53.5	35.2	28.5	29.8	36.8
10	N ₂	20.2	19.6	14.4	17.1	17.8	38.8	35.6	33.4	32.4	35.0
11	N ₂ P	18.6	18.9	13.4	15.8	16.7	36.6	35.7	28.8	30.4	32.9
12	N ₂ PS	21.2	20.4	14.8	16.4	18.2	42.4	40.1	32.7	32.3	36.9
13	N ₂ PK	23.1	17.5	14.4	15.3	17.6	55.2	42.1	33.8	35.2	41.6
14	N ₂ PM	24.6	19.9	14.1	17.1	18.9	52.7	35.3	29.1	32.4	37.4
15	N ₂ *C	17.8	16.5	12.7	12.8	15.0	37.1	34.3	28.8	29.3	32.4
16	N ₂ 'C	25.4	18.5	11.8	13.5	17.3	69.4	48.9	43.6	44.9	51.7
17	N ₂ and C in (C) alternate (N ₂) years	13.7	10.1	5.6	7.9	9.3	31.6	24.4	18.8	21.8	24.2
18		22.5	17.1	14.6	13.5	16.9	43.8	38.1	34.8	25.9	35.6
19	R	21.8	17.5	12.4	16.2	17.0	42.6	32.6	27.1	29.2	32.9
20	N ₂ C'	-	19.3	13.2	-	16.2	-	42.6	32.7	-	37.6

97th season, 1940

Plot	Section	IV	V	II	I		IV	V	II	I	
2A	D	33.1	31.9	27.7	15.4 ⁺	27.0	55.7	44.6	42.2	39.3	45.4
2B	D	37.0	29.0	32.3	23.8 ⁺	30.5	57.8	42.4	46.5	43.3	47.5
3	O	21.5	13.6	12.6	15.4	15.8	28.6	18.7	16.7	18.3	20.6
5	C	25.5	12.7	14.4	16.3	17.2	38.8	17.0	17.9	22.6	24.1
6	N ₁ C	30.5	17.1	17.8	20.8	21.6	47.3	25.6	23.6	28.2	31.2
7	N ₁ C	34.6	21.1	24.4	26.7	26.7	61.7	35.2	37.7	42.4	44.2
8	N ₂ C	32.2	27.1	27.4	29.9	29.2	61.8	51.7	47.5	55.3	54.1
9	N ₃ 'C	31.7	19.0	20.9	26.3	24.5	50.7	30.3	33.0	43.4	39.4
10	N ₂	14.8	18.4	17.5	19.8	17.6	26.0	24.3	21.9	27.8	25.0
11	N ₂ P	27.9	19.0	20.0	20.9	22.0	41.4	31.5	25.3	30.0	32.0
12	N ₂ PS	32.9	20.3	22.9	25.5	25.4	48.1	29.8	30.9	36.2	36.2
13	N ₂ PK	35.7	17.8	22.5	24.0	25.0	59.7	33.3	33.6	37.7	41.1
14	N ₂ PM	34.4	18.7	22.1	24.3	24.9	54.9	30.2	30.4	36.2	37.9
15	N ₂ *C	32.6	15.1	22.6	25.8	24.0	51.8	25.1	33.3	38.3	37.1
16	N ₂ 'C	33.8	26.8	29.6	27.6	29.4	60.8	40.1	43.0	45.3	47.3
17	N ₂ and C in (N ₂) alternate (C) years	33.7	25.2	26.2	28.6	28.4	56.4	39.1	39.3	42.2	44.2
18		24.4	11.0	11.5	12.2	14.8	33.5	17.1	17.3	15.3	20.8
19	R	34.2	19.0	21.0	20.7	23.7	52.3	28.5	29.3	31.6	35.4
20	N ₂ C'	-	-	20.4	23.0	21.7	-	-	25.4	33.4	29.4

Wheat - Broadbalk

A/1.5

Years after fallow		Total Grain: cwt. per acre					Total Straw [†] : cwt. per acre				
		1	2	3	4	Mean	1	2	3	4	Mean
98th season, 1941											
Plot	Section	III	IV	V	II		III	IV	V	II	
2A	D	16.8	13.7	14.4	9.5	13.6	40.1	26.6	25.0	25.4	29.3
2B	D	19.2	16.0	16.2	9.1	15.1	41.8	30.8	28.3	25.7	31.6
3	O	14.3	4.5	6.6	4.7	7.5	23.9	7.6	8.7	7.8	12.0
5	C	14.9	5.0	7.2	5.9	8.2	23.3	10.3	12.4	10.0	14.0
6	N ₁ C	15.7	9.6	9.8	7.6	10.7	29.1	16.1	15.2	14.0	18.6
7	N ₂ C	17.1	14.2	12.6	11.3	13.3	39.7	31.7	27.2	22.2	30.2
8	N ₃ C	20.9	14.3	13.1	16.2	16.1	46.2	37.6	32.1	35.0	37.7
9	N ₁ 'C	17.3	12.1	11.9	11.6	13.4	34.6	22.8	22.2	22.8	25.6
10	N ₂	13.8	14.3	12.5	12.2	13.2	26.6	27.2	19.6	17.4	22.7
11	N ₂ P	16.7	14.5	10.9	11.1	13.3	30.2	26.1	19.4	18.0	23.4
12	N ₂ PS	18.2	16.0	14.1	13.8	15.5	35.0	27.0	23.6	20.2	26.4
13	N ₂ PK	19.3	14.5	9.9	11.3	13.3	28.1	28.0	22.1	18.6	24.2
14	N ₂ PM	19.7	16.7	11.8	12.3	15.1	34.4	28.3	22.3	18.4	25.8
15	N ₂ *C	18.4	12.0	5.6	9.8	11.4	36.7	20.1	11.7	14.2	20.7
16	N ₂ 'C	21.7	15.9	14.9	15.7	17.0	49.7	36.2	31.7	31.0	37.2
17	N ₂ and C in (C) alternate years (N ₂)	15.6	5.7	4.0	4.1	7.4	35.5	8.4	6.2	6.4	14.1
18		18.8	12.2	12.4	10.1	13.4	31.8	21.3	21.6	18.0	23.2
19	R	18.0	13.5	9.3	9.2	12.5	29.9	22.7	13.3	11.6	19.4
20	N ₂ C'					11.6	11.6			19.6	19.6
99th season, 1942											
Plot	Section	I	III	IV	V		I	III	IV	V	
2A	D	20.7	23.7	24.5	27.9	24.2	52.2	43.2	45.1	43.6	46.0
2B	D	19.6	26.8	28.6	26.6	25.4	53.5	43.2	45.1	42.4	47.3
3	O	15.7	10.1	15.9	15.0	14.2	22.7	13.5	20.1	18.6	18.7
5	C	21.6	11.7	16.7	17.0	16.8	34.7	15.5	23.7	20.7	23.6
6	N ₁ C	25.1	16.3	18.6	21.5	20.4	42.3	22.7	26.7	29.4	30.3
7	N ₂ C	28.9	23.0	27.1	26.4	26.4	43.3	33.9	37.1	36.4	39.0
8	N ₃ C	25.8	25.7	28.8	24.8	26.3	52.4	39.8	41.6	39.7	43.4
9	N ₁ 'C	27.1	20.5	17.8	18.0	20.8	45.2	29.1	24.3	24.0	30.6
10	N ₂	22.9	23.4	18.5	17.2	20.5	32.2	28.6	22.8	21.4	26.2
11	N ₂ P	18.2	21.3	17.9	16.6	18.5	29.2	28.2	23.1	31.2	27.9
12	N ₂ PS	24.5	24.4	25.4	21.0	23.8	37.0	36.3	29.7	26.3	32.3
13	N ₂ PK	23.0	22.7	23.3	23.8	24.5	48.5	31.3	33.1	32.9	36.4
14	N ₂ PM	25.0	22.4	25.0	22.3	23.7	35.4	32.2	32.2	28.6	32.1
15	N ₂ *C	28.6	20.9	23.7	21.2	23.6	49.8	33.5	33.5	32.6	37.4
16	N ₂ 'C	28.1	25.6	24.3	23.5	25.4	46.6	36.2	35.5	33.9	38.0
17	N ₂ and C in (N ₂) alternate years (C)	23.8	22.2	25.8	24.0	25.2	43.4	31.6	37.2	34.7	36.7
18		21.6	10.4	11.5	13.6	14.3	29.6	14.6	15.5	22.1	20.4
19	R	31.1	22.2	23.0	24.7	25.2	42.7	29.7	33.1	33.3	34.7
20	N ₂ C'	24.0				24.0	35.0			35.0	

[†] Includes straw, cavings and chaff. * Sulphate of ammonia applied in autumn.

Years after fallow		Total Grain: cwt. per acre					Total Straw [†] : cwt. per acre				
		1	2	3	4	Mean	1	2	3	4	Mean
100th season, 1943											
Plot	Section	II	I	III	IV		II	I	III	IV	
2A	D	23.5	20.0	15.9	6.6	17.7	62.5	49.8	41.0	46.8	50.0
2B	D	29.7	21.4	23.1	14.4	22.2	65.9	57.5	48.6	49.4	55.4
3	O	25.0	13.4	10.1	8.6	14.3	37.9	17.6	14.2	11.1	20.2
5	C	24.9	12.3	11.0	8.8	14.2	43.3	17.6	17.3	14.1	23.1
6	N ₁ C	25.0	17.0	14.7	13.3	17.5	46.3	28.2	26.1	25.3	31.5
7	N ₂ C	28.5	26.1	21.7	21.3	24.4	54.4	50.7	44.8	43.9	48.4
8	N ₃ C	27.3	23.6	20.5	18.9	22.6	58.7	60.0	54.3	54.3	56.8
9	N ₁ ³ C	26.5	20.8	16.4	16.5	20.0	46.8	40.5	32.8	33.1	38.3
10	N ₂	24.1	25.1	20.6	19.0	22.2	36.4	41.2	34.3	31.0	35.7
11	N ₂ P	23.8	24.7	19.6	19.3	21.8	36.2	42.4	30.7	30.9	35.0
12	N ₂ PS	25.8	23.4	21.7	22.9	23.4	44.2	43.3	36.2	40.5	41.0
13	N ₂ PK	30.0	21.3	19.5	20.3	22.8	53.7	47.5	38.7	42.3	45.6
14	N ₂ PM	26.4	25.1	20.9	21.6	23.5	41.1	45.9	35.4	38.0	40.1
15	N ₂ *C	26.5	22.2	19.1	19.5	21.8	45.1	41.9	37.3	36.0	40.1
16	N ₂ 'C	33.2	25.6	28.7	23.5	27.8	56.1	52.2	50.8	43.5	50.6
17	N ₂ and C in (C) alternate (N ₂) years	23.3	11.4	9.7	8.5	13.2	36.4	17.3	14.4	12.5	20.2
18		27.8	22.1	22.2	22.2	23.6	45.3	38.1	39.9	39.8	40.8
19	R	28.9	23.6	19.1	20.2	23.0	43.6	37.8	31.5	35.3	37.0
20	N ₂ 'C'	23.2	19.9			21.6	36.2	37.2			36.7
101st season, 1944											
Plot	Section	V	II	I	III		V	II	I	III	
2A	D	29.8	22.7	16.6	15.8	21.2	62.4	48.5	41.8	36.4	47.3
2B	D	29.2	27.5	22.5	23.2	25.6	53.3	51.6	50.0	48.8	50.9
3	O	18.6	8.3	7.9	9.1	11.0	21.4	12.1	12.1	12.0	14.4
5	C	20.2	9.0	9.1	10.2	12.1	29.3	11.8	11.8	14.1	16.8
6	N ₁ C	25.1	11.7	14.9	13.1	16.2	38.7	18.7	22.9	19.2	24.9
7	N ₂ C	29.5	19.3	22.8	20.6	23.0	46.3	36.9	33.6	33.9	38.9
8	N ₃ C	29.9	23.6	24.9	27.0	26.4	54.5	44.4	53.2	45.4	49.4
9	N ₁ 'C	23.9	13.9	15.4	19.3	18.2	35.1	22.2	23.2	31.3	28.0
10	N ₂	16.2	18.4	17.4	17.6	17.4	26.2	27.4	29.4	25.2	27.0
11	N ₂ P	15.8	15.7	14.2	13.8	14.9	25.4	26.1	27.8	24.6	26.0
12	N ₂ PS	22.6	16.5	17.2	16.6	18.2	36.9	25.3	31.8	27.0	30.2
13	N ₂ PK	28.7	20.1	20.2	17.7	21.7	46.7	36.6	41.0	31.9	39.0
14	N ₂ PM	27.2	16.0	17.3	15.6	19.0	42.9	27.9	31.5	26.6	32.2
15	N ₂ *C	25.8	19.3	20.7	20.1	21.5	45.8	32.0	40.9	31.7	37.6
16	N ₂ 'C	31.4	23.4	23.7	26.5	26.2	51.6	30.8	36.8	37.1	39.1
17	N ₂ and C in (N ₂) alternate (C) years	26.9	18.9	19.2	20.0	21.2	44.3	28.2	31.1	29.2	33.2
18		27.2	8.9	6.1	7.7	12.5	42.1	9.2	8.5	10.6	17.6
19	R	29.3	18.1	16.1	15.4	19.7	50.9	28.4	23.4	27.5	32.6
20	N ₂ 'C'	20.7	13.9			177.33		33.4	21.3		27.4

X173 [†]Includes straw, cavings and chaff. * Sulphate of ammonia applied in autumn.

Wheat - Broadbalk

Years after fallow		Total Grain: cwt. per acre					Total Straw: cwt. per acre				
		1	2	3	4	Mean	1	2	3	4	Mean
102nd season, 1945											
Plot	Section	IV	V	II	I		IV	V	II	I	
2A	D	25.1	28.9	23.1	12.3	22.4	44.7	46.1	43.3	47.7	45.4
2B	D	23.4	31.0	25.0	17.5	24.2	52.5	49.0	46.5	52.5	50.1
3	O	14.6	11.6	6.5	7.6	10.1	22.1	14.6	10.8	12.0	14.9
5	C	20.9	14.9	9.9	10.5	14.0	34.0	17.6	13.3	15.5	20.1
6	N ₁ C	23.5	18.9	11.4	13.7	16.9	40.2	30.7	19.4	22.1	28.1
7	N ₁ ¹ C	23.3	20.8	16.1	22.3	20.6	42.0	36.2	29.5	43.4	37.8
8	N ₁ ² C	24.0	27.8	22.5	25.2	24.9	43.9	49.4	40.9	55.2	47.4
9	N ₁ ³ C	15.2	15.4	13.2	14.7	14.6	30.3	25.1	22.5	27.6	26.4
10	N ₂	14.6	21.7	15.3	13.7	16.3	24.4	31.8	24.1	24.8	26.3
11	N ₂ P	14.2	17.5	13.9	16.2	15.4	29.5	26.2	22.9	29.2	27.0
12	N ₂ PS	18.5	22.0	16.9	16.8	18.6	36.0	34.2	27.6	31.5	32.3
13	N ₂ PK	23.3	22.3	15.1	15.4	19.0	45.1	40.2	27.9	31.9	36.3
14	N ₂ PM	16.0	23.8	14.4	14.4	17.2	38.4	39.1	25.1	29.5	33.0
15	N ₂ *C	22.4	20.0	17.7	18.4	19.6	41.2	33.3	30.6	36.4	35.4
16	N ₂ 'C	21.0	23.6	20.8	20.4	21.4	41.8	41.5	39.3	43.4	41.5
17	N ₂ and C in (C) alternate (N ₂) years	16.6	10.3	7.7	9.2	11.0	22.9	9.1	15.9	12.4	11.5
18		20.8	16.5	17.2	16.4	17.7	38.3	26.5	28.4	28.4	32.9
19	R	23.2	18.0	14.9	16.3	18.1	41.1	27.3	25.2	28.3	30.4
20	N ₂ C'			13.6	15.2	14.4			28.1	28.6	28.4

Plot	Section	103rd season, 1946									
		III	IV	V	II		III	IV	V	II	
2A	D	31.7	15.4	20.0	17.9	21.2	71.6	55.1	55.7	45.1	56.9
2B	D	29.3	18.2	21.0	22.2	22.7	75.5	52.2	48.3	48.6	56.2
3	O	18.7	7.3	7.3	7.3	10.2	35.9	16.0	14.6	15.9	20.6
5	C	20.4	9.2	9.7	10.0	12.3	40.0	22.0	19.7	24.3	26.5
6	N ₁ C	24.4	9.8	9.9	8.8	13.2	53.6	29.2	25.8	26.8	33.8
7	N ₂ C	26.4	15.9	15.3	14.4	18.0	63.9	41.2	38.1	36.4	44.9
8	N ₂ C	31.4	20.2	20.1	17.0	22.2	72.0	59.9	57.7	52.6	60.6
9	N ₁ ³ C	26.6	18.2	14.6	12.1	17.9	52.8	40.3	34.2	31.9	39.8
10	N ₂	25.6	19.5	12.3	16.9	18.6	51.8	40.8	33.9	33.1	39.9
11	N ₂ P	20.2	15.2	15.2	14.5	16.3	39.3	44.2	34.1	32.1	37.4
12	N ₂ PS	22.4	15.3	11.9	16.0	16.4	45.3	37.7	34.0	33.2	37.6
13	N ₂ PK	29.0	17.4	14.9	14.7	19.0	62.0	44.0	39.6	38.0	45.9
14	N ₂ PM	25.3	20.9	16.8	15.1	19.5	47.8	45.5	41.4	32.6	41.8
15	N ₂ *C	25.3	17.9	15.5	14.7	18.4	58.1	38.5	32.8	34.5	41.0
16	N ₂ 'C	31.9	20.6	19.2	16.2	22.0	67.6	53.3	47.9	43.1	53.0
17	N ₂ and C in (C) alternate (N ₂) years	24.5	19.1	19.6	15.9	19.8	60.6	42.2	46.5	36.0	46.3
18		18.9	9.7	7.9	3.3	10.0	42.3	20.5	18.3	10.2	22.8
19	R	24.3	18.1	17.0	13.3	18.2	50.3	39.4	37.6	27.7	38.8
20	N ₂ C'					11.1			28.1		28.1

Years after fallow		Total Grain: cwt. per acre					Total Straw [†] : cwt. per acre				
		1	2	3	4	Mean	1	2	3	4	Mean
104th season, 1947											
Plot	Section	I	III	IV	V		I	III	IV	V	
2A	D	19.1	14.6	19.4	14.6	16.9	28.3	19.7	25.9	20.7	23.6
2B	D	20.6	16.9	15.0	14.5	16.8	31.9	25.7	20.3	20.6	24.6
3	O	10.1	4.4	5.7	4.9	6.3	12.0	5.3	7.3	6.7	7.8
5	C	16.5	6.2	9.5	7.9	10.0	24.3	9.1	12.8	12.9	14.8
6	NN ₁ C	19.0	9.0	8.8	7.3	11.0	25.0	13.2	15.6	13.8	16.9
7	N ₂ ¹ C	22.3	12.5	7.8	7.1	12.4	28.4	18.1	13.9	15.6	19.0
8	N ₂ ² C	20.7	16.7	12.4	11.5	15.3	34.3	23.6	22.3	23.0	25.8
9	N ₂ ¹ 'C	15.1	9.4	6.5	6.9	9.5	25.5	11.3	10.0	11.2	14.5
10	N ₂	8.1	8.1	5.3	4.1	6.4	16.7	11.5	7.7	6.6	10.6
11	N ₂ P	9.4	9.8	5.8	4.0	7.2	22.9	12.0	7.5	8.1	12.6
12	N ₂ PS	13.0	10.0	7.1	5.5	8.9	23.9	14.0	11.5	13.6	15.8
13	N ₂ PK	22.2	9.7	6.7	6.1	11.2	30.8	15.0	12.2	11.3	17.3
14	N ₂ PM	14.5	10.7	7.3	6.6	9.8	25.3	17.5	11.1	12.5	16.6
15	N ₂ *C	20.5	11.3	9.1	7.0	12.0	29.2	20.6	13.5	11.6	18.7
16	N ₂ 'C	19.2	13.6	11.3	11.1	13.8	28.5	19.3	19.4	18.8	21.5
17	N ₂ and C in (C)	14.6	5.0	5.6	6.4	7.9	21.7	9.7	8.4	9.7	12.4
18	alternate (N ₂) years	16.1	10.3	14.9	15.3	14.2	21.2	15.2	19.6	19.4	18.8
19	R	15.7	10.0	8.5	10.2	11.1	21.6	15.1	15.9	16.9	17.4
20	N ₂ C'	9.7				9.7	15.0				15.0

[†] Includes straw, cavings and chaff

* Sulphate of ammonia applied in autumn

⁺ Yields from small areas left after cutting green wheat and wild oats.
(only happened in 1940 and 1946)

BARLEY - HOOSFIELD

Experiments on barley grown continuously on the same land were begun on Hoosfield in 1852, the general purpose being the same as on Broadbalk. On Hoosfield however the mineral manures in various combinations were laid on strips running lengthways along the field, and these were crossed at right angles by strips of various nitrogenous manures, to give plots of approximately 1/6 acre - one for each treatment. Dung alone was applied to a plot outside this factorial arrangement.

Since 1919, Plumage Archer has been the variety grown, except that from 1929 to 1932 Plumage Archer and Spratt Archer were grown in strips running through all the plots. Weeds have been troublesome on Hoosfield, as on Broadbalk, and it was found necessary to fallow the whole field in 1912, 1933 and 1943. Commencing in 1944 a yearly spraying with DNOC (dinitro-ortho-cresol) was given in late May or early June to check broad-leaved weeds, but wild oat (*Avena fatua*) has become a serious pest in recent years and hand pulling is regularly carried out.

Weed surveys have been made by the Botany Department.

Treatments

Symbols and amounts per acre

Cross Dressings

- O Unmanured
- A Sulphate of ammonia, 43 lb. N
- AA Nitrate of soda, 43 lb. N
- AAS Nitrate of soda, 43 lb. N; silicate of soda, 400 lb.
- C Rape cake (castor bean meal from 1940 onwards), 1,000 lb.

Strip Dressings

- P Superphosphate, 65 lb. P₂O₅
- K Sulphate of potash, 98 lb. K₂O
- S Sulphate of soda, 100 lb.
- M Sulphate of magnesia, 100 lb.

Plot Dressings

- D Farmyard manure, 14 tons
- D' Unmanured, following farmyard manure 1852-71
- F Unmanured, following furnace ash 1852-1933
- N' Nitrate of soda, 43 lb. N
- N'' Nitrate of soda, 43 lb. N, following double dressing 1852-57

Dung is ploughed in in winter; all other manures are broadcast in spring during the preparation of the seedbed.

Crop Notes

Year	Sown	Harvested	Year	Sown	Harvested
1939	Mar. 10	Aug. 25	1944	Mar. 29	Aug. 18
1940	Mar. 12	Aug. 6	1945	Mar. 9	Aug. 8
1941	Mar. 17	Aug. 25	1946	Mar. 25	Aug. 23
1942	Mar. 29	Aug. 15	1947	Apr. 17	Aug. 19

Total grain: cwt. per acre

Plot	Cross Dressing	Strip Dressing	1939	1940	1941	1942	1944	1945	1946	1947
1	0	0	9.2	6.4	2.1	8.8	14.8	11.2	10.2	5.4
2	0	P	12.5	9.4	4.4	11.1	22.7	11.7	10.3	6.7
3	0	KSM	12.0	7.8	3.3	12.0	17.9	18.1	11.8	8.3
4	0	PKSM	14.7	10.9	4.7	14.9	27.8	21.1	13.3	9.1
5	0	PK	11.9	6.8	3.8	14.9	28.6	20.8	13.4	10.8
1	A	0	17.1	12.2	5.7	9.4	10.4	15.6	15.0	7.1
2	A	P	20.7	15.8	8.5	8.0	15.3	22.9	14.4	9.0
3	A	KSM	18.4	13.5	6.6	10.2	11.8	21.9	15.6	11.8
4	A	PKSM	24.9	14.4	12.2	12.2	21.1	22.0	16.0	12.5
5	A	PK	22.2	18.2	8.5	9.0	13.9	14.8	22.4	17.4
1	AA	0	16.5	13.9	5.5	6.0	12.1	18.1	19.6	7.7
2	AA	P	24.3	20.8	8.8	9.9	21.3	25.0	16.4	12.0
3	AA	KSM	18.0	14.3	6.8	10.8	11.4	24.2	18.8	11.5
4	AA	PKSM	24.6	21.2	12.2	14.9	24.4	28.7	22.5	13.4
1	AAS	0	19.8	19.5	7.1	9.6	14.4	18.4	23.3	11.0
2	AAS	P	24.5	20.5	8.2	10.2	24.4	24.0	20.0	17.4
3	AAS	KSM	22.4	19.4	8.4	12.8	12.4	22.6	12.5	13.6
4	AAS	PKSM	26.2	23.2	12.9	16.2	26.5	26.8	24.8	16.6
1	C	0	21.0	16.6	7.7	7.4	16.4	19.4	14.7	14.6
2	C	P	21.4	16.6	7.6	7.5	28.5	27.4	17.9	18.6
3	C	KSM	19.6	16.8	9.5	10.3	22.1	24.9	40.5	16.7
4	C	PKSM	24.1	19.3	10.6	13.2	28.5	27.5	38.9	16.1
7-1	D'		15.3	9.7	4.2	13.8	25.4	23.6	14.0	10.5
7-2	D		32.8	20.8	7.6	12.7	30.9	36.6	32.6	21.0
6-1	O		10.9	6.3	2.1	8.7	17.8	14.6	8.1	6.9
6-2	F		10.4	5.3	1.9	6.9	15.1	13.2	11.1	8.1
1	N'		17.4	14.8	7.1	8.1	13.1	18.5	17.8	12.9
2	N"		21.4	17.8	7.8	8.0	17.8	23.1	17.8	19.6

X
Vertical LinesBarley - Hoosfield

A/2 3

Barley - Hoosfield

Total straw: cwt. per acre

Plot	Cross Dressing	Strip Dressing	1939	1940	1941	1942	1944	1945	1946	1947
1	0	0	11.8	8.6	8.2	10.9	14.8	12.1	9.8	5.0
2	0	P	12.3	7.3	7.1	10.8	18.2	10.8	9.4	6.4
3	0	KSM	19.0	9.6	9.9	15.2	17.4	17.0	13.0	8.3
4	0	PKSM	20.0	13.7	12.1	19.5	23.2	18.1	14.8	9.1
5	0	PK	15.5	9.9	14.7	19.0	27.8	19.9	14.4	10.0
1	A	0	21.1	18.8	18.5	16.1	11.6	15.8	15.4	5.7
2	A	P	23.0	20.7	21.5	16.6	14.9	25.9	16.5	9.5
3	A	KSM	23.0	16.1	18.7	17.8	12.7	24.7	21.5	13.4
4	A	PKSM	25.0	20.9	19.3	23.1	21.6	31.0	22.2	10.6
5	A	PK	29.1	28.7	25.2	29.8	24.0	39.2	31.4	20.0
1	AA	0	27.2	27.6	18.9	19.9	15.3	25.7	22.4	9.6
2	AA	P	27.9	24.7	25.2	22.1	21.2	30.0	22.4	14.5
3	AA	KSM	25.1	21.2	23.8	20.7	13.1	28.4	25.3	15.4
4	AA	PKSM	25.7	24.5	21.8	27.6	22.4	32.0	28.0	11.1
1	AAS	0	28.3	26.0	22.2	26.3	19.8	30.4	27.1	12.6
2	AAS	P	27.8	29.0	21.5	25.2	24.6	32.1	26.7	15.0
3	AAS	KSM	27.2	22.4	22.3	27.0	15.9	32.4	20.1	13.0
4	AAS	PKSM	27.5	25.4	21.7	28.2	27.3	35.2	31.9	12.7
1	C	0	25.7	25.1	21.5	21.6	18.1	29.4	24.1	15.2
2	C	P	25.4	23.9	21.0	21.4	26.9	32.0	23.6	17.8
3	C	KSM	23.0	21.1	21.5	24.3	24.0	29.3	61.5	15.4
4	C	PKSM	25.8	23.1	20.1	27.2	32.6	33.4	52.3	26.0
7-1	D'		21.0	16.0	13.2	16.6	23.9	23.5	17.9	11.8
7-2	D		41.4	35.6	32.2	31.7	32.5	45.2	47.3	21.4
6-1	O		14.1	10.1	7.9	13.8	18.7	14.5	10.7	8.4
6-2	F		13.9	6.4	7.8	11.4	13.8	13.5	11.3	8.3
1	N'		24.0	23.4	22.8	23.3	16.6	26.8	21.7	13.9
2	N"		28.5	25.4	22.7	24.8	19.8	29.3	21.8	18.2

35.6

X

WHEAT AFTER FALLOW - HOOSFIELD

Comparison of a Three-Year Fallow with a One-Year Fallow

The land has been unmanured since 1851. There are two $\frac{1}{2}$ -acre strips, each divided into four parts, and the strips are fallowed in alternate years. In 1932 and after, one quarter of the strip in crop has also been fallowed, different quarters being selected in successive years, thus providing in every year from 1934 onwards a comparison of the effect of a three-year fallow with the effect of a one-year fallow. Half the experiment is under wheat after a one-year fallow, so that continuity with previous results is maintained.

Cropping of strips A and B

C = Crop. F = Fallow.

Year	A1	A2	A3	A4	B1	B2	B3	B4
1932, 40	F	C	C	C	F	F	F	F
1933, 41	F	F	F	F	C	C	F	C
1934, 42	C	F	C	C	F	F	F	F
1935, 43	F	F	F	F	C	C	C	F
1936, 44	C	C	F	C	F	F	F	F
1937, 45	F	F	F	F	F	C	C	C
1938, 46	C	C	C	F	F	F	F	F
1939, 47	F	F	F	F	C	F	C	C

Crop Notes

Crop Year	Date Sown	Date Harvested	Variety
1939	Oct. 24	Aug. 24	Red Standard
1940	Oct. 23	Aug. 8	Squareheads Master
1941	Oct. 28	Aug. 21	Squareheads Master
1942	Oct. 21	Aug. 17	Sherriffs "Stand Up"
1943	Oct. 15	Aug. 3	Squareheads Master
1944	Oct. 8	Aug. 10	Red Standard
1945	Oct. 30	Aug. 11	Red Standard
1946	Oct. 15	Aug. 16	Squareheads Master 13/4
1947	Oct. 18	Aug. 12	Squareheads Master 13/4

N

Yields: cwt per acre

Year	Total Grain			Mean	Total Straw*			Mean
	After 1 year fallow	After 3 years fallow	Mean		After 1 year fallow	After 3 years fallow	Mean	
1939	B3 12.5	B4 9.7	B1 11.2	11.1	B3 20.1	B4 16.2	B1 19.6	18.6
1940	A2 11.4	A3 12.2	A4 12.3	12.0	A2 17.9	A3 16.3	A4 18.2	17.5
1941	B4 6.9	B1 7.9	B2 9.3	8.0	B4 10.9	B1 12.0	B2 14.8	12.6
1942	A3 12.4	A4 10.6	A1 14.1	12.4	A3 16.3	A4 16.0	A1 20.0	17.4
1943	B1 14.9	B2 16.5	B3 19.7	17.0	B1 21.6	B2 23.9	B3 29.3	24.9
1944	A4 13.6	A1 17.4	A2 15.0	15.3	A4 18.2	A1 23.9	A2 20.4	20.8
1945	B2 11.2	B3 9.5	B4 9.9	10.2	B2 18.0	B3 16.7	B4 16.6	17.1
1946	A1 11.1	A2 9.9	A3 11.2	10.7	A1 23.1	A2 21.4	A3 26.0	23.5
1947	B3 6.2	B4 5.3	B1 8.9	6.8	B3 8.5	B4 8.1	B1 11.6	9.4

*Includes straw, cavings and chaff.

CROPS GROWN IN ROTATION - AGDELL FIELD

The experiment, which began in 1848, compares two crop rotations, one with clover and the other with fallow, and the effect of mineral manure with and without nitrogen applied to the root crops.

Area of each plot, 2/5 acre.

Rotations: Turnips or swedes, barley, clover or beans, wheat.
Turnips or swedes, barley, fallow, wheat.

Until 1904, half of each plot of turnips was fed on the land, the remainder being carted off. Subsequently all the roots were carted.

Treatments and rates of dressing.

Applied to the root crops only (all amounts per acre):

None;

Mineral manure, with no nitrogen: Superphosphate, 84 lb. P₂O₅;
Sulphate of potash, 245 lb. K₂O; sulphate of soda, 100 lb;
sulphate of magnesia, 200 lb.

Complete mineral and nitrogenous manure: As above, together with sulphate of ammonia, 43 lb. N and 2,000 lb. castor bean meal or rape dust.
per acre.

Crop Notes.

Year	Crop	Sown	Harvested	Variety
1939	Wheat	Oct. 27, 1938	Aug. 15	Red Standard
1940	Turnips	May 31	Nov. 28	Bruce
1941	Barley	March 19	Sept. 10	Plumage Archer
1942	Clover	May 9, 1941	June 29	Montgomery Red
1943	Wheat	Oct. 15, 1942	Aug. 2	Red Standard
1944	Swedes	June 23	Nov. 5	Mixed seed
1945	Barley	March 9	Aug. 14	Plumage Archer
1946	Clover	April 6, 1945	June 24	Montgomery Red
1947	Winter wheat	Nov. 6, 1946	Aug. 12	Squareheads Master 13/4
1947	Spring wheat	April 18	Aug. 21	Atle

In 1940 the turnips, especially those on plots 1 and 2, were badly attacked by Finger and Toe disease.

In 1944 three varieties of swedes and eleven varieties of turnips were sown in strips down each plot. The turnips on all plots failed, as did also the swedes on plots 5 and 6; all plots were attacked by disease, plots 3 and 4 less severely than the rest.

In 1947 the wheat wintered badly, and it was decided to resow all of plots 1, 3 and 5 and the west side of plots 2, 4 and 6 with spring wheat.

Produce per acre

Year	Crop	O		M		C		
		Unmanured since 1848		Mineral Manure		Complete mineral and nitrogenous manure		
		5	6	3	4	1	2	
Fallow	beans	Fallow	beans	Fallow	beans	Fallow	beans	
Twenty-third Course								
1936	Roots (Turnips)	tons	1.22	0.47	2.69	2.55	5.63	3.26
1937	Barley	cwt.	0.5	0.3	3.1	0.5	0.6	1.0
	Grain	cwt.	3.4	2.1	2.5	4.7	2.7	3.4
	Straw*	cwt.	-	8.3	-	29.1	-	26.4
1938	Clover hay	cwt.	14.9	11.9	18.2	22.2	12.8	20.9
1939	Wheat	cwt.	26.3	19.9	32.9	36.6	24.9	33.2
Twenty-fourth Course								
1940	Root (Turnips)	tons	0.52	0.14	3.96	5.09	2.76	2.71
1941	Barley	cwt.	7.4	3.2	9.6	11.5	10.3	4.2
	Grain	cwt.	11.9	10.8	16.6	20.2	12.9	18.6
	Straw*	cwt.	-	10.0	-	39.9	-	32.9
1942	Clover hay	cwt.	19.7	17.7	24.7	27.4	21.2	27.1
1943	Wheat	cwt.	30.3	29.2	39.4	41.7	34.6	42.4
Twenty-fifth Course								
1944	Roots (Swedes)	tons	Failed		5.26	5.01	3.53	2.09
1945	Barley	cwt.	12.4	9.5	19.8	20.9	24.9	20.6
	Grain	cwt.	11.4	10.1	21.5	16.7	25.1	24.9
	Straw*	cwt.	-	7.4	-	44.9	-	25.0
1946	Clover hay	cwt.	-	7.0	-	16.0	-	12.3
1947	Winter wheat	cwt.	-	13.3	-	24.9	-	17.3
1947	Spring wheat	cwt.	12.8	10.4	20.3	18.6	18.5	16.8
	Grain	cwt.	15.8	15.1	26.2	25.0	22.3	19.3
	Straw*	cwt.						

*Includes straw, cavings and chaff.

MANGOLDS AND SUGAR BEET - BARNFIELD

Experiments on root crops have been conducted on Barnfield since 1843. White turnips were grown from 1843-1848, followed by swedes until 1870 except for three years 1853-1855 when the crop was barley. Sugar beet was grown from 1871-1875 and then in 1876 the field was established under mangolds, variety Yellow Globe, with practically the present scheme of manuring. Since 1946 four rows of sugar beet, variety Klein E, have been drilled on each plot in order to compare the growth of mangolds and sugar beet under a wide range of fertilizer treatments. The leaves of the crops are spread and ploughed in on their respective plots.

Area of each plot, 1/7 to 1/5 acre.

Symbols and amounts per acre.

Cross Dressings

O	Unmanured
N	Nitrate of soda, 86 lb. N
A	Sulphate of ammonia, 86 lb. N
C	Rape cake, (castor bean meal from 1940 onwards) 2000 lb.

Strip Dressings

P	Superphosphate, 65 lb. P ₂ O ₅
K	Sulphate of potash, 245 lb. K ₂ O
S	Sodium chloride, 200 lb.
M	Sulphate of magnesia, 200 lb.
Ca	Calcium chloride, 190 lb; potassium nitrate, 570 lb; calcium nitrate, 100 lb.
D	Farmyard manure, 14 tons

Treatments

Strip	Strip	Cross Dressings
1 D	6 PK	O
2 DPK	7 PSM	N
4 PKSM	8 O	A
5 P	9 NKSM	AC
		C

From 1904 onwards, plot 4N has been divided, 4(a) receiving the dressing NPKSM, 4(b) receiving the dressing NPCa.

Dung is ploughed down in winter. The mineral manures, castor bean meal, and one third of the sulphate of ammonia and nitrate of soda are applied after the first cultivation but before the seed is drilled. The remaining two-thirds of the nitrogenous fertilizers are applied as a top-dressing about the time of singling.

Crop Notes

Year	Date sown	Date lifted	Year	Date sown	Date lifted
1939	May 9	Nov. 11	1944	May 3	Nov. 16
1940	May 10	Nov. 21	1945	April 18	Oct. 26
1941	April 18	Nov. 25	1946 ⁽¹⁾	May 3	Nov. 20
1942	May 1	Dec. 7	1946 ⁽²⁾	May 3	Nov. 11
1943	April 27	Nov. 5	1947 ⁽¹⁾	May 16	Oct. 31
			1947 ⁽²⁾	May 16	Oct. 30

(1) Mangolds

(2) Sugar beet

Investigations by the Soil Microbiology Department are described by A.V.Garcia, "Contribuicoes para o connecimento da Microbiologia do solo. Estudos sobre as bacterias autoctones e amibas de tres talhoes do campo experimental de Barnfield", Anais do Instituto Superior de Agronomia 18. (1951) 1, and by B.N.Singh, "The effect of artificial fertilizers and dung on the numbers of amoebae in Rothamsted soils", J.Gen.Microbiol., 3, (1949) 204.

Mangolds and Sugar Beet - Barnfield

Mangolds

Strip Dressing	Roots: tons per acre					Leaves: tons per acre				
	Cross Dressing					Cross Dressing				
	O	N	A	AC	C	O	N	A	AC	C
1939										
1 D	10.86	18.69	16.87	23.55	22.18	3.17	4.63	4.30	5.39	5.23
2 DPK	14.07	22.14	19.06	28.67	24.00	3.46	5.03	4.65	6.09	5.00
4 PKSM	3.62	18.05	15.42	25.60	18.99	1.20	4.09	3.54	5.71	3.72
4(b)PCa		17.76					4.82			
5 P	3.69	19.18	10.16	12.00	12.45	1.17	4.50	3.50	4.06	3.74
6 PK	3.07	17.41	15.91	23.09	17.76	0.95	4.34	3.25	5.62	3.72
7 PSM	3.51	19.35	18.19	23.90	20.76	1.10	4.63	3.70	4.92	4.29
8 O	2.29	9.71	6.48	10.03	10.31	0.92	3.57	2.90	3.91	3.98
9 NKSM	17.10	-	-	-	-	3.68	-	-	-	-
1940										
1 D	15.03	25.82	20.64	23.65	23.77	2.44	4.61	4.30	4.62	3.78
2 DPK	18.11	30.35	27.17	30.61	26.37	2.82	6.03	5.40	5.75	4.17
4 PKSM	4.43	21.81	18.51	28.19	22.65	1.13	4.38	3.47	5.21	3.86
4(b)PCa		21.85					4.56			
5 P	4.64	19.91	12.69	15.60	14.64	0.91	2.92	2.91	4.17	3.29
6 PK	4.33	20.67	16.31	26.09	20.24	0.86	2.84	2.41	4.43	3.17
7 PSM	3.87	22.42	18.52	21.48	21.79	0.99	4.29	3.93	5.71	4.75
8 O	5.53	17.73	10.30	12.11	13.22	1.02	3.90	3.20	4.01	3.49
9 NKSM	19.54	-	-	-	-	3.02	-	-	-	-
1941										
1 D	3.63	13.97	12.10	18.17	15.68	0.87	4.27	3.54	4.37	3.80
2 DPK	4.49	17.28	19.35	22.67	20.60	1.29	4.45	4.41	5.02	4.04
4 PKSM	0.12	8.54	9.40	19.39	17.12	0.08	2.90	2.91	4.15	3.02
4(b)PCa		6.03					2.18			
5 P	0.24	8.30	8.48	11.22	12.39	0.18	2.06	2.76	3.57	3.12
6 PK	0.43	8.59	12.42	15.29	15.16	0.25	1.91	2.50	3.72	2.87
7 PSM	0.16	6.47	10.20	18.57	15.89	0.12	1.65	2.55	4.06	3.26
8 O	0.70	3.50	2.61	3.06	2.97	0.34	1.54	1.46	0.91	1.16
9 NKSM	11.81	-	-	-	-	2.56	-	-	-	-
1942										
1 D	10.38	25.31	20.02	24.00	21.09	1.83	3.13	3.30	3.47	2.35
2 DPK	15.55	31.06	29.16	32.34	27.36	2.10	4.45	4.09	4.67	3.35
4 PKSM	1.64	24.04	22.56	31.08	24.95	0.32	3.60	2.84	4.28	2.30
4(b)PCa		22.93					4.31			
5 P	1.97	17.43	8.40	6.40	9.92	0.66	2.45	2.94	1.66	2.69
6 PK	1.14	20.81	18.62	23.46	19.62	0.34	2.98	1.88	3.74	2.01
7 PSM	0.75	22.25	17.77	18.74	19.94	0.10	3.77	2.54	3.03	3.33
8 O	2.40	15.78	8.61	6.10	8.17	0.78	4.13	2.69	2.45	2.779
9 NKSM	20.98	-	-	-	-	2.84	-	-	-	-

Mangolds

Strip Dressing	Roots: tons per acre					Leaves: tons per acre				
	Cross Dressing					Cross Dressing				
	O	N	A	AC	C	O	N	A	AC	C
1943										
1 D	12.58	25.71	19.14	30.08	28.10	3.71	6.43	4.89	7.36	4.67
2 DPK	14.14	22.34	21.55	32.26	25.69	3.91	6.39	6.65	6.90	5.70
4 PKSM	2.28	11.24	12.79	29.92	18.27	1.08	7.12	4.53	6.97	5.82
4(b)PCa		13.36					6.36			
5 P	3.13	12.89	11.69	12.44	14.07	1.32	5.33	3.47	5.70	5.21
6 PK	2.18	17.64	15.84	27.85	19.47	0.66	4.21	3.62	6.17	3.47
7 PSM	1.86	17.85	13.72	22.91	23.06	0.68	6.09	4.67	6.17	5.43
8 O	1.82	17.52	10.85	11.47	12.21	1.05	5.26	4.21	7.88	5.24
9 NKSM	19.97	-	-	-	-	4.57	-	-	-	-
1944										
1 D	8.59	14.62	13.79	14.09	10.34	3.77	5.24	4.89	5.63	5.82
2 DPK	9.85	12.40	13.85	20.86	18.34	4.01	5.43	4.40	7.58	6.56
4 PKSM	1.48	8.06	6.76	16.30	9.53	0.76	4.31	3.91	6.51	4.40
4(b)PCa		9.16					3.96			
5 P	0.91	3.61	1.96	4.98	5.45	1.08	2.89	2.57	3.28	3.87
6 PK	1.27	7.16	8.31	19.66	14.17	1.13	4.26	4.99	5.99	3.62
7 PSM	1.31	4.70	4.88	16.27	14.18	0.95	3.16	3.28	6.19	4.75
8 O	0.44	5.69	2.67	5.71	5.21	0.61	3.87	2.59	4.35	4.06
9 NKSM	12.25	-	-	-	-	4.33	-	-	-	-
1945										
1 D	12.18	22.31	17.82	30.10	25.36	3.65	3.74	5.48	4.89	4.70
2 DPK	14.06	21.36	17.59	34.63	26.68	3.69	5.95	4.26	6.02	4.89
4 PKSM	3.61	16.99	18.09	32.45	20.52	1.52	5.82	6.21	7.12	4.09
4(b)PCa		15.57					5.33			
5 P	3.02	9.19	11.51	15.11	15.66	0.95	2.84	2.91	4.48	2.96
6 PK	3.20	14.41	14.47	33.32	24.51	1.20	5.31	3.89	5.21	3.06
7 PSM	2.98	10.43	13.53	21.36	20.83	0.81	3.96	5.02	4.53	4.45
8 O	3.19	12.16	8.06	13.41	15.51	1.08	5.19	4.01	2.76	3.43
9 NKSM	22.91	-	--	-	-	4.35	-	-	-	-
1946										
1 D	8.16	19.30	14.73	12.09	16.12	1.49	4.77	4.87	4.31	4.84
2 DPK	9.13	20.00	17.97	13.31	18.92	2.72	6.14	5.19	5.82	4.28
4 PKSM	2.14	12.76	8.72	16.20	13.50	7.29	4.87	2.98	4.01	3.77
4(b)PCa		15.35					4.72			
5 P	1.14	10.47	3.20	13.70	11.51	3.82	3.77	1.76	2.91	2.42
6 PK	1.50	9.02	8.97	28.85	13.54	3.57	2.25	3.28	4.48	3.57
7 PSM	1.46	12.02	8.58	11.19	9.75	3.28	3.65	2.52	4.50	2.40
8 O	1.46	4.03	2.71	12.69	13.33	3.07	1.49	2.25	2.25	3.33
9 NKSM	8.48	-	-	-	-	3.25	-	-	-	-

Strip Dressings	Roots: tons per acre					Leaves: tons per acre				
	O	N	A	AC	C	O	N	A	AC	C
1947	Mangolds									
1 D	6.41	11.29	10.91	10.01	10.01	1.79	2.10	1.91	2.18	2.79
2 DPK	9.54	14.39	12.29	12.87	13.15	1.44	3.11	2.62	3.45	2.40
4 PKSM	3.10	10.57	7.29	7.86	9.84	1.44	2.62	2.25	2.45	2.45
4(b) PCa		11.52					2.18			
5 P	1.93	8.60	1.63	2.64	3.69	0.69	1.83	0.64	0.83	1.57
6 PK	2.27	7.76	6.07	7.36	6.88	0.61	2.45	1.61	2.23	2.35
7 PSM	1.89	8.22	7.11	5.25	4.49	0.59	2.37	1.93	1.83	1.57
8 O	0.57	0.71	0.42	3.51	3.67	0.44	0.59	0.49	0.98	1.13
9 NKSM	3.67	-	-	-	-	1.79	-	-	-	-
1946	Sugar beet									
	Roots (washed): tons per acre					Tops: tons per acre				
2 DPK	6.01	7.24	6.41	13.28	9.20	5.82	12.48	8.71	15.71	12.09
4 PKSM	1.27		5.06	8.54	6.80	2.69		4.26	10.42	5.97
4(b) PCa		8.12					9.39			
5 P	1.24	7.19	1.59	11.11	6.95	3.91	10.86	2.50	10.62	5.24
6 PK	1.17	6.39	4.89	9.59	10.13	2.40	9.54	3.77	15.02	7.93
7 PSM	1.43	7.66	5.50	6.12	8.32	3.62	6.85	5.63	12.09	12.87
8 O	0.38	2.79	2.10	8.12	7.24	1.50	4.11	2.50	11.35	11.06
1947										
1 D	6.28	7.65	7.60	6.34	8.02	4.65	6.41	6.46	5.48	7.19
2 DPK	4.92	7.05	5.65	6.26	8.24	5.97	6.56	5.72	5.58	6.70
4 PKSM	1.90		5.59	6.53	6.23	1.22		5.33	7.29	4.70
4(b) PCa		7.49					5.48			
5 P	2.28	5.65	1.90	3.01	4.06	1.03	6.80	2.50	3.13	3.91
6 PK	1.35	5.66	4.04	6.84	6.20	1.32	6.46	2.54	7.00	4.21
7 PSM	1.75	5.87	4.43	3.88	4.64	1.47	7.05	3.82	5.28	4.26
8 O	1.55	1.86	0.49	1.38	1.28	1.37	2.50	1.17	3.23	3.57

No Sugar beet was grown on strip 1 in 1946.

HAY - THE PARK GRASS PLOTS

Long period effects of fertilizers and manures on the yield and botanical composition of meadow hay

This field has been under grass for centuries. The experimental treatments were first applied in 1856. Until 1874 the aftermath was grazed by sheep, but since then a second cut of hay has been taken in late summer, and no stock has been admitted.

In 1903 most of the plots were divided, the southern halves receiving 18 cwt. of ground lime per acre, every fourth year.

Most plots are $\frac{1}{4}$ acre, but the sizes range from $\frac{1}{2}$ to $\frac{1}{12}$ acre.

The experiment is discussed, and early departures from the present manuring system described, by R.O.Cashen, J.Agric.Sci., 37, (1947), 1.

Plot		Treatments (all amounts per acre)
1	N ₁	Sulphate of ammonia (43 lb.N); (with dung (14 tons) also, 1856-63).
2	O	Unmanured, following dung (14 tons), 1856-63.
3	O	Unmanured.
4-1	P	Superphosphate (65 lb. P ₂ O ₅).
4-2	N ₂ P	Sulphate of ammonia (86 lb.N), superphosphate (65 lb.P ₂ O ₅).
5-1	O	Unmanured, following ammonium salts (86 lb.N), 1856-97.
5-2	PK	Superphosphate (65 lb. P ₂ O ₅), sulphate of potash (245 lb. K ₂ O), following ammonium salts (86 lb.N) 1856-97.
6	PKM	Complete minerals as Plot 7, following ammonium salts (86 lb. N), 1856-68.
7	PKM	Complete minerals: Superphosphate (65 lb. P ₂ O ₅). Sulphate of potash (245 lb. K ₂ O). Sulphate of soda (100 lb.) Sulphate of magnesia (100 lb.)
8	PM	Complete minerals without sulphate of potash.
9	N ₂ PKM	Complete minerals, with sulphate of ammonia (86 lb. N).
10	N ₂ PM	Complete minerals, without sulphate of potash; with sulphate of ammonia (86 lb. N).

Plot

- 11-1 N₃PKM Complete minerals, with sulphate of ammonia (129 lb. N).
- 11-2 N₃PKMS As plot 11-1, with silicate of soda (400 lb.) since 1862.
- 12 C Unmanured.
- 13 DF Dung (14 tons) in 1905 and every fourth year since (omitted in 1917), fish guano (6 cwt.) in 1907 and every fourth year since; following complete minerals with ammonium salts (86 lb. N), 1856-1904.
- 14 N₂'PKM Complete minerals, with nitrate of soda (86 lb. N).
- 15 PKM Complete minerals, following nitrate of soda (86 lb. N), 1858-63.
- 16 N₁'PKM Complete minerals, with nitrate of soda (43 lb. N).
- 17 N₁' Nitrate of soda (43 lb. N).
- 18 N₂KM Complete minerals without superphosphate, with sulphate of ammonia (86 lb. N); following minerals and ammonium salts supplying the constituents of 1 ton of hay, 1865-1904.
- 19 D Dung (14 tons) in 1905 and every fourth year since except 1917, following nitrate of soda (43 lb. N), superphosphate (65 lb. P₂O₅) and sulphate of potash (142 lb. K₂O), 1872-1904.
- 20 D;C Dung (14 tons) in 1905 and every fourth year since except in 1917; in each intervening year nitrate of soda (27 lb. N), superphosphate (33 lb. P₂O₅) and sulphate of potash (49 lb. K₂O); following superphosphate (66 lb. P₂O₅) and nitrate of potash (327 lb.), 1872-1904.

Ground lime was applied to the southern portion (limed) of the plots at the rate of 2,000 lb. per acre in the winters of 1903-4, 1907-8, 1915-16, 1923-24, 1927-28, 1931-32, 1935-36, 1939-40, 1943-44 and at the rate of 2,500 lb. per acre in the winter of 1920-21, except on plots 18, 19 and 20, where part received light liming and part heavy liming as follows (weights in lb. per acre):-

Plot	Light liming (LL)	Heavy Liming (HL)
18	3,951	6,788
19	570	3,150
20	570	2,772

Hay - Park Grass

Dung and lime are applied in winter, minerals as early in spring as possible, and nitrogenous fertilizers in March. On plots 11-1, 11-2 and 16 the nitrogenous fertilizer is given in two dressings one month apart.

Dates of cutting

Year	1st cut	2nd cut	Year	1st cut	2nd cut
1939	July 1	Sept. 28	1944	June 21	Nov. 22
1940	June 20	Jan. 1941	1945	June 19	Nov. 9
1941	June 25	Oct. 1	1946	June 22	Dec. 24
1942	June 10	Sept. 19	1947	June 13	Sept. 2
1943	June 21	Nov. 25			

The second crop is carted green, and hay yields are estimated from the dry matter.

Yield of hay: cwt. per acre

Plot	1939									1940								
	Not limed			Limed			Not limed			Limed			Crtcp			Crop		
	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total	1st	2nd	Crop	Total		
1	8.4	16.8	25.2	20.4	11.8	32.2	10.6		10.6	16.0		16.0						
2	13.0	11.4	24.4	13.8	9.0	22.8	13.0		13.0	14.5		14.5						
3	11.1	9.9	21.0	12.8	6.5	19.3	13.6		13.6	14.5		14.5						
4-1	19.0	14.4	33.4	17.7	10.6	28.3	14.7		14.7	16.0		16.0						
4-2	12.2	20.2	32.4	29.6	12.0	41.6	14.1		14.1	27.9		27.9						
5-1	9.5	9.1	18.6	-	-	-	9.6		9.6	-		-						
5-2	20.9	12.9	33.8	-	-	-	21.9		21.9	-		-						
6	32.9	20.9	53.8	-	-	-	22.4		22.4	-		-						
7	35.5	23.6	59.1	41.1	26.2	67.3	24.2		24.2	33.3		33.3						
8	20.7	12.0	32.7	14.3	10.9	25.2	16.5		16.5	19.4		19.4						
9	32.8	26.6	59.4	54.0	22.5	76.5	36.7	3.8	40.5	39.3	1.2	40.5						
10	23.8	17.2	41.0	43.6	19.0	62.6	16.7		16.7	31.1	1.2	32.3						
11-1	25.6	29.2	54.8	54.8	31.8	86.6	31.3	18.2	49.5	48.9	6.6	55.5						
11-2	36.1	28.0	64.1	56.8	35.2	92.0	40.8	17.8	58.6	51.6	9.4	61.0						
12	15.0	9.6	24.6	-	-	-	11.5		11.5									
13	37.5	23.4	60.9	32.0	20.5	52.5	27.7	2.0	29.7	24.1	0.9	25.0						
14	52.4	23.1	75.5	52.3	19.1	71.4	51.5	2.6	54.1	48.6	1.1	49.7						
15	20.0	15.1	35.1	31.1	17.8	48.9	18.0		18.0	28.5		28.5						
16	39.5	19.4	58.9	37.4	19.4	56.8	35.7	1.6	37.3	39.3	0.6	39.9						
17	17.8	12.9	30.7	21.8	9.0	30.8	19.8		19.8	25.9		25.9						
18	12.1	17.1	29.2	29.5*	9.6*	39.1*	6.8	2.9	9.7	24.3*		24.3*						
				25.7†	8.2†	33.9†				20.2†		20.2†						
19	25.0	18.1	43.1	26.7*	17.4*	44.1*	27.2		27.2	23.8*		23.8*						
				25.8†	16.0†	41.8†				23.4†		23.4†						
20	37.6	19.2	56.8	36.1*	22.4*	58.5*	33.5		33.5	38.4*		38.4*						
				33.2†	18.1†	51.3†				35.0†		35.0†						

* Heavy liming.

† Light liming.

Hay - Park Grass

Yield of hay: cwt. per acre

Plot	1941						1942					
	Not limed			Limed			Not limed			Limed		
	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total
1	6.7	6.3	13.0	18.5	5.5	24.0	5.8	4.0	9.8	15.0	4.1	19.1
2	11.4	6.6	18.0	11.7	5.8	17.5	4.0	1.8	5.8	7.3	1.2	8.5
3	9.8	7.1	16.9	12.5	11.9	24.4	3.3	0.6	3.9	5.8	0.6	6.4
4-1	17.3	6.4	23.7	14.7	7.3	22.0	8.5	0.9	9.4	8.7	1.4	10.1
4-2	10.8	4.6	15.4	23.7	4.0	27.7	15.0	4.8	19.8	27.2	3.6	30.8
5-1	6.5	6.2	12.7	-	-	-	7.5	1.4	8.9	-	-	-
5-2	12.8	8.8	21.6	-	-	-	13.6	4.6	18.2	-	-	-
6	21.4	13.8	35.2	-	-	-	8.1	4.8	12.9	-	-	-
7	27.8	20.3	48.1	31.4	14.6	46.0	25.5	4.8	30.3	35.4	3.0	38.4
8	19.4	15.8	35.2	13.6	13.4	27.0	12.2	2.6	14.8	9.7	1.9	11.6
9	15.8	6.7	22.5	39.4	16.6	56.0	46.6	13.8	60.4	40.2	3.8	44.0
10	13.1	14.1	27.2	28.7	10.6	39.3	11.8	7.4	19.2	29.4	4.6	34.0
11-1	9.4	24.0	33.4	45.0	24.3	69.3	35.6	25.1	60.7	45.8	6.4	52.2
11-2	31.2	29.2	60.4	50.5	34.0	84.5	46.6	23.8	70.4	54.2	8.4	62.6
12	11.4	8.5	19.9	-	-	-	7.2	2.8	10.0	-	-	-
13	29.2	32.6	61.8	25.2	20.6	45.8	32.6	3.1	35.7	21.8	2.1	23.9
14	52.6	16.3	68.9	50.8	9.2	60.0	52.4	6.5	58.9	46.9	6.0	52.9
15	19.3	9.6	28.9	29.9	7.7	37.6	14.7	2.8	17.5	23.6	1.2	24.8
16	30.8	10.7	41.5	33.9	8.3	42.2	33.3	3.6	36.9	35.2	3.4	38.6
17	19.3	7.0	26.3	21.4*	3.8	25.2	12.5	5.2	17.7	22.2	2.9	25.1
18	7.2	12.8	20.0	22.1*	7.2*	29.3*	7.8	4.5	12.3	21.3*	1.6*	22.9*
				21.1†	7.0†	28.1†				24.3†	2.2†	26.5†
19	30.9	15.5	46.4	28.9*	14.7*	43.6*	22.5	5.1	27.6	23.5*	4.9*	28.4*
				31.3†	17.9†	49.2†				25.5†	2.9†	28.4†
20	30.4	14.6	45.0	31.9*	13.8*	45.7*	34.0	4.2	38.2	37.7*	2.9*	40.6*
				28.9†	11.0†	39.9†				40.7†	4.9†	45.6†

* Heavy liming.

† Light liming.

Yield of hay: cwt. per acre

Plot	1943						1944					
	Not limed			Limed			Not limed			Limed		
	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total
1	8.4	-	8.4	13.8	-	13.8	0.7	-	0.7	5.4	-	5.4
2	11.4	-	11.4	9.9	-	9.9	3.8	-	3.8	2.0	-	2.0
3	10.9	-	10.9	9.8	-	9.8	3.5	-	3.5	2.8	-	2.8
4-1	11.9	-	11.9	11.6	-	11.6	5.6	-	5.6	3.4	-	3.4
4-2	13.8	0.5	14.3	28.7	1.6	30.3	0.5	-	0.5	10.7	3.0	13.7
5-1	8.4	0.5	8.9	-	-	-	3.4	1.4	4.8	-	-	-
5-2	14.2	1.9	16.1	-	-	-	7.4	3.4	10.8	-	-	-
6	22.6	2.4	25.0	-	-	-	18.3	4.6	22.9	-	-	-
7	27.5	2.5	30.0	33.7	1.1	34.8	16.8	6.5	23.3	32.0	6.9	38.9
8	22.9	1.1	24.0	17.0	0.5	17.5	8.9	2.4	11.3	6.5	1.0	7.5
9	54.7	9.1	63.8	43.7	3.5	47.2	24.0	15.9	39.9	28.2	6.4	34.6
10	22.5	4.4	26.9	33.7	3.0	36.7	7.6	7.2	14.8	18.2	4.0	22.2
11-1	61.9	15.5	77.4	54.4	69.8	64.2	18.5	31.5	50.0	48.6	12.2	60.8
11-2	73.9	13.8	87.7	56.7	11.9	68.6	37.2	25.8	63.0	45.3	15.6	60.9
12	16.7	10.6	17.3	-	-	6.5	6.5	-	6.5	-	-	-
13	30.9	4.9	35.8	28.8	1.8	30.6	17.5	4.8	22.3	20.8	-	20.8
14	47.4	3.0	50.4	46.1	2.1	48.2	30.7	6.9	37.6	12.1	4.0	16.1
15	17.8	-	17.8	27.6	-	27.6	4.5	2.8	7.3	7.7	1.4	9.1
16	31.5	-	31.5	33.8	0.5	34.3	15.5	3.2	18.7	18.2	3.6	21.8
17	16.8	-	16.8	19.1	-	19.1	4.2	-	4.2	3.9	1.1	5.0
18	15.7	1.9	17.6	36.2*	-	36.2*	5.0	4.1	9.1	19.0*	2.1*	21.1*
				29.8†	-	29.8†				12.7†	2.5†	15.2†
19	25.1	-	25.1	19.1*	-	19.1*	11.9	6.8	18.7	9.7†	4.4†	14.1†
				21.8†	-	21.8†				11.4†	4.2†	15.6†
20	38.0	1.1	39.1	33.7*	0.5*	34.2*	20.3	5.5	25.8	17.4*	2.9*	20.3*
				38.4†	2.0†	40.4†				23.0†	3.6†	26.6†

* Heavy liming.

† Light liming.

Hay - Park Grass

Yield of hay: cwt per acre

Plot	1945						1946					
	Not limed			Limed			Not limed			Limed		
	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total
1	6.0	4.0	10.0	17.7	2.2	19.9	13.6	6.4	20.0	20.3	3.7	24.0
2	11.8	-	11.8	10.5	-	10.5	12.9	3.0	15.9	14.2	2.3	16.5
3	11.9	-	11.9	11.1	-	11.1	12.3	3.6	15.9	12.8	1.8	14.6
4-1	17.2	1.2	18.4	16.0	0.6	16.6	16.5	3.3	19.8	14.3	3.8	18.1
4-2	15.9	1.5	17.4	26.9	1.5	28.4	19.2	6.1	25.3	43.5	7.9	51.4
5-1	6.7	1.4	8.1	-	-	-	7.3	2.4	9.7	-	-	-
5-2	16.6	3.9	20.5	-	-	-	11.7	5.8	17.5	-	-	-
6	33.5	11.5	45.0	-	-	-	26.2	11.2	37.4	-	-	-
7	30.4	10.0	40.4	48.8	10.1	58.9	22.5	11.0	33.5	52.2	11.9	64.1
8	26.8	5.6	32.4	19.7	3.8	23.5	20.2	8.4	28.6	13.8	6.5	20.3
9	38.2	15.5	53.7	42.2	4.0	46.2	19.5	24.2	43.7	40.5	11.5	52.0
10	30.9	6.9	37.8	34.1	3.4	37.5	18.5	13.4	31.9	30.6	9.0	39.6
11-1	30.0	19.1	49.1	53.5	7.0	60.5	30.6	3.4	34.0	48.1	20.6	68.7
11-2	44.8	23.0	67.8	54.9	10.8	65.7	32.0	3.2	35.2	52.5	2.0	54.5
12	16.8	2.8	19.6	-	-	-	11.1	6.0	17.1	-	-	-
13	37.9	9.1	47.0	52.3	13.8	66.1	24.4	14.2	38.6	36.7	10.4	47.1
14	49.7	11.8	61.5	52.8	6.8	59.6	52.5	12.9	65.4	54.7	11.4	66.1
15	22.0	8.2	30.2	30.2	6.5	36.7	21.3	7.4	28.7	25.9	5.6	31.5
16	17.5	6.2	23.7	34.6	7.9	42.5	35.8	9.3	45.1	37.7	10.8	48.5
17	28.0	7.5	35.5	26.0	5.2	31.2	24.9	7.1	32.0	23.2	5.8	29.0
18	17.0	4.0	21.0	33.2*	-	33.2*	6.5	13.8	20.3	21.9*	3.8*	25.7*
				32.0†	-	32.0†				19.5†	6.7†	26.2†
19	40.3	10.1	50.4	38.8*	5.0*	43.8*	28.7	10.9	39.6	27.4*	7.4*	34.8*
				41.2†	5.5†	46.7†				31.9†	8.2†	40.1†
20	38.6	5.2	43.8	40.6*	3.1*	43.7*	43.5	10.7	54.2	46.6*	7.2*	53.8*
				42.7†	8.8†	51.5†				51.1†	11.9†	63.0†

* Heavy liming.

† Light liming.

Yield of hay: cwt per acre

Plot	1947					
	Not Limed			Limed		
	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total
1	9.0	3.9	12.9	23.2	6.0	29.2
2	18.4	1.1	19.5	21.2	4.2	25.4
3	19.1	1.4	20.5	20.4	2.1	22.5
4-1	24.8	2.6	27.4	23.4	3.4	26.8
4-2	15.4	5.9	21.3	19.9	7.8	27.7
5-1	13.3	1.6	14.9	-	-	-
5-2	28.2	5.0	33.2	-	-	-
6	40.3	8.9	49.2	-	-	-
7	41.2	8.2	49.4	51.4	9.2	60.6
8	29.4	6.0	35.4	24.6	4.6	29.2
9	20.2	22.2	42.4	30.1	9.0	39.1
10	18.0	10.8	28.8	26.4	8.5	34.9
11-1	9.8	22.2	32.0	44.1	11.9	56.0
11-2	18.3	30.9	49.2	52.1	17.2	69.3
12	22.1	3.0	25.1	-	-	-
13	33.8	7.6	41.4	43.2	13.1	56.3
14	45.8	13.8	59.6	40.2	12.0	52.2
15	33.5	6.0	39.5	45.3	7.0	52.3
16	34.9	9.5	44.4	41.2	9.5	50.7
17	23.1	7.4	30.5	26.9	6.8	33.7
18	15.5	3.9	19.4	32.4*	10.5*	42.9*
				28.9†	9.8†	38.7†
19	29.8	6.6	36.4	33.8*	5.2*	39.0*
				33.6†	5.2†	38.8†
20	41.9	8.0	49.9	42.9†	7.1*	50.0*
				37.4†	9.9†	47.3†

* Heavy liming.

† Light liming.

X
Code Numbers of botanical species

	<u>Gramineae</u>	<u>Miscellaneous</u>
1		
1	<i>Agrostis vulgaris</i>	101 <i>Ranunculus acris</i> and <i>bulbosus</i>
2	<i>Aira caespitosa</i>	104 <i>Cerastium vulgatum</i>
3	<i>Alopecurus pratensis</i>	105 <i>Stellaria gramineae</i>
4	<i>Anthoxanthum odoratum</i>	106 <i>Linum catharticum</i>
5	<i>Arrhenatherum avenaceum</i>	107 <i>Agrimonia eupatoria</i>
6	<i>Avena flavescens</i>	109 <i>Potentilla reptans</i>
7	<i>Avena pubescens</i>	110 <i>Poterium sanguisorba</i>
8	<i>Briza media</i>	111 <i>Spiraea ulmaria</i>
9	<i>Bromus mollis</i>	112 <i>Epilobium angustifolium</i>
10	<i>Cynosurus cristatus</i>	113 <i>Anthriscus sylvestris</i>
11	<i>Dactylis glomerata</i>	114 <i>Conopodium denudatum</i>
12	<i>Festuca ovina</i>	115 <i>Heracleum sphondylium</i>
13	<i>Festuca pratensis</i>	116 <i>Pimpinella saxifraga</i>
14	<i>Holcus lanatus</i>	117 <i>Galium verum</i>
15	<i>Lolium perenne</i>	118 <i>Scabiosa arvensis</i>
16	<i>Poa pratensis</i>	119 <i>Achillea millefolium</i>
17	<i>Poa trivialis</i>	120 <i>Centaurea nigra</i>
18	<i>Phleum pratense</i>	122 <i>Hieracium pilosella</i>
	<u>Leguminosae</u>	
51	<i>Lathyrus pratensis</i>	123 <i>Hypochaeris radicata</i>
52	<i>Lotus corniculatus</i>	124 <i>Leontodon hispidus</i>
53	<i>Ononis arvensis</i>	125 <i>Senecio jacobaea</i>
54	<i>Trifolium pratense</i>	126 <i>Taraxacum vulgare</i>
55	<i>Trifolium repens</i>	127 <i>Tragopogon pratensis</i>
56	<i>Vicia sepium</i>	129 <i>Plantago lanceolata</i>
		130 <i>Veronica chamaedrys</i>
		131 <i>Ajuga reptans</i>
		132 <i>Prunella vulgaris</i>
		133 <i>Thymus serpyllum</i>
		134 <i>Rumex acetosa</i>
		135 <i>Rumex crispus</i>
		136 <i>Luzula campestris</i>
		137 <i>Carex praecox</i>
		138 <i>Urtica dioica</i>

X
U = Unlimed

L = Limed

LL = Light lime

HL = Heavy lime

A/6.9

Hay - Park Grass

Sp. No.	Botanical Composition per cent. 1st Crop										Plot 4 1949 U L	
	Plot 1					Plot 2						
	1939		1940		1947		1948		1949			
	U	L	U	L	U	L	U	L	U	L		
1	52.4	3.7	24.1	3.2	76.0	1.1	75.3	1.5	10.0	0.5	2000 0.4	
3	0.2	4.0	0.2	4.0	0.2	1.4	0.1	2.5	9.6	4.1	2.2 2.4	
4	0.7	2.4	0.9	5.6	0.3	6.0	0.2	1.4	1.1	0.4	3.4 1.1	
5	0.1	2.2	0.1	0.8	0.2	4.2	-	3.0	0.4	0.2	7.3 0.1	
6	-	2.2	-	2.4	-	1.3	-	1.7	0.2	1.5	0.9 2.7	
7	-	34.7	-	27.7	0.3	27.4	0.2	12.1	3.5	22.5	5.5 18.1	
8	-	-	-	-	-	-	-	-	1.8	3.9	0.5 1.8	
9	-	-	-	0.1	-	-	-	-	-	-	0.1 -	
11	0.8	13.0	0.2	12.2	2.0	15.9	3.2	18.3	7.9	7.5	7.1 4.9	
12	40.9	15.2	72.2	19.3	14.0	6.1	15.6	15.4	15.5	7.4	9.1 3.6	
13	-	-	-	-	-	-	-	-	0.1	-	0.3 1.1	
14	-	3.5	0.1	3.3	-	5.9	-	5.8	3.3	2.0	6.7 4.5	
15	-	-	-	-	-	-	-	-	-	0.1	0.3 -	
16	0.2	2.3	0.2	2.5	-	1.3	-	1.3	0.1	1.0	0.9 1.6	
17	-	0.2	-	0.9	-	0.3	-	0.3	-	0.3	0.5 1.8	
51	-	1.5	-	1.7	-	1.9	-	2.0	2.2	2.4	3.0 7.3	
52	-	0.5	-	0.6	-	1.2	-	2.4	9.4	9.9	3.8 7.7	
53	-	-	-	-	-	-	-	-	-	-	2.4 -	
54	-	0.1	-	0.2	-	0.6	-	0.2	3.6	4.8	2.3 4.9	
55	-	-	-	-	-	-	-	-	0.4	0.7	2.9 2.5	
101	-	1.1	-	1.2	-	1.1	-	0.8	0.9	2.8	3.8 2.3	
104	-	0.2	-	0.1	-	0.3	-	0.6	0.2	0.3	0.4 -	
105	-	-	-	-	-	-	-	-	0.1	-	- -	
106	-	-	-	-	-	-	-	-	-	0.6	- -	
109	-	-	-	0.2	-	1.9	0.1	-	-	-	- -	
110	-	-	-	-	-	-	-	-	0.6	0.1	0.9 2.1	
111	-	-	-	-	-	-	-	-	-	0.3	- -	
114	-	0.1	-	-	-	0.4	0.1	0.1	3.2	0.2	1.1 -	
116	-	0.5	-	0.8	-	0.1	-	0.4	0.1	0.3	0.1 0.3	
117	2.1	-	-	-	-	-	-	-	1.6	-	- -	
118	-	0.9	-	-	-	0.9	-	2.0	0.5	1.7	0.2 2.2	
119	-	1.1	-	1.7	-	0.5	--	1.7	1.9	1.2	3.8 3.4	
120	0.4	4.7	0.7	1.4	0.5	0.7	0.8	2.0	1.3	2.5	1.6 3.6	
123	-	-	-	-	-	-	-	-	-	0.1	- -	
124	-	0.1	-	-	-	0.8	-	1.3	12.5	8.7	11.0 6.3	
125	-	-	-	-	-	-	-	0.2	-	-	- -	
126	-	0.5	-	0.5	-	0.5	-	1.0	0.4	0.5	0.6 0.1	
127	-	-	-	-	-	0.8	-	-	-	0.2	- 0.7	
129	-	3.5	-	6.6	0.6	10.7	-	19.8	5.7	10.2	8.5 9.7	
130	-	-	-	0.1	-	-	-	-	-	0.1	0.1 0.3	
131	-	-	-	0.1	-	-	-	-	-	-	- -	
134	2.0	1.7	0.8	2.8	5.7	8.5	2.5	2.1	1.4	0.5	5.8 2.1	
136	0.2	0.1	0.5	0.2	-	0.1	0.1	-	0.4	0.5	0.9 0.4	
137	-	-	-	-	-	-	-	-	0.1	-	- -	

Botanical Composition per cent. 1st Crop

Sp. No.	Plot 3				Plot 4				Plot 5 ¹		Plot 5 ²	
	1939		1940		1941		1947		1948		1949	
	U	L	U	L	U	L	U	L	U	L	U	L
1	7.7	1.2	12.1	2.0	1.4		8.4	1.1	15.6	0.8	17.8	7.1
2	-	-	-	-	-		-	-	-	-	0.7	-
3	2.0	3.3	2.6	7.0	4.1		2.4	2.8	7.9	6.2	0.3	9.1
4	1.7	0.3	2.8	0.9	0.3		5.1	2.6	0.6	0.7	1.1	0.1
5	-	0.2	0.7	0.6	-		0.6	0.1	0.1	0.1	0.5	1.8
6	0.3	2.0	0.2	2.0	1.2		0.6	1.9	0.2	1.2	-	-
7	5.2	16.1	5.7	18.2	18.9		3.4	13.5	2.7	13.4	0.5	3.5
8	3.1	4.9	0.9	1.5	3.2		4.7	3.6	0.2	1.0	-	-
9	0.1	-	-	-	-		-	-	-	-	-	-
10	0.1	-	0.3	0.1	0.1		-	-	-	-	-	-
11	3.0	3.1	3.5	3.7	3.7		12.1	3.5	4.5	3.3	4.1	4.7
12	11.3	4.7	19.4	7.7	8.5		10.4	3.0	16.6	4.1	55.8	11.5
14	3.5	1.8	4.6	3.1	2.6		5.3	5.0	4.3	2.5	0.7	4.1
X15	-	0.1	0.1	-	-		-	-	-	-	-	-
16	-	1.3	0.1	2.2	1.8		0.3	1.6	0.2	2.1	0.7	2.6
17	-	0.3	0.1	0.7	1.2		-	0.8	-	0.3	-	-
51	0.5	1.9	0.7	2.3	3.5		2.7	2.5	0.9	2.0	-	16.8
52	4.4	10.9	6.3	13.7	9.6		3.3	5.0	3.8	7.0	3.1	5.2
54	1.6	4.7	2.2	7.0	4.1		4.9	7.5	2.3	6.4	-	1.1
55	0.1	0.3	0.3	0.3	0.5		0.1	0.2	0.3	0.7	-	1.3
56	-	-	-	-	-		-	-	-	-	-	2.1
101	0.1	2.8	0.1	1.9	1.1		1.1	8.6	0.1	3.0	-	0.3
104	0.2	0.2	0.1	0.1	0.3		0.6	0.4	0.2	0.2	-	-
105	0.01X	-	0.1	-	-		-	-	0.2	-	0.2	0.1
106	-	0.8	-	0.2	0.3		-	0.1	-	0.1	-	-
109	-	0.1	-	-	-		-	-	0.3	-	-	-
110	14.6	12.6	9.1	4.1	9.8		5.0	13.4	5.6	11.7	-	-
111	-	-	-	0.3	0.1		-	0.2	-	0.4	-	-
114	2.2	0.3	1.6	-	0.1		5.7	0.9	1.6	0.1	2.0	0.1
115	-	-	-	-	-		-	-	-	-	-	0.6
116	0.3	0.6	0.3	0.6	0.4		-	0.2	0.1	0.2	-	0.3
117	-	-	0.1	-	-		0.1	-	1.1	-	0.6	-
118	1.1	3.7	0.4	3.8	3.2		0.7	2.1	0.9	2.1	1.7	0.9
119	1.4	0.7	1.9	0.8	1.0		0.9	0.5	1.3	1.0	1.6	12.8
120	2.9	1.5	1.3	0.4	0.4		0.6	0.5	1.0	2.0	2.0	5.4
122	-	-	0.1	0.1	-		-	-	-	-	-	-
123	-	-	-	-	-		-	0.1	0.1	-	0.4	-
124	18.2	9.5	12.3	7.0	6.4		12.0	8.8	17.9	12.5	1.6	0.1
126	0.1	0.2	0.2	0.4	-		0.3	0.5	0.1	0.2	-	-
127	0.3	0.1	-	-	-		0.4	0.3	0.1	-	-	-
129	11.8	7.7	6.3	5.3	9.7		3.7	6.2	6.2	13.0	2.4	0.9
130	0.1	0.3	0.3	0.6	0.6		-	0.4	0.1	0.2	-	-
131	-	-	-	0.1	-		-	-	-	-	-	-
132	-	0.1	0.6	0.1	0.1		-	-	-	-	-	-
133	-	-	0.1	-	-		-	-	0.3	-	-	-
134	1.3	1.2	0.2	0.3	1.4		3.7	1.8	2.4	1.3	1.0	5.7
136	0.3	0.4	0.6	0.8	0.3		0.3	0.2	0.1	0.2	1.2	1.8
137	0.4	0.1	1.7	0.1	0.1		0.6	0.1	0.1	-	-	-

A/6.11 X

Hay - Park Grass

Sp. No.	Botanical Composition per cent.								1st Crop			
	Plot 4 ²				Plot 10							
	1947		1949		1939		1940		1947		1948	
	U	L	U	L	U	L	U	L	U	L	U	L
1	68.8	1.8	36.2	2.2	30.4	0.9	33.9	2.1	31.3	1.9	51.9	1.9
3	0.2	32.4	0.7	24.3	0.1	63.8	0.1	50.3	0.4	25.2	0.3	28.6
4	14.5	4.6	10.0	1.2	43.7	0.9	31.5	2.9	52.4	11.3	10.3	1.8
5	-	2.3	-	2.5	1.1	3.5	0.9	7.6	5.4	2.7	5.1	4.1
7	-	-	-	0.3	-	-	-	-	0.1	-	-	-
11	-	0.1	-	0.3	0.1	0.2	0.1	0.1	0.1	0.6	0.3	-
12	9.6	29.8	35.4	57.4	0.7	22.2	2.1	26.3	2.6	43.7	10.2	54.5
14	4.8	1.0	17.5	0.1	23.9	0.2	31.3	0.2	6.6	0.6	21.6	0.5
X16	-	5.3	-	6.3	-	6.0	-	6.2	0.1	3.3	-	3.4
51	-	0.1	-	-	-	-	0.1	-	-	-	-	-
101	-	0.6	-	0.1	-	-	-	-	-	-	-	-
112	0.8	-	-	-	-	-	-	-	-	-	-	-
116	-	-	-	0.1	-	-	-	-	-	-	-	-
117	-	0.8	-	0.4	-	-	-	-	-	1.0	-	-
119	-	0.2	-	0.8	-	-	-	-	-	0.2	-	0.7
126	-	-	-	-	0.3	-	-	-	-	0.5	-	0.4
127	-	-	-	-	-	-	-	-	-	0.1	-	-
129	-	-	-	0.1	-	-	-	-	-	0.4	-	-
134	1.3	20.6	0.2	3.9	-	2.0	-	4.3	1.1	8.4	0.3	5.0
135	-	0.4	-	-	-	-	-	-	-	-	-	-

A/6.12

Sp. No.	Plot 6 1949 U	Botanical Composition per cent. 1st Crop										Plot 15 1949 U	Plot 15 1949 L		
		1939		1940		1947		1948							
		U	L	U	L	U	L	U	L	U	L				
1	2.5	7.2	0.1	3.1	0.2	4.7	0.1	3.8	0.4	2.8	0.4				
3	6.1	1.2	15.6	6.1	22.9	4.5	15.7	8.2	13.2	18.5	6.8				
4	2.2	1.5	0.1	2.6	0.1	6.7	0.3	3.6	0.3	1.7	0.3				
5	2.2	1.5	11.0	3.1	10.9	4.0	19.2	2.4	11.7	0.6	13.9				
6	0.1	0.5	2.2	0.6	2.5	0.8	2.1	1.0	2.5	0.2	0.2				
7	3.1	1.8	5.9	1.8	2.7	1.2	3.9	1.1	5.2	1.1	12.6				
9	0.2	-	2.2	0.2	7.3	-	0.2	0.2	2.1	-	1.1				
11	11.8	21.8	18.9	26.3	13.8	20.8	20.7	15.7	13.2	8.4	4.4				
12	5.3	7.7	2.6	6.5	1.2	4.1	1.0	4.6	1.0	7.0	3.7				
13	-	-	1.6	-	0.4	0.2	3.9	0.3	3.3	-	-				
14	2.6	1.0	1.4	5.8	0.9	1.8	0.6	2.8	1.2	1.4	1.6				
15	-	-	-	-	-	-	-	-	0.2	-	-				
16	1.7	1.1	1.5	1.7	1.3	2.1	1.3	2.6	1.2	0.5	1.2				
17	-	-	5.6	0.3	9.3	0.2	2.0	0.3	1.1	-	0.4				
51	20.7	28.8	13.1	8.8	6.4	11.4	5.2	11.3	15.6	22.3	13.4				
52	2.6	2.3	0.1	2.6	-	0.3	-	0.5	-	0.9	-				
54	5.3	4.6	1.4	4.9	2.1	4.3	0.1	4.6	3.4	1.7	2.8				
55	2.2	3.9	7.8	3.0	5.7	1.6	3.8	3.4	6.2	2.8	16.5				
56	0.5	-	-	-	-	-	-	-	-	-	-				
101	0.8	0.8	1.2	1.0	1.2	1.3	3.2	0.7	1.9	0.4	1.8				
104	-	-	0.1	-	-	-	0.1	-	-	-	0.1				
111	-	0.8	-	2.3	-	-	-	-	-	-	-				
113	-	-	0.2	-	-	-	-	-	0.3	-	-				
114	4.1	1.4	0.2	2.0	0.1	4.4	0.1	1.4	0.1	1.1	0.4				
115	0.6	1.4	1.2	4.8	4.4	1.1	4.6	1.0	4.5	-	3.2				
117	0.1	-	-	0.1	-	-	-	0.2	-	-	-				
118	1.1	0.1	0.7	-	1.6	-	-	-	0.3	-	-				
119	8.9	2.8	0.2	1.2	0.1	7.6	0.3	14.0	1.0	10.0	1.2				
120	5.5	4.0	0.8	3.1	0.1	7.0	1.5	4.3	0.5	7.6	1.4				
124	0.2	0.4	-	-	-	0.1	-	-	-	0.3	-				
126	0.2	-	1.7	0.1	3.3	0.1	1.6	0.4	1.9	0.1	1.9				
127	0.3	0.1	-	-	0.1	0.9	3.5	1.0	2.6	-	0.3				
129	7.4	1.6	1.0	1.3	0.5	5.4	1.8	6.5	4.6	9.7	9.6				
130	-	-	0.1	-	0.1	0.1	0.1	-	0.1	-	0.1				
134	1.5	1.3	1.5	5.6	0.8	2.3	3.1	2.2	0.4	0.8	0.6				
136	0.2	0.4	-	1.1	-	0.1	-	0.2	-	0.1	0.1				
138	-	-	-	-	-	0.9	-	1.7	-	-	-				

Hay - Park Grass

Botanical Composition per cent. 1st Crop

Sp. No.	Plot 8										Plot 16	
	1939		1940		1941		1947		1948		1949	
	U	L	U	L	U	L	U	L	U	L	U	L
1	5.9	0.8	4.0	1.3	2.4	0.7	4.2	0.9	3.0	1.3	2.3	0.2
3	0.4	1.5	0.8	2.8	0.4	1.5	1.1	2.7	3.6	2.7	22.5	10.9
4	1.2	1.0	2.3	1.4	2.8	0.9	5.8	5.6	2.4	1.4	4.2	0.2
5	10.8	11.6	5.4	9.5	4.8	8.2	12.4	8.3	14.2	14.7	22.0	19.4
6	1.8	4.2	1.7	3.5	0.7	2.4	1.4	2.0	0.5	2.1	0.6	0.6
7	3.5	16.1	3.6	14.7	4.3	20.5	1.7	16.7	1.9	11.3	5.8	15.3
8	0.2	1.5	0.1	0.7	-	1.4	0.2	2.4	0.1	0.9	-	-
9	-	0.1	-	-	-	-	-	-	-	-	0.2	0.6
10	-	-	-	0.1	-	-	-	-	-	-	-	-
11	4.5	4.7	3.7	4.7	2.9	4.8	10.9	6.2	12.6	5.6	9.8	13.4
12	9.5	4.6	11.6	6.8	23.6	8.7	4.6	1.5	5.2	2.1	6.2	14.4
13	-	4.0	0.1	1.6	-	2.1	6.3	6.3	0.4	6.9	-	-
14	6.2	3.5	7.8	3.6	4.3	2.4	6.5	4.2	7.4	2.1	1.1	-
15	0.2	0.3	0.2	0.2	-	0.1	0.3	-	0.1	-	-	-
16	0.7	1.1	0.6	1.3	0.7	1.8	1.4	1.7	1.1	1.5	0.4	0.5
17	0.7	1.1	0.6	3.0	0.3	3.2	0.3	1.7	0.1	0.8	0.1	0.4
51	1.6	0.8	0.3	0.5	0.6	0.6	0.4	0.8	0.6	1.5	11.7	8.5
52	4.3	5.4	6.1	6.2	3.1	5.2	1.8	2.1	3.3	5.9	-	-
54	8.1	8.6	17.9	17.3	6.2	8.5	6.5	7.3	2.7	3.5	-	0.7
55	0.5	0.1	0.9	-	1.8	0.2	0.5	0.1	0.7	0.1	0.2	0.4
101	2.3	0.8	2.7	0.8	1.4	0.6	9.8	4.0	5.7	2.1	0.2	0.4
104	0.5	0.5	0.2	0.1	0.7	0.8	0.4	0.3	0.1	0.4	-	-
105	0.2	-	0.1	-	0.1	0.1	0.1	-	0.1	-	-	-
107	-	0.8	-	-	-	-	-	-	-	-	-	-
111	-	-	-	1.0	-	1.1	-	0.6	-	0.6	-	-
113	-	-	-	-	-	-	-	-	-	-	-	1.5
114	0.7	0.1	0.9	-	0.6	0.1	2.1	0.1	0.8	0.2	0.7	0.4
115	0.1	0.1	-	-	-	-	-	-	-	-	-	0.9
116	0.2	0.6	0.3	0.4	0.3	0.5	-	0.2	0.1	0.2	-	-
117	0.2	0.1	0.2	0.1	1.5	0.1	0.2	0.1	0.4	0.2	-	-
118	2.2	3.5	0.9	4.6	1.4	5.6	0.7	4.7	0.9	3.6	-	-
119	4.3	0.8	3.4	0.7	6.3	1.4	4.1	1.2	6.9	4.0	3.9	1.1
120	4.0	5.4	2.8	1.7	2.1	2.6	1.0	1.0	2.0	2.5	-	-
124	4.4	3.2	2.0	1.6	3.0	1.5	4.9	3.4	4.5	5.6	-	-
126	0.2	-	0.4	0.1	0.3	0.2	0.4	0.2	0.1	0.1	0.9	3.3
127	-	-	-	-	-	0.1	-	0.7	-	0.9	0.7	1.4
129	13.0	10.7	11.1	6.8	18.6	9.9	10.0	9.5	15.0	13.1	6.3	5.3
130	0.3	0.3	0.5	0.8	0.3	0.6	0.2	0.6	0.3	0.2	-	-
131	-	-	-	0.1	-	-	-	-	-	-	-	-
132	-	-	-	-	-	0.1	-	-	-	-	-	-
134	6.8	1.8	4.0	1.4	2.5	1.2	5.7	2.6	2.9	1.8	0.2	0.2
136	0.5	0.3	2.8	0.6	2.0	0.3	0.2	0.3	0.3	0.1	-	-
138	-	-	-	-	-	-	0.2	-	-	-	-	-

A/6.14

Botanical Composition per cent. 1st Crop

Plot 9

Sp. No.	1939		1940		1941		1947		1948	
	U	L	U	L	U		U	L	U	L
1	2.0	0.8	5.3	2.6	5.6		23.8	3.4	7.8	4.3
3	0.1	69.4	0.1	55.0	-		0.8	31.8	-	38.1
4	3.2	0.7	0.3	2.4	8.0		15.7	12.4	0.4	4.2
5	0.3	14.3	0.9	21.9	0.2		4.3	13.2	0.6	14.7
6	-	-	-	-	-		-	-	-	0.1
7	-	-	-	0.2	-		-	0.5	-	0.5
9	-	0.1	-	1.0	-		-	0.1	-	0.1
11	-	4.3	-	4.1	0.2		0.2	12.5	-	11.6
12	0.1	1.9	0.1	4.7	0.1		0.1	2.8	0.1	4.3
14	94.1	2.5	93.3	2.1	85.1		51.3	3.4	90.6	2.5
16	-	4.4	-	2.1	-		0.1	3.8	-	9.4
17	-	-	-	0.1	-		-	-	-	-
51	-	0.8	-	0.7	-		-	6.0	-	3.5
55	-	-	-	-	0.1		-	-	-	-
112	-	-	-	-	-		1.6	-	0.5	-
113	-	0.1	-	0.2	-		-	0.1	-	-
114	-	-	-	-	-		-	0.8	-	0.1
115	-	0.2	-	2.0	-		-	1.4	-	1.5
119	-	-	-	-	-		-	0.2	-	0.5
124	-	-	-	-	-		0.1	-	-	-
126	-	0.5	-	0.6	-		0.1	4.8	-	3.3
127	-	-	-	-	-		-	0.3	-	0.1
129	0.1	-	-	-	-		0.1	0.2	-	0.1
134	0.1	-	-	0.3	0.7		1.8	2.3	-	1.1

A/6.15

Hay - Park Grass

Sp. No.	Botanical Composition per cent. 1st Crop										
	Plot 11 ¹				Plot 11 ²				Plot 12 1949 U	Plot 17 1949 U	
	1947 U	1947 L	1949 U	1949 L	1947 U	1947 L	1949 U	1949 L			
1	4.5	0.8	-	-	44.1	0.2	5.3	-	3.5	1.2	0.6
3	1.0	78.9	0.1	82.0	0.8	70.2	0.3	57.5	7.6	14.5	7.3
4	0.8	0.7	-	-	0.1	0.1	-	-	3.4	8.7	0.8
5	0.3	2.9	-	2.3	12.7	11.3	0.8	17.5	0.8	-	1.7
6	-	-	-	-	-	-	-	-	0.2	-	1.5
7	-	-	-	-	-	-	-	-	4.1	1.7	20.5
8	-	-	-	-	-	0.1	-	-	6.0	0.6	0.7
9	-	0.1	-	-	-	-	-	-	-	-	0.7
11	-	2.5	-	4.9	-	7.8	-	10.3	8.8	25.4	21.1
12	0.2	-	-	0.4	0.2	-	0.1	-	19.4	9.4	22.3
13	-	-	-	-	-	-	-	-	1.7	-	0.8
14	81.1	7.6	99.6	3.5	40.8	2.4	92.8	1.6	2.8	8.4	2.1
15	-	-	-	-	-	-	-	-	-	0.6	2.4
16	-	4.5	--	5.1	0.1	6.3	-	10.6	0.2	-	0.1
17	-	-	-	-	-	0.1	-	-	-	-	0.6
51	-	-	-	-	-	0.1	-	-	1.1	0.1	-
52	-	-	-	-	-	-	-	-	7.4	-	2.8
54	0.2	-	-	-	-	-	-	-	4.6	-	-
55	-	-	-	-	-	-	-	-	0.2	-	-
101	-	-	-	-	-	-	-	-	0.7	0.4	2.1
104	-	-	-	-	-	-	-	-	0.1	0.1	0.1
105	-	-	-	-	-	-	-	-	0.2	-	-
109	-	-	-	-	-	-	-	-	0.1	-	-
111	-	-	-	-	-	-	-	-	0.2	-	-
112	11.9	-	0.3	-	-	-	-	-	-	-	-
114	-	0.1	-	-	-	-	-	-	1.3	0.4	0.1
115	-	-	-	-	-	0.1	-	0.1	-	-	0.2
116	-	-	-	-	-	-	-	-	0.1	-	-
117	-	-	-	-	-	-	-	-	0.9	-	0.4
118	-	-	-	-	-	-	-	-	1.7	-	-
119	-	-	-	-	-	-	-	-	1.8	1.2	1.0
120	-	-	-	-	-	-	-	-	1.8	5.5	1.4
123	-	-	-	-	-	-	-	-	0.1	-	-
124	-	-	-	-	-	-	-	-	10.0	4.0	1.7
126	-	1.6	-	0.5	-	1.3	-	1.3	0.2	1.4	0.6
127	-	-	-	-	-	-	-	-	0.1	-	0.5
129	-	-	-	-	-	-	-	-	6.8	14.0	5.6
130	-	-	-	-	-	-	-	-	0.1	-	-
134	-	0.3	-	1.3	1.2	-	0.7	1.1	1.0	0.6	0.2
136	-	-	-	-	-	-	-	-	0.6	-	0.1
137	-	-	-	-	-	-	-	-	0.4	1.8	-

Botanical Composition per cent. 1st Crop
 Plot 13

Sp. No.	1944		1945		1946		1947		1948	
	U	L	U	L	U	L	U	L	U	L
1	8.4	-	6.3	-	8.0	0.1	10.9	-	15.7	0.2
3	56.7	5.8	46.0	6.8	32.5	8.6	27.6	13.7	31.9	10.4
4	4.8	0.5	4.9	0.3	15.5	1.2	14.2	0.8	6.3	0.3
5	0.9	5.9	2.1	7.0	1.4	10.9	2.1	14.2	3.4	25.9
6	0.1	-	-	-	-	-	-	-	-	0.1
7	-	-	-	0.2	-	0.4	0.1	0.7	-	0.3
9	0.1	0.1	-	0.1	-	0.1	-	-	-	0.1
11	6.7	8.6	5.6	7.8	6.8	27.5	9.8	24.2	9.1	21.5
12	3.6	1.1	3.4	1.6	6.0	1.1	4.5	1.0	4.0	0.9
13	-	-	-	-	-	-	-	0.1	-	-
14	2.7	1.8	4.5	2.1	5.7	5.4	6.2	5.3	2.5	4.4
15	0.1	-	-	-	-	-	-	-	-	-
16	1.3	1.4	0.8	1.9	0.1	1.4	0.9	2.3	0.9	1.8
17	0.4	1.2	0.1	8.2	0.1	5.8	0.2	4.6	0.1	1.2
51	0.3	31.0	0.5	26.6	0.2	4.4	0.7	6.3	0.5	7.9
52	-	0.2	-	0.3	-	0.1	-	-	-	-
54	-	7.3	-	6.0	0.1	2.4	0.1	1.2	-	2.4
55	-	2.6	-	4.0	-	0.8	-	-	-	0.3
101	-	0.9	-	2.1	0.1	1.0	0.6	2.3	0.1	2.2
104	0.3	0.5	-	0.2	0.1	0.1	0.1	0.2	0.1	0.2
105	0.2	-	0.6	-	0.6	0.1	0.2	-	0.1	-
113	0.2	2.0	-	-	-	0.3	-	0.2	-	0.7
114	0.9	-	1.6	0.2	1.0	0.1	2.0	-	0.8	0.1
115	0.2	-	1.3	-	-	-	-	-	-	-
117	-	-	-	-	0.4	-	-	-	-	-
119	0.6	0.2	3.0	0.5	2.9	0.4	3.8	0.5	4.0	0.3
120	0.3	4.0	-	0.4	0.5	0.4	0.7	1.6	0.2	0.7
123	0.2	-	-	-	-	-	-	-	-	-
124	-	0.9	0.3	1.2	0.4	1.3	0.3	1.5	1.7	1.3
126	0.3	6.2	0.7	3.6	1.0	4.1	0.9	5.4	1.5	3.2
127	-	0.5	-	2.2	-	1.0	-	0.4	0.3	1.3
129	6.2	16.2	12.1	14.5	14.0	19.3	10.3	10.8	14.7	11.5
130	0.5	0.7	0.4	0.9	0.9	0.4	0.7	0.2	0.2	-
131	-	0.1	-	-	-	-	0.2	-	-	-
134	3.9	0.3	5.8	1.3	1.7	1.3	2.9	2.5	1.9	0.8
136	0.1	-	-	-	-	-	-	-	-	-

Hay - Park Grass

Botanical Composition per cent. 1st Crop

Plot 14

Sp. No.	1939						1940						1941						1947						1948						
	U	L	L	sun	shade		U	L	L	sun	shade		U	L	L	sun	shade		U	L	L	sun	shade		U	L	L	sun	shade		
1	-	-	0.1	0.1	0.1	0.2	-	-	-	0.1	0.5	0.2	1.0	0.6	0.1	0.4															
3	53.1	23.9	13.7	49.1	18.5	12.4	59.2	23.1	18.5	27.6	10.5	5.0	31.8	12.1	6.7																
4	-	-	0.4	-	-	0.5	-	-	-	0.5	0.2	0.3	0.5	0.6	-	0.2															
5	31.1	48.2	12.4	30.9	52.0	13.1	28.9	44.5	18.7	36.8	43.6	27.1	36.2	45.0	34.7																
6	-	-	0.6	-	0.1	0.6	-	-	0.2	0.4	-	0.1	1.2	0.2	0.5	1.1															
7	0.6	0.9	12.6	0.1	0.7	11.9	0.1	0.8	14.4	0.5	3.9	12.6	1.4	5.3	11.6																
9	0.1	0.4	0.3	-	0.1	0.6	0.1	0.3	0.6	-	-	-	-	0.3	-	-															
11	6.3	4.3	1.0	5.7	6.0	1.9	7.2	7.2	1.1	14.5	10.7	5.2	14.2	13.6	5.0																
12	0.2	4.8	38.0	0.1	5.3	36.6	-	11.4	34.6	-	9.0	27.1	-	13.4	27.4																
14	0.2	-	0.2	-	-	0.1	-	-	-	0.8	-	0.3	-	-	2.3																
15	-	-	0.1	-	-	-	-	-	-	0.1	-	-	-	-	-																
16	1.2	1.9	3.6	0.3	1.0	1.6	0.7	1.3	3.1	3.9	2.9	222	4.7	2.9	2.9	2.9															
17	3.5	7.3	3.1	9.8	10.5	2.3	1.5	3.9	1.5	0.3	0.5	0.9	2.4	1.0	1.0																
51	1.6	5.6	12.1	0.7	1.5	15.0	0.3	4.0	4.9	3.3	13.2	12.2	2.1	3.2	3.6																
54	-	-	-	-	-	-	-	-	0.1	-	-	0.1	-	-	0.3																
55	-	-	0.3	-	-	-	-	-	-	-	-	-	-	-	0.5																
101	-	-	0.1	-	-	0.1	-	-	-	-	-	0.1	0.1	-	-																
107	-	-	-	-	-	-	-	-	0.3	-	-	-	-	-	-																
113	0.6	1.2	0.6	1.3	2.3	1.9	0.8	0.1	0.6	5.3	0.8	0.3	0.2	0.7	0.5																
114	-	-	0.1	-	-	-	-	-	-	-	-	-	-	-	-																
116	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1																
119	-	-	0.2	-	-	0.1	-	-	-	-	-	-	-	-	0.1																
120	-	-	-	-	-	0.1	-	-	-	0.2	-	-	-	-	-																
126	0.1	0.2	0.2	0.6	0.9	0.7	0.7	2.0	0.6	3.0	2.9	3.6	3.2	1.1	1.7																
127	-	-	-	-	-	-	-	-	-	-	0.9	-	-	0.5	-																
129	0.1	-	-	-	-	0.2	-	0.1	-	1.1	0.3	0.1	1.6	-	-																
134	1.3	1.3	0.3	1.3	1.0	0.1	0.5	0.8	0.2	2.0	0.1	0.1	0.4	0.6	-																

Sp. No.	Botanical Composition per cent.						1st Crop					
	Plot 18			Plot 19			Plot 18			Plot 19		
	1946 U	1946 LL	1946 HL	1948 U	1948 LL	1948 HL	1946 U	1946 LL	1946 HL	1948 U	1948 LL	1948 HL
1	74.6	2.0	1.4	76.9	3.4	1.2	8.1	2.8	0.1	11.8	4.3	0.4
2	-	-	3.1	-	-	-	-	-	-	-	-	-
3	1.8	2.9	2.9	0.1	2.8	3.8	13.1	18.3	19.7	6.2	25.2	16.7
4	4.2	2.3	0.2	0.8	0.8	-	9.1	2.7	0.1	7.3	1.5	-
5	-	10.1	10.3	0.5	24.6	25.0	9.8	6.4	13.9	5.7	5.5	14.3
6	-	-	-	-	-	-	1.4	0.5	2.4	1.7	3.0	3.1
7	-	1.0	0.7	-	3.2	0.8	0.5	1.7	2.7	0.9	1.9	5.4
9	-	0.4	0.1	-	0.3	0.1	-	0.5	0.1	-	0.2	0.3
11	-	12.5	36.5	1.3	35.0	48.3	3.6	4.4	4.3	3.3	10.9	13.6
12	12.5	9.8	5.1	8.9	6.8	2.2	4.6	2.5	3.3	9.8	5.0	4.5
13	-	-	-	-	0.1	-	2.4	-	4.1	1.1	0.4	5.7
14	0.5	0.3	0.3	-	0.2	-	2.7	2.0	0.4	1.3	1.7	-
15	-	0.1	0.1	-	0.1	0.1	-	-	-	-	-	0.3
16	--	1.1	1.0	-	0.8	1.1	0.2	0.9	1.4	0.6	0.9	1.9
17	0.1	0.6	0.3	-	0.2	0.2	0.5	1.0	1.3	0.5	1.8	0.7
18	-	1.6	-	-	2.6	0.1	-	-	-	-	-	-
51	-	0.1	1.4	-	-	0.5	9.5	10.5	13.2	13.5	7.4	7.5
52	-	-	-	-	-	-	-	-	2.6	1.1	-	1.7
54	-	0.4	0.1	-	0.1	0.1	0.4	1.2	0.8	0.2	-	0.5
55	-	0.1	-	-	-	-	0.9	2.9	1.5	2.7	1.0	0.7
101	-	-	-	-	-	-	5.9	10.4	2.2	5.7	7.9	4.0
104	-	0.6	0.1	-	0.1	-	0.3	0.4	0.1	0.2	-	-
105	-	-	-	-	-	-	-	-	-	0.2	-	-
107	-	0.2	-	-	-	-	-	-	-	-	-	-
111	-	-	-	-	0.1	-	-	-	-	-	-	-
113	-	-	-	-	-	-	-	0.4	0.1	-	-	-
114	-	0.1	-	-	0.3	0.1	0.4	0.4	0.2	0.3	0.4	-
115	-	1.5	2.4	-	0.6	1.6	0.1	1.5	1.7	1.0	0.4	1.1
117	-	0.4	-	-	-	-	-	-	-	-	-	-
119	-	0.1	0.5	-	0.2	-	6.3	5.4	1.9	10.2	5.2	2.3
120	-	10.2	0.9	-	3.5	--	-	0.7	1.8	-	1.0	1.4
123	-	-	-	-	-	-	0.3	-	-	-	-	-
124	-	0.1	-	-	0.1	-	0.1	-	-	0.4	-	-
126	-	22.6	23.4	0.2	8.9	10.8	1.3	2.5	1.7	0.6	0.6	1.0
127	-	1.5	3.4	-	1.2	2.6	-	0.1	0.8	-	0.4	2.9
129	0.1	17.3	4.5	-	3.8	1.0	15.8	14.2	16.3	11.8	10.4	9.7
130	-	-	-	-	-	-	0.9	1.8	0.9	0.2	0.7	-
131	-	-	-	-	-	-	-	-	-	0.2	-	-
134	6.2	0.5	0.9	11.2	0.2	0.4	1.8	3.9	0.4	1.3	2.3	0.3
136	-	-	-	-	-	-	-	-	-	0.2	-	-
138	-	-	-	0.1	-	-	-	-	-	-	-	-

A/6.19

Hay - Park GrassBotanical Composition per cent. 1st Crop
Plot 20

Sp. No.	1946			1948		
	U	LL	HL	U	LL	HL
1	2.9	1.5	0.1	4.1	2.5	0.1
3	33.9	18.0	12.6	39.2	21.8	17.6
4	1.3	3.2	0.7	1.1	1.9	0.2
5	9.9	26.9	15.5	15.1	21.7	17.4
6	1.5	2.3	2.6	4.0	1.9	3.3
7	2.6	4.0	9.1	0.7	3.0	6.8
9	-	0.3	0.2	0.2	0.1	0.2
11	10.3	7.9	4.3	14.6	14.4	14.2
12	4.8	3.6	9.4	3.7	2.8	4.8
13	0.5	-	0.5	-	-	-
14	3.0	2.9	1.4	1.0	5.7	1.6
15	0.1	0.2	0.2	-	0.1	0.8
16	0.3	0.3	1.2	1.3	0.4	3.0
17	0.9	4.0	3.6	1.3	3.9	4.3
51	5.5	4.8	7.1	4.3	2.1	8.8
52	0.3	-	0.4	0.1	-	-
54	-	-	0.6	-	0.1	0.2
55	-	0.7	3.0	0.6	2.8	0.8
101	5.0	1.2	1.9	1.2	1.9	1.3
104	0.1	0.2	0.3	-	0.3	-
113	0.7	2.7	0.7	0.5	-	0.3
114	0.2	0.1	0.1	-	-	-
115	-	-	0.6	0.1	--	0.2
117	-	-	0.3	-	-	0.4
119	1.7	1.8	3.7	0.4	1.8	2.7
120	-	1.9	2.8	-	0.7	0.2
126	1.3	2.6	3.0	0.7	1.1	2.3
127	3.7	0.2	3.1	3.3	1.1	3.8
129	3.2	6.7	6.3	0.9	6.2	4.1
130	1.2	0.5	3.0	0.1	0.3	0.1
134	5.1	1.5	1.7	1.5	1.4	0.5

A/6.20

Botanical Composition per cent. 1st Crop

Plot	1939				1940			
	Gram- ineae	Legum- inosae	Other orders	Dom. Sp.	Gram- ineae	Legum- inosae	Other orders	Dom. Sp.
1 U	95.3	-	4.7	117 134	98.0	-	2.0	ND
1 L	83.4	2.1	14.5	120	82.0	2.5	15.5	129
3 U	37.9	6.7	55.4	124	53.1	9.6	37.3	124 110
3 L	39.4	17.8	42.8	110	49.8	23.2	27.0	124 129
7 U	45.3	39.7	15.0	120	58.1	19.3	22.6	134 115
7 L	68.7	22.4	8.9	126	73.6	14.2	12.2	115 126
8 U	45.6	14.4	40.0	129	42.4	25.3	32.3	129
8 L	56.2	14.8	29.0	129	55.2	24.0	20.8	129 118
9 U	99.9	-	0.1	ND	100.0	-	-	-
9 L	98.4	0.8	0.8	126	96.2	0.7	3.1	115
10 U	100.0	-	-	-	99.9	0.1	-	-
10 L	97.6	-	2.4	134	95.7	-	4.3	134
14 U	96.3	1.6	2.1	134	96.0	0.7	3.3	113 134
14 L Sun	91.7	5.6	2.7	134	94.3	1.5	4.2	134 113
14 L Shade	86.1	12.5	1.4	ND	81.9	15.0	3.1	113
18 U	98.6	0.1	1.3	134	98.3	0.1	1.6	ND
18 LL	68.9	0.2	30.9	126	62.2	0.2	37.6	126
18 HL	73.6	0.3	26.1	126	61.9	0.4	37.7	126
19 U	65.6	19.9	14.5	ND	80.6	7.0	12.4	ND
19 LL	59.1	26.3	14.6	ND	84.1	3.9	12.0	ND
19 HL	72.8	19.0	8.2	ND	87.4	6.7	5.9	ND
20 U	84.4	9.0	6.6	134	83.6	5.9	10.5	ND
20 LL	65.5	23.4	11.1	ND	84.7	5.4	9.9	ND
20 HL	68.1	15.6	16.3	ND	79.2	7.0	13.8	ND

U - Unlimed; L - Limed; LL - Light lime; HL - Heavy lime; ND - None

Dominant

Columns headed "Dom. Sp." give the code numbers of the dominant non-gramineous or leguminous species.

A/6.21

Hay - Park Grass

Botanical Composition per cent. 1st Crop

Plot	1941				1942			
	Gram- ineae	Legum- inosae	Other orders	Dom. Sp.	Gram- ineae	Legum- inosae	Others orders	Dom. Sp.
1 U	95.9	0.2	3.9	ND	96.7	0.1	3.2	ND
1 L	73.4	3.7	22.9	ND	72.6	3.5	23.9	ND
3 U	46.8	12.4	40.8	ND	48.1	8.0	43.9	110
3 L	47.0	17.7	35.3	129 110	52.2	11.1	36.7	129
7 U	22.8	28.5	48.7	ND	67.4	9.8	22.8	ND
7 L	63.2	25.7	11.1	ND	63.9	11.4	24.7	ND
8 U	47.2	11.7	41.1	129	40.4	6.9	52.7	129
8 L	58.7	14.5	26.8	129 118	64.8	9.9	25.3	ND
9 U	99.2	0.1	0.7	ND	99.6	-	0.4	ND
9 L	98.9	0.3	0.8	ND	94.6	0.1	5.3	ND
10 U	98.3	-	1.7	ND	99.4	0.1	0.5	ND
10 L	98.6	0.1	1.3	ND	90.7	-	9.3	ND
14 U	97.6	0.3	2.1	ND	94.7	0.3	5.0	ND
14 L Sun	92.6	4.1	3.3	126	91.7	0.9	7.4	ND
14 L Shade	93.5	5.0	1.5	ND	91.9	3.7	4.4	ND
18 U	96.7	0.1	3.2	134	90.3	-	9.7	ND
18 LL	57.1	0.3	42.6	ND	62.7	0.7	36.6	126
18 HL	59.7	0.1	40.2	ND	62.7	0.4	36.9	ND
19 U	82.3	5.9	11.8	ND	77.7	2.4	19.9	ND
19 LL	86.7	4.3	9.0	ND	79.5	1.5	19.0	ND
19 HL	90.8	4.2	5.0	ND	94.7	0.5	4.8	ND
20 U	86.1	6.1	7.8	ND	67.0	3.3	29.7	ND
20 LL	82.4	3.7	13.9	ND	81.7	0.6	17.7	ND
20 HL	90.8	2.3	6.9	ND	87.0	1.1	11.9	ND

U - Unlimed; L - Limed; LL - Light lime; HL - Heavy lime; ND - None Dominant



Botanical Composition per cent. 1st Crop

Plot	1943				1944			
	Gram- ineae	Legum- inosae	Other orders	Dom. Sp.	Gram- ineae	Legum- inosae	Other orders	Dom. Sp.
1 U	99.0	-	1.0	ND	96.5	0.2	3.3	120
1 L	70.3	2.5	27.2	129	65.7	2.0	32.3	ND
3 U	54.2	8.2	37.6	ND	60.7	5.3	34.0	ND
3 L	41.3	12.7	46.0	110	28.4	17.3	54.3	110
7 U	54.0	19.4	26.6	114	38.2	31.9	29.9	ND
7 L	71.1	16.6	12.3	ND	52.3	34.3	13.4	ND
8 U	46.6	14.6	38.8	ND	35.0	16.8	48.2	ND
8 L	48.3	19.6	32.1	ND	46.4	20.2	33.4	ND
9 U	100.0	-	-	-	100.0	-	-	-
9 L	97.9	0.1	2.0	ND	95.8	-	4.2	115
10 U	99.8	0.1	0.1	ND	99.8	0.1	0.1	ND
10 L	98.7	-	1.3	ND	93.5	-	6.5	134
13 U	not sampled				85.8	0.3	13.9	129
13 L					26.4	41.1	32.5	129
14 U	98.1	0.5	1.4	ND	98.3	0.6	1.1	ND
14 L Sun	97.1	1.3	1.6	ND	93.8	4.0	2.2	ND
14 L Shade	95.1	2.6	2.3	ND	94.9	2.5	2.6	126
18 U	98.7	0.1	1.2	134	80.2	0.3	19.5	134
18 LL	78.8	0.1	21.1	126	43.2	0.9	55.9	ND
18 HL	85.9	0.4	13.7	126	64.1	0.6	35.3	126
19 U	75.5	7.7	16.8	ND	62.4	8.0	29.6	ND
19 LL	79.7	4.5	15.8	ND	67.6	10.7	21.7	ND
19 HL	79.0	9.3	11.7	ND	76.7	5.6	17.7	ND
20 U	91.7	2.3	6.0	ND	88.6	2.8	8.6	ND
20 LL	90.1	1.4	8.5	115	84.0	1.1	14.9	ND
20 HL	83.8	1.5	14.7	ND	78.4	2.7	18.9	ND

U - Unlimed; L - Limed; LL - Light lime; HL - Heavy lime; ND - None Dominant

A/6.23

Hay - Park Grass

Botanical Composition per cent. 1st Crop

Plot	1945				1946			
	Gram- ineae	Legum- inosaee	Other orders	Dom. Sp.	Gram- ineae	Legum- inosaee	Other orders	Dom. Sp.
1 U	95.1	0.1	4.8	109	98.5	-	1.5	ND
1 L	53.2	6.9	39.9	129	64.8	6.2	29.0	ND
3 U	52.8	12.4	34.8	110 124	45.6	14.0	40.4	124
3 L	28.3	20.9	50.8	ND	27.2	21.0	51.8	124
7 U	26.1	40.1	33.8	120	28.6	25.9	45.5	ND
7 L	60.7	20.4	18.9	ND	65.4	15.5	19.1	127
8 U	31.7	17.5	50.8	129. 124 120	27.3	16.0	56.7	124
8 L	33.2	24.2	42.6	ND	33.0	24.0	43.0	124 129
9 U	100.0	-	-	-	99.6	-	0.4	ND
9 L	89.7	3.4	6.9	115	95.6	0.2	4.2	ND
10 U	100.0	-	-	-	99.6	0.1	0.3	134
10 L	97.5	-	2.5	134	93.3	-	6.7	134
13 U	73.7	0.5	25.8	129	76.0	0.3	23.6	129
13 L	36.1	36.9	27.0	129	62.5	7.7	29.8	129
14 U	97.7	1.6	0.7	126	92.5	2.1	5.4	113 126
14 L	90.2	6.6	3.2	126	87.6	9.2	3.2	126
14 L	92.4	4.4	3.2	126	89.8	7.4	2.8	126
18 U	83.9	-	16.1	134	93.7	-	6.3	134
18 LL	52.4	0.1	47.5	126 129	44.7	0.6	54.7	126
18 HL	62.3	1.2	36.5	126 127	61.9	1.5	36.6	126
19 L	52.9	16.3	30.8	ND	56.0	10.8	33.2	129
19 LL	50.2	22.5	27.3	129	43.7	14.6	41.7	129
19 HL	70.0	15.4	14.6	ND	53.9	18.0	28.1	129
20 L	65.4	18.4	16.2	119	72.1	5.7	22.2	134
20 LL	65.3	12.5	22.2	119	74.9	5.6	19.5	129
20 HL	57.3	13.1	29.6	119	61.5	11.1	27.4	129

U - Unlimed; L - Limed; LL - Light lime; HL - Heavy lime; ND - None Dominant

A/6.24

Botanical Composition per cent. 1st Crop
1947

Plot	Gram- ineae	Legum- inosae	Other orders	Dom. Sp.	Plot	Gram- ineae	Legum- inosae	Other orders	Dom. Sp.
1 U	93.1	-	6.9	134	2 U	58.0	10.7	31.3	124
1 L	70.9	3.7	25.4	129	2 L	58.6	15.4	26.0	ND
3 U	53.3	11.0	35.7	124	4 ¹ U	46.8	13.5	39.7	134
3 L	39.5	15.2	45.3	110	4 ¹ L	42.0	21.2	36.8	124
7 U	51.2	17.7	31.1	119	4 ² U	97.9	-	2.1	ND
				120	4 ² L	77.4	0.1	22.5	134
7 L	71.0	9.1	19.9	115					
8 U	50.8	9.2	40.0	129	5 ¹	72.8	1.1	26.1	ND
				101	5 ²	49.1	11.0	39.9	119
8 L	60.2	10.3	29.5	129					
9 U	96.4	-	3.6	134	6	59.6	16.3	24.1	119
9 L	83.9	6.0	10.1	126					
10 U	98.8	-	1.2	134	11 ¹ U	87.8	0.2	12.0	112
10 L	89.4	-	10.6	134	11 ¹ L	98.0	-	2.0	126
13 U	76.6	0.7	22.7	129	11 ² U	98.8	-	1.2	134
13 L	66.9	7.5	25.6	129	11 ² L	98.4	0.1	1.5	126
14 U	85.1	3.3	11.6	113	12	53.2	8.5	38.3	124
14 L Sun	81.6	13.2	5.2	126					
14 L Shade	83.2	12.6	4.2	126					
18 U	93.4	-	6.6	134	15 U	61.9	14.1	24.0	129
18 LL	65.3	0.7	34.0	126	15 L	62.1	11.6	26.3	129
				127					
18 HL	64.7	1.6	33.7	126					
19 U	50.7	17.0	32.3	119	16 U	68.2	13.0	18.8	ND
19 LL	51.2	9.9	38.9	ND	16 L	72.3	7.5	20.2	129
19 HL	60.5	13.5	26.0	129					
20 U	68.6	8.5	22.9	134	17 U	69.7	0.1	30.2	129
20 LL	70.2	4.4	25.4	129	17 L	76.3	2.7	21.0	119
20 HL	52.4	10.3	37.3	127					

U - Unlimed; L - Limed; LL - Light lime; HL - Heavy lime; ND - None Dominant

Botanical Composition per cent. 1st Crop

Plot	1948			Plot	1949		
	Gram- ineae	Legum- nosae	Other orders		Gram- ineae	Legum- nosae	Other orders
1 U	94.7	-	5.3	2 U	53.6	15.4	31.0
1 L	63.3	4.6	32.1	2 L	51.4	17.8	30.8
3 U	53.0	7.2	39.8	4 ¹ U	46.7	14.4	38.9
3 L	35.8	16.1	48.1	4 ¹ L	44.3	22.4	33.3
7 U	46.6	19.8	33.6	4 ² U	99.8	-	0.2
7 L	56.5	25.3	18.2	4 ² L	94.7	-	5.3
8 U	52.5	7.3	40.2	5 ¹	82.2	3.0	14.8
8 L	53.4	11.1	35.5	5 ²	44.6	26.5	28.9
9 U	99.5	-	0.5	6	37.7	31.3	31.0
9 L	89.9	3.5	6.6				
10 U	99.7	-	0.3	11 ¹ U	99.7	-	0.3
10 L	93.8	-	6.2	11 ¹ L	98.2	-	1.8
13 U	73.8	0.5	25.7	11 ² U	99.4	-	0.6
13 L	67.0	10.6	22.4	11 ² L	97.5	-	2.5
14 U	92.4	2.0	5.6	12	58.5	13.4	28.1
14 L Sun	93.8	3.2	2.9	1			
14 L Shade	93.4	4.3	2.3				
				15 U	42.2	27.8	30.0
				15 L	46.6	32.7	20.7
18 U	88.5	-	11.5	16 U	75.3	11.9	12.8
18 LL	80.9	0.1	19.0	16 L	76.0	9.6	14.4
18 HL	82.9	0.6	16.5				
19 U	50.3	17.4	32.3	17 U	70.6	0.1	29.3
19 LL	62.1	8.5	29.4	17 L	83.2	2.8	14.0
19 HL	66.9	10.4	22.7				
20 U	86.3	5.0	8.7				
20 LL	80.3	4.9	14.8				
20 HL	74.2	9.9	15.9				

These results have not been previously published. Sampling was discontinued in 1950.

U - Unlimed; L - Limed; LL - Light lime; HL - Heavy lime; ND - None Dominant

WHEAT AND BARLEY - WOBURN STACKYARD

These are two almost identical, but distinct experiments which were for many years carried out on similar lines to those on Broadbalk and Hoosfield at Rothamsted. The field was under continuous wheat and barley from 1877 to 1926, and the treatments and results for this period are given in the 1928 Station Report p.103. From 1927-1940 no manures were applied, wheat and barley being grown each year except 1927, 28, 34, 35 when the field was fallowed. Results for 1929 to 1938 are given annually in the Station Reports in the sections devoted to Woburn. In 1941 and 42 a top dressing of 2 cwt. sulphate of ammonia was applied.

In 1943 a new scheme was started. The plots were divided into sets of three according to their previous manurial treatments (omitting plots 2, 5 and 8 of each crop, which were so acid as to give negligible crops). In 1943 the wheat was so weedy that it was ploughed up and the land fallowed for the rest of the season. In 1947 and 1948 the field was fallowed and no treatments were applied.

Area of one whole plot, $\frac{1}{4}$ acre.

The experiments are discussed by H. H. Mann, "The influence of fallowing on the yield of wheat or barley on very exhausted land", J. Agric. Sci., 33, (1943), 207.

Treatments

In the present system of manuring, of each set of three plots one receives a top-dressing of nitrochalk at 2 cwt. per acre (N1), one at 4 cwt. (N2), and the third at 6 cwt. (N3). The dressings rotate in cyclical order.

Summary of treatments 1877-1926 (plots arranged in the present sets of three)

Plot 1 Unmanured

3 Nitrate of Soda; 3a, (2N) since 1877; 3b, (1N) since 1907 only.

3aa, as 3a with lime in 1921; 3bb, as 3b with lime in 1921

7 Unmanured

4a Minerals; 4b, as 4a with lime in 1915

6 Minerals and Nitrate of Soda, (1N)

9a, 9b Minerals and, in alternate years, Nitrate of Soda (1N)

10a Superphosphate and Nitrate of Soda (1N)

10b Rape cake (1N)

11a Sulphate of Potash and Nitrate of Soda (1N)

11b Dung (4N)

- 2 Sulphate of Ammonia (1N); 2a unlimed, 2aa, limed in 1905, 09, 10, 11,
2b, limed in 1897 2bb limed in 1897 and 1905
- 5 Minerals and Sulphate of Ammonia (1N); 5a, unlimed 5aa, limed in
1905 and 1916 5b, limed in 1905
- 8 Minerals and, in alternate years, Sulphate of Ammonia (2N); 8a, 8b
unlimed, 8aa and 8bb limed in 1905 and 1918

N provides 25 lb. per acre of Ammonia

Crop Notes

Wheat-Red Standard			Barley - Plumage Archer		
Year	Sown	Harvested	Year	Sown	Harvested
1939	17.10.38	Aug. 15	1939	Feb. 27	Aug. 17
1940	2.11.39	Aug. 17	1940	Mar. 9	Aug. 8
1941	15.10.40	Aug. 19	1941	Mar. 18	Sept. 10
1942	19.11.41	Aug. 17	1942	Apr. 13	Sept. 4
1943		Ploughed in, May 6	1943	Mar. 12	Aug. 27
1944	27.9.43	Aug. 15	1944	Mar. 8	Sept. 4
1945	23.10.44	Aug. 2	1945	Mar. 6	Aug. 2
1946	2.10.45	Aug. 22	1946	Mar. 13	Aug. 15

Wheat and Barley - Woburn Stackyard

Yields of Wheat (lb. per acre)

Plot	Grain 1939	Straw	Grain 1940	Straw	Grain 1941	Straw	Grain 1942	Straw
1	450	976	294	609			213	680
2a	85	341	Nil	81	Nil	Nil	Nil	Nil
2aa	416	993	8	147	77	242	Nil	Nil
2b	263	847	98	214	113	255	Nil	Nil
2bb	337	674	278	459	528	1117	48	161
3a	289	632	447	721	935	1865	302	893
3b	155	386	239	506	641	1123	325	675
4	480	1124			975	2522	671	1396
5a	652	1362	201	423	258	610	79	417
5b	575	1030	217	602	1016	1815	171	364
6	435	1019	695	1362	1347	2246	797	2043
7	367	761	402	932	680	1909	334	901
8a	101	222	36	81	Nil	Nil	Nil	Nil
8aa	625	1117	355	649	69	177	67	419
8b	218	444	73	113	56	177	Nil	Nil
8bb	621	1194	218	452	278	661	173	706
9a	344	1052	544	1171	1115	2421	764	1929
9b	560	1194	474	919	919	1800	526	1339
10a	272	597	186	273			382	598
10b	289	554	231	446			371	888
11a	363	950	591	880	1197	2148	828	1516
11b	443	1312	918	1720	1211	2807	1168	2732

Yields of plots damaged by rabbits, or otherwise invalidated, are omitted.

Yields of Barley (lb. per acre)

Plot	Grain 1939	Straw	Grain 1940	Straw	Grain 1941	Straw	Grain 1942	Straw
1	385	1071	21	350	284	896	194	483
2a	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
2aa	367	1316	24	453	284	933	146	435
2b	341	877	24	325	442	1017	363	827
2bb	349	910	24	340	566	1200	151	384
3a	361	747	20	373	412	850	411	762
3aa	363	887	24	302	234	600	185	556
3b	593	1266	20	390	420	933	383	823
3bb	259	755	24	208	142	367	125	384
4a	337	976	12	234	691	1203	417	894
4b	228	929	12	226	304	1138	404	890
5a	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
5aa	175	774	Nil	Nil	Nil	Nil	323	762
5b	417	885	28	655	606	1057	560	947
6	717	1342	12	250	924	1484	1086	1384
7	368	1028	19	372	388	721	246	586
8a	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
8aa	435	1065	20	468	Taken with 8bb	658	1313	
8b	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
8bb	577	1113	20	500	686	1242	483	1100
9a	727	1412	37	677	1006	1675	975	1600
9b	572	1417	79	744	862	1500	1281	1583
10a	345	932	17	280	297	744	204	504
10b	288	1212	17	280	192	686	136	620
11a	444	967	76	543	676	1496	915	1393
11b	668	1635	135	912	886	876	1405	1909

Wheat and Barley - Woburn Stackyard

X New system of Manuring - Yields, (lb. per acre).
Plots arranged in their groups of three.

Wheat

Plot	Grain	1944	Straw	Grain	1945	Straw	Grain	1946	Straw
1	N2	1980	4887	N1	737	1334	N3		
3	N1	2259	5411	N3	1466	2559	N2	469	1927
7	N3	2494	5794	N2	841	2140	N1		
4	N3	2104	4371	N2	664	1558	N1		
6	N1	2318	5539	N3	1377	2689	N2	379	1862
9	N2	3625	6782	N1	601	1229	N3	744	2205
10a	N2	2250	5645	N1	587	1151	N3	569	2446
10b	N3	2109	4960	N2	833	1635	N1		
11a	N1	2504	7917	N3	1693	3183	N2	425	2014
11b(1)	N3	1872	4829	N2	1451	2841	N1	372	1598
11b(2)	N1	2452	6079	N3	1591	3037	N2	585	2354
11b(3)	N2	1689	4421	N1	762	1524	N3	750	2384

Yield of plots damaged by rabbits, or otherwise invalidated, are omitted.

Barley

Plot	1943		1944		1945		1946		
	Grain	Straw	Grain	Straw	Grain	Straw	Grain	Straw	
1	N1	87	311	N3	906	1766	N2	539	1839
3	N2	328	863	N1	529	1037	N3	908	2015
7	N3	458	977	N2	549	1115	N1	533	1594
4	N3	519	1051	N2	1343	1936	N1	676	2053
6	N1	551	1144	N3	821	1535	N2	1019	2717
9	N2	928	1661	N1	988	2152	N3	1020	2144
10a	N3	107	422	N2	404	1122	N1	304	1393
10b	N1	99	1045	N3	635	1929	N2	519	2081
11a	N2	702	1764	N1	668	2227	N3	857	2143
11b(1)	N2	1237	2537	N1	896	2530	N3	1006	2959
11b(2)	N3	800	2075	N2	604	2067	N1	727	2265
11b(3)	N1	606	2475	N3	1192	2545	N2	1087	2628
							N2	757	2482
							N3	1253	3279
							N1	1178	2586

| verso |
|-------|-------|-------|-------|-------|-------|-------|-------|
| verso |
TSCF	921	CH	461	TR	17	703	1000
	922	CH	865	DE	17	704	1001
	923	CH	945	LE	17	705	1002
	924	CH	946	LE	17	706	1003
	925	CH	821	LE	17	707	1004
	926	CH	962	LE	17	708	1005
	927	CH	963	LE	17	709	1006
	928	CH	1051	LE	17	710	1007
	929	CH	1052	LE	17	711	1008
	930	CH	1053	LE	17	712	1009
	931	CH	1054	LE	17	713	1010
	932	CH	1055	LE	17	714	1011
	933	CH	1056	LE	17	715	1012
	934	CH	1057	LE	17	716	1013
	935	CH	1058	LE	17	717	1014
	936	CH	1059	LE	17	718	1015
	937	CH	1060	LE	17	719	1016
	938	CH	1061	LE	17	720	1017
	939	CH	1062	LE	17	721	1018
	940	CH	1063	LE	17	722	1019
	941	CH	1064	LE	17	723	1020
	942	CH	1065	LE	17	724	1021
	943	CH	1066	LE	17	725	1022
	944	CH	1067	LE	17	726	1023
	945	CH	1068	LE	17	727	1024
	946	CH	1069	LE	17	728	1025
	947	CH	1070	LE	17	729	1026
	948	CH	1071	LE	17	730	1027
	949	CH	1072	LE	17	731	1028
	950	CH	1073	LE	17	732	1029
	951	CH	1074	LE	17	733	1030
	952	CH	1075	LE	17	734	1031
	953	CH	1076	LE	17	735	1032
	954	CH	1077	LE	17	736	1033
	955	CH	1078	LE	17	737	1034
	956	CH	1079	LE	17	738	1035
	957	CH	1080	LE	17	739	1036
	958	CH	1081	LE	17	740	1037
	959	CH	1082	LE	17	741	1038
	960	CH	1083	LE	17	742	1039
	961	CH	1084	LE	17	743	1040
	962	CH	1085	LE	17	744	1041
	963	CH	1086	LE	17	745	1042
	964	CH	1087	LE	17	746	1043
	965	CH	1088	LE	17	747	1044
	966	CH	1089	LE	17	748	1045
	967	CH	1090	LE	17	749	1046
	968	CH	1091	LE	17	750	1047
	969	CH	1092	LE	17	751	1048
	970	CH	1093	LE	17	752	1049
	971	CH	1094	LE	17	753	1050
	972	CH	1095	LE	17	754	1051
	973	CH	1096	LE	17	755	1052
	974	CH	1097	LE	17	756	1053
	975	CH	1098	LE	17	757	1054
	976	CH	1099	LE	17	758	1055
	977	CH	1100	LE	17	759	1056
	978	CH	1101	LE	17	760	1057
	979	CH	1102	LE	17	761	1058
	980	CH	1103	LE	17	762	1059
	981	CH	1104	LE	17	763	1060
	982	CH	1105	LE	17	764	1061
	983	CH	1106	LE	17	765	1062
	984	CH	1107	LE	17	766	1063
	985	CH	1108	LE	17	767	1064
	986	CH	1109	LE	17	768	1065
	987	CH	1110	LE	17	769	1066
	988	CH	1111	LE	17	770	1067
	989	CH	1112	LE	17	771	1068
	990	CH	1113	LE	17	772	1069
	991	CH	1114	LE	17	773	1070
	992	CH	1115	LE	17	774	1071
	993	CH	1116	LE	17	775	1072
	994	CH	1117	LE	17	776	1073
	995	CH	1118	LE	17	777	1074
	996	CH	1119	LE	17	778	1075
	997	CH	1120	LE	17	779	1076
	998	CH	1121	LE	17	780	1077
	999	CH	1122	LE	17	781	1078
	1000	CH	1123	LE	17	782	1079
	1001	CH	1124	LE	17	783	1080
	1002	CH	1125	LE	17	784	1081
	1003	CH	1126	LE	17	785	1082
	1004	CH	1127	LE	17	786	1083
	1005	CH	1128	LE	17	787	1084
	1006	CH	1129	LE	17	788	1085
	1007	CH	1130	LE	17	789	1086
	1008	CH	1131	LE	17	790	1087
	1009	CH	1132	LE	17	791	1088
	1010	CH	1133	LE	17	792	1089
	1011	CH	1134	LE	17	793	1090
	1012	CH	1135	LE	17	794	1091
	1013	CH	1136	LE	17	795	1092
	1014	CH	1137	LE	17	796	1093
	1015	CH	1138	LE	17	797	1094
	1016	CH	1139	LE	17	798	1095
	1017	CH	1140	LE	17	799	1096
	1018	CH	1141	LE	17	800	1097
	1019	CH	1142	LE	17	801	1098
	1020	CH	1143	LE	17	802	1099
	1021	CH	1144	LE	17	803	1100
	1022	CH	1145	LE	17	804	1101
	1023	CH	1146	LE	17	805	1102
	1024	CH	1147	LE	17	806	1103
	1025	CH	1148	LE	17	807	1104
	1026	CH	1149	LE	17	808	1105
	1027	CH	1150	LE	17	809	1106
	1028	CH	1151	LE	17	810	1107
	1029	CH	1152	LE	17	811	1108
	1030	CH	1153	LE	17	812	1109
	1031	CH	1154	LE	17	813	1110
	1032	CH	1155	LE	17	814	1111
	1033	CH	1156	LE	17	815	1112
	1034	CH	1157	LE	17	816	1113
	1035	CH	1158	LE	17	817	1114
	1036	CH	1159	LE	17	818	1115
	1037	CH	1160	LE	17	819	1116
	1038	CH	1161	LE	17	820	1117
	1039	CH	1162	LE	17	821	1118
	1040	CH	1163	LE	17	822	1119
	1041	CH	1164	LE	17	823	1120
	1042	CH	1165	LE	17	824	1121
	1043	CH	1166	LE	17	825	1122
	1044	CH	1167	LE	17	826	1123
	1045	CH	1168	LE	17	827	1124
	1046	CH	1169	LE	17	828	1125
	1047	CH	1170	LE	17	829	1126
	1048	CH	1171	LE	17	830	1127
	1049	CH	1172	LE	17	831	1128
	1050	CH	1173	LE	17	832	1129
	1051	CH	1174	LE	17	833	1130
	1052	CH	1175	LE	17	834	1131
	1053	CH	1176	LE	17	835	1132
	1054	CH	1177	LE	17	836	1133
	1055	CH	1178	LE	17	837	1134
	1056	CH	1179	LE	17	838	1135
	1057	CH	1180	LE	17	839	1136
	1058	CH	1181	LE	17	840	1137
	1059	CH	1182	LE	17	841	1138
	1060	CH	1183	LE	17	842	1139
	1061	CH	1184	LE	17	843	1140
	1062	CH	1185	LE	17	844	1141
	1063	CH	1186	LE	17	845	1142
	1064	CH	1187	LE	17	846	1143
	1065	CH	1188	LE	17	847	1144
	1066	CH	1189	LE	17	848	1145
	1067	CH	1190	LE	17	849	1146
	1068	CH	1191	LE	17	850	1147
	1069	CH	1192	LE	17	851	1148
	1070	CH	1193	LE	17	852	1149
	1071	CH	1194	LE	17	853	1150
	1072	CH	1195	LE	17	854	1151
	1073	CH	1196	LE	17	855	1152
	1074	CH	1197	LE	17	856	1153
	1075	CH	1198	LE	17	857	1154
	1076	CH	1199	LE	17	858	1155
	1077	CH	1200	LE	17	859	1156
	1078	CH	1201	LE	17	860	1157
	1079	CH	1202	LE	17	861	1158
	1080	CH	1203	LE	17	862	1159
	1081	CH	1204	LE	17	863	1160
	1082	CH	1205	LE	17	864	1161
	1083	CH	1206	LE	17	865	1162
	1084	CH	1207	LE	17	866	1163
	1085	CH	1208	LE	17	867	1164
	1086	CH	1209	LE	17	868	1165
	1087	CH	1210	LE	17	869	1166
	1088	CH	1211	LE	17	870	1167
	1089	CH	1212	LE	17	871	1168
	1090	CH	1213	LE	17	872	1169
	1091	CH	1214	LE	17	873	1170
	1092	CH	1215	LE	17	874	1171
	1093	CH	1216	LE	17	875	1172
	1094	CH	1217	LE	17	876	1173
	1095	CH	1218	LE	17	877	1174
	1096	CH	1219	LE	17	878	1175
	1097	CH	1220	LE	17	879	1176
	1098	CH	1221	LE	17	880	1177
	1099	CH	1222	LE	17	881	1178
	1100	CH	1223	LE	17	882	1179
	1101	CH	1224	LE	17	883	1180
	1102	CH	1225	LE	17	884	1181
	1103	CH	1226	LE	17	885	1182
	1104	CH	1227	LE	17	886	1183
	1105	CH	1228	LE	17	887	1184
	1106	CH	1229	LE	17	888	1185

TWO COURSE ROTATION EXPERIMENT

Long Hoos (begun in 1942)Cumulative Effects of Agricultural Salt

The experiment was designed to test the cumulative effects of agricultural salt and muriate of potash, and to compare two methods of application of the salt.

The rotation is 1st year Sugar beet, 2nd year Barley.

There are 2 series (one for each crop in any particular year), each consisting of 4 blocks of 12 plots each. Area of each plot in acres:

Series	Total	Sugar Beet	Barley
		Harvested	
1	0.0219	0.0190	0.0207
2	0.0200	0.0173	0.0189

Treatments: all combinations of

- (1) Agricultural salt (rates for sugar beet): None (0), $2\frac{1}{2}$ cwt. (Na 1), 5 cwt. (Na 2) and $7\frac{1}{2}$ cwt. (Na 3) per acre.
- (2) Muriate of potash: None (0), the equivalent of half the single dressing of salt (K1) (approximately 1 cwt. K₂O per acre), the equivalent of the double dressing of salt (K2), applied to sugar beet at sowing.
- (3) Time of application of salt: in seed bed before sowing (A), before ploughing during the winter (B).
- (4) Salt applied to sugar beet only (0), salt repeated at half rate on barley (R).

The following interactions are partially confounded with block differences:-

(A-B) x R; (3-2+1-0) Na x (A-B); (3-2+1-0) Na x R;

and the interactions of these with K.

Basal dressings, applied to all plots at sowing:

Barley:	0.3 cwt. N per acre as sulphate of ammonia
Sugar beet:	0.8 cwt. N per acre as sulphate of ammonia
	0.6 cwt. P ₂ O ₅ per acre as superphosphate.

Crop Notes

	Sugar beet:	Klein E Sown	Lifted	Barley:	Plumage Archer Sown	Harvested
1942	April 25	Dec. 2		(Series 2 grew preliminary wheat)		
1943	April 14	Nov. 12		March 13		Aug. 10
1944	April 22	Oct. 26, Nov. 23		March 8		Aug. 10
1945	April 12	Nov. 5		March 27		Aug. 8
1946	April 15	Nov. 22		March 21		Aug. 14
1947	April 25	Oct. 30		April 11		Aug. 11

In 1942, 1943, samples were taken at intervals from one end of each plot for physiological and chemical study. The yields given are corrected for the reduced area per plot. In 1944, 12 of the 48 plots of sugar beet were lifted Oct. 26th. This had an appreciable effect on tops and sugar percentage, which have been adjusted by means of a regression. There was no effect on the other results.

Two Course Rotation Experiment

		Sugar Beet					
		Year					
Salt cwt.		1942	1943	1944	1945	1946	1947
Muriate of potash K ₂ O cwt. per acre		Roots (washed): tons per acre					
0.0	0	14.98	15.46	8.51	16.18	15.34	5.92
	2.5	16.95	17.60	10.96	17.96	17.71	9.04
	5.0 (a)	17.12	17.63	10.61	17.15	16.86	9.33
	7.5	16.72	16.68	9.69	16.86	16.32	9.23
	Mean (b)	16.44	16.84	9.94	17.04	16.56	8.38
1.0	0	14.73	16.97	8.04	16.97	15.19	7.48
	2.5	16.08	17.03	11.25	16.88	17.08	8.50
	5.0 (a)	16.86	17.12	10.53	17.34	17.61	9.99
	7.5	17.61	16.08	11.34	16.39	17.16	9.74
	Mean (b)	16.32	16.80	10.29	16.90	16.76	8.93
2.0	0	15.68	16.87	9.34	16.57	16.93	7.54
	2.5	16.28	16.45	10.96	17.12	17.56	8.72
	5.0 (a)	16.65	16.91	10.28	17.55	17.21	9.26
	7.5	17.28	16.87	11.24	16.21	17.02	10.20
	Mean (b)	16.47	16.78	10.46	16.86	17.18	8.93
Salt applied Winter		2.5	16.04	17.43	10.71	17.27	17.66
		5.0 (d)	16.58	17.32	10.09	17.21	17.35
		7.5	17.10	16.23	11.10	16.63	16.24
		Mean (e)	16.57	16.99	10.63	17.04	17.08
		2.5	16.83	16.62	11.40	17.37	17.24
		5.0 (d)	17.17	17.12	10.86	17.48	17.11
In seed bed		7.5	17.31	16.86	10.41	16.35	17.43
		Mean (e)	17.10	16.87	10.89	17.07	17.26
Mean		0	15.13	16.43	8.63	16.57	15.82
		2.5	16.44	17.03	11.06	17.32	17.45
		5.0 (c)	16.88	17.22	10.47	17.35	17.23
		7.5	17.20	16.54	10.76	16.49	16.83
		Mean	16.41	16.81	10.23	16.93	16.83
Standard errors (28 d.f. in 1942, 22 d.f. in other years)		(a)	0.517	0.577	0.503	0.384	0.427
		(b)	0.259	0.289	0.251	0.192	0.213
		(c)	0.299	0.333	0.290	0.222	0.247
		(d)	0.395	0.441	0.384	0.294	0.326
		(e)	0.239	0.266	0.232	0.177	0.197
		Per plot	1.035	1.154	1.005	0.769	0.854
		%age	6.3	6.9	9.8	4.5	5.1

Sugar Beet

	Salt cwt. per acre	Year					
		1942	1943	1944	1945	1946	1947
Sugar Percentage							
Muriate of potash							
K ₂ O cwt. per acre	0	18.19	18.44	16.24	18.03	17.04	19.34
0.0	2.5	18.02	18.84	16.93	18.47	17.04	20.42
	5.0 (a)	18.54	18.85	17.02	18.81	17.14	21.07
	7.5	18.31	18.81	16.60	18.57	17.18	20.40
	Mean (b)	18.27	18.74	16.70	18.47	17.10	20.31
	0	18.19	18.48	16.62	18.49	16.96	20.86
	2.5	18.30	18.97	16.84	18.46	16.79	20.48
1.0	5.0 (a)	18.16	18.72	16.76	18.38	17.29	20.58
	7.5	17.98	18.60	16.66	18.64	17.16	20.54
	Mean (b)	18.16	18.69	16.72	18.49	17.05	20.62
	0	18.26	18.78	16.68	18.57	17.49	20.48
	2.5	18.53	18.83	16.92	18.83	17.38	20.76
2.0	5.0 (a)	18.38	18.84	16.78	18.58	17.21	20.43
	7.5	18.26	18.60	16.89	18.70	16.98	21.03
	Mean (b)	18.35	18.76	16.82	18.67	17.26	20.68
Salt applied	2.5	18.30	19.00	16.85	18.64	17.08	20.33
Winter	5.0 (d)	18.34	18.85	16.72	18.53	17.10	20.76
	7.5	18.08	18.61	16.83	18.63	17.10	20.94
	Mean (e)	18.24	18.82	16.80	18.60	17.09	20.68
In seed bed	2.5	18.27	18.76	16.94	18.53	17.05	20.78
	5.0 (d)	18.37	18.76	16.98	18.65	17.33	20.63
	7.5	18.28	18.73	16.60	18.65	17.12	20.38
	Mean (e)	18.31	18.75	16.84	18.61	17.17	20.60
Mean	0	18.21	18.57	16.51	18.36	17.16	20.23
	2.5	18.28	18.88	16.90	18.59	17.07	20.55
	5.0 (c)	18.36	18.80	16.85	18.59	17.21	20.69
	7.5	18.18	18.67	16.72	18.64	17.11	20.66
	Mean	18.26	18.73	16.74	18.54	17.14	20.54
Standard errors (28 d.f. in 1942, 21 d.f. in 1944, 22 d.f. in other years)	(a)	0.180		0.220	0.165	0.262	
	(b)	0.090		0.110	0.082	0.131	
	(c)	0.104		0.127	0.095	0.151	
	(d)	0.137		0.168	0.126	0.200	
	(e)	0.083		0.102	0.076	0.121	
	Per plot	0.360		0.441	0.329	0.524	

Two Course Rotation ExperimentSugar Beet

	Salt cwt. per acre	Year					
		1942	1943	1944	1945	1946	1947
Muriate of potash K_2O cwt. per acre		Total sugar:					
0.0	0	54.5	57.0	27.2	58.3	52.3	23.7
	2.5	61.1	66.3	36.6	66.3	60.2	37.2
	5.0 (a)	63.5	66.5	36.1	64.5	57.9	39.5
	7.5	61.2	62.7	32.2	62.7	56.0	37.9
	Mean (b)	60.1	63.1	33.0	63.0	56.6	34.6
	0	53.6	62.7	27.1	62.8	51.5	31.3
1.0	2.5	58.8	64.6	37.4	62.3	57.5	35.1
	5.0 (a)	61.2	64.1	35.8	63.8	60.9	41.1
	7.5	63.3	59.8	37.7	61.1	58.8	40.4
	Mean (b)	59.2	62.8	34.5	62.5	57.2	37.0
	0	57.3	63.3	31.6	61.5	59.2	31.1
	2.5	60.4	62.0	37.1	64.4	61.0	36.5
2.0	5.0 (a)	61.1	63.9	34.5	65.2	59.2	38.4
	7.5	63.1	62.8	37.9	60.6	57.8	42.9
	Mean (b)	60.5	63.0	35.3	62.9	59.3	37.2
	0	58.6	66.2	35.8	64.3	60.3	35.4
	2.5	60.8	65.3	34.1	63.9	59.4	40.0
	5.0 (a)	61.8	60.3	37.3	61.7	55.5	42.5
In seed bed	7.5	60.4	63.9	35.7	63.3	58.4	39.3
	2.5	61.6	62.4	38.3	64.4	58.9	37.2
	5.0 (a)	63.1	64.3	36.9	65.1	59.3	39.4
	7.5	63.3	63.2	34.6	61.3	59.7	38.3
	Mean (e)	62.6	63.3	36.6	63.6	59.3	38.3
	0	55.1	61.0	28.6	60.9	54.3	28.7
Mean	2.5	60.1	64.3	37.0	64.3	59.6	36.3
	5.0 (c)	61.9	64.8	35.5	64.5	59.3	39.7
	7.5	62.5	61.8	35.9	61.5	57.5	40.4
	Mean	59.9	63.0	34.3	62.8	57.7	36.3
	(a)	1.85	2.17	1.74	1.52	1.91	2.50
	(b)	0.92	1.08	0.87	0.76	0.96	1.25
Standard errors (28 d.f. in 1942, 22 d.f. in other years)	(c)	1.07	1.25	1.00	0.87	1.10	1.44
	(d)	1.41	1.66	1.32	1.16	1.46	1.91
	(e)	0.85	1.00	0.80	0.71	0.88	1.15
	Per plot	3.69	4.34	3.47	3.03	3.82	5.00
	%age	6.2	6.9	10.1	4.8	6.6	13.8

Sugar Beet

	Salt cwt. per acre	1942	1943	1944	1945	1946	1947
Tops: tons per acre							
Muriate of potash	0	10.06	9.55	10.69	12.20	17.05	4.59
K ₂ O cwt. per acre	2.5	11.23	10.76	13.20	11.82	18.01	6.94
0.0	5.0 (a)	10.56	11.55	13.01	11.80	17.71	7.23
	7.5	11.42	11.60	12.52	12.74	17.21	7.01
	Mean (b)	10.82	10.86	12.36	12.14	17.50	6.44
1.0	0	10.41	10.62	12.40	11.89	17.73	5.81
	2.5	12.19	11.10	13.80	13.56	17.73	6.40
	5.0 (a)	10.20	10.50	11.98	12.43	16.65	7.66
	7.5	13.30	10.74	14.27	12.20	18.18	7.26
	Mean (b)	11.53	10.74	13.11	12.52	17.57	6.76
2.0	0	10.43	10.24	11.70	11.76	17.31	5.83
	2.5	11.74	10.17	12.06	12.05	16.93	6.59
	5.0 (a)	11.42	10.34	11.49	11.49	17.07	7.30
	7.5	11.77	11.12	12.63	11.45	16.49	7.48
	Mean (b)	11.34	10.47	11.97	11.69	16.95	6.80
Salt applied Winter	2.5	11.55	10.49	12.69	12.08	17.83	6.63
	5.0 (a)	10.58	10.80	12.47	11.62	17.54	7.52
	7.5	12.17	11.06	13.89	11.91	17.48	7.62
	Mean (e)	11.43	10.78	13.02	11.87	17.62	7.26
In seed bed	2.5	11.88	10.87	13.34	12.89	17.29	6.65
	5.0 (a)	10.87	10.79	11.86	12.19	16.75	7.27
	7.5	12.17	11.25	12.39	12.35	17.10	6.88
	Mean (e)	11.64	10.97	12.53	12.48	17.05	6.93
Mean	0	10.30	10.14	11.59	11.95	17.36	5.41
	2.5 (c)	11.72	10.68	13.02	12.48	17.56	6.64
	5.0	10.73	10.80	12.16	11.91	17.14	7.40
	7.5	12.17	11.15	13.14	12.13	17.29	7.25
	Mean	11.23	10.69	12.48	12.12	17.34	6.68
Standard errors (28 d.f. in 1942, 21 d.f. in 1944, 22 d.f. in other years)	(a)	0.669	0.600	0.680	0.382	0.813	0.353
	(b)	0.334	0.300	0.340	0.191	0.407	0.177
	(c)	0.386	0.346	0.392	0.221	0.470	0.204
	(d)	0.511	0.458	0.519	0.292	0.621	0.270
	(e)	0.309	0.277	0.314	0.176	0.375	0.163
	Per plot %age	1.338	1.200	1.359	0.764	1.627	0.707
		11.9	11.2	10.9	6.3	9.4	10.6



	Salt cwt. per acre	Sugar Beet					
		1942	1943	1944	1945	1946*	1947
Muriate of potash		Plant Number: thousands per acre					
K ₂ O cwt. per acre	0	25.9	23.5	27.7	27.6		16.3
0.0	2.5	25.3	25.6	29.1	28.5		19.5
	5.0 (a)	26.0	25.3	27.8	28.4		19.1
	7.5	26.3	24.5	24.6	28.1		19.1
	Mean (b)	25.9	24.7	27.3	28.2		18.5
	0	25.2	23.9	27.9	28.3		19.2
1.0	2.5	23.4	24.5	27.7	27.7		19.7
	5.0 (a)	27.4	24.1	29.2	28.6		21.2
	7.5	27.3	22.6	26.7	27.9		20.6
	Mean (b)	25.8	23.8	27.9	28.1		20.2
	0	24.8	25.1	27.8	28.2		18.1
2.0	2.5	24.5	22.7	27.7	27.9		18.8
	5.0 (a)	27.1	24.0	28.5	27.6		19.7
	7.5	26.5	23.5	27.7	27.0		22.1
	Mean (b)	25.7	23.8	27.9	27.7		19.7
Salt applied Winter	2.5	24.8	24.1	28.3	27.8		18.8
	5.0 (d)	27.3	24.3	28.7	27.8		19.6
	7.5	26.5	23.8	26.2	27.9		21.4
	Mean (e)	26.2	24.1	27.7	27.8		19.9
In seed bed	2.5	24.0	24.4	28.0	28.3		19.8
	5.0 (d)	26.3	24.6	28.4	28.6		20.4
	7.5	26.9	23.2	26.4	27.4		19.8
	Mean (e)	25.7	24.1	27.6	28.1		20.0
Mean	0	25.3	24.1	27.8	28.0		17.9
	2.5	24.4	24.3	28.2	28.0		19.3
	5.0 (c)	26.8	24.5	28.5	28.2		20.0
	7.5	26.7	23.5	26.3	27.7		20.6
	Mean	25.8	24.1	27.7	28.0		19.5
In seed bed	2.5	24.0	24.4	28.0	28.3		19.8
	5.0 (d)	26.3	24.6	28.4	28.6		20.4
	7.5	26.9	23.2	26.4	27.4		19.8
	Mean (e)	25.7	24.1	27.6	28.1		20.0
Standard errors (28 d.f. in 1942, 22 d.f. in other years)	(a)	0.93	1.00	0.94	0.56		
	(b)	0.46	0.50	0.47	0.28		
	(c)	0.54	0.58	0.55	0.32		
	(d)	0.71	0.76	0.72	0.43		
	(e)	0.43	0.46	0.44	0.26		
	Per plot	1.86	2.00	1.89	1.12		
	%age	7.2	8.3	6.8	4.0		

* The plant numbers were not counted in 1946.

Barley

	Salt applied in previous year cwt. per acre	Grain: cwt. per acre				
		1943	1944	1945	1946	1947
Muriate of potash applied in previous year K ₂ O cwt. per acre	0	24.0	30.0	27.1	32.6	23.1
	2.5	24.9	31.1	27.8	31.9	23.9
	5.0 (a)	23.8	30.2	27.5	33.0	22.1
	0.0	7.5	23.7	29.8	26.5	32.0
		Mean (b)	24.1	30.3	27.2	32.4
						22.7
		0	22.5	30.3	26.4	34.3
		2.5	26.9	29.6	29.4	32.7
	1.0	5.0 (a)	22.7	30.5	26.6	35.5
		7.5	26.6	30.1	29.5	32.4
		Mean (b)	24.7	30.1	28.0	33.7
Salt in current year	0	23.3	29.4	26.1	32.8	23.7
	2.5	25.7	30.7	28.0	33.0	24.1
	5.0 (a)	23.9	29.4	27.2	33.3	22.7
	0.0	7.5	26.4	30.7	29.3	31.9
		Mean (b)	24.8	30.0	27.6	32.8
						23.5
	2.5	24.4	30.8	27.6	33.2	23.4
	5.0 (c)	22.7	30.0	26.4	33.4	22.7
	None	7.5	26.1	29.8	28.7	31.3
		Mean (d)	24.4	30.2	27.6	32.6
Half Rate	2.5	27.3	30.1	29.1	31.9	26.2
	5.0 (c)	24.2	30.1	27.8	34.5	22.0
	0.0	7.5	25.1	30.5	28.2	32.9
		Mean (d)	25.5	30.2	28.4	33.1
						23.5
	2.5	25.1	30.0	27.9	32.7	25.4
	5.0 (c)	23.2	30.0	26.8	34.3	22.4
	0.0	7.5	27.2	30.5	29.5	32.4
		Mean (d)	25.2	30.2	28.1	33.1
						23.2
In seed bed	2.5	26.6	31.0	28.9	32.4	24.2
	5.0 (c)	23.8	30.1	27.4	33.5	22.3
	0.0	7.5	23.9	29.8	27.4	31.8
		Mean (d)	24.8	30.3	27.9	32.6
						23.8
	Mean	0	23.3	29.9	26.5	33.2
		2.5	25.8	30.5	28.4	32.5
		5.0 (e)	23.5	30.0	27.1	33.9
		7.5	25.6	30.2	28.4	32.1
		Mean	24.5	30.1	27.6	32.9
Standard errors (22 d.f.)	(a)	0.92	0.68	0.66	1.43	0.82
	(b)	0.46	0.34	0.33	0.72	0.41
	(c)	0.71	0.52	0.50	1.09	0.63
	(d)	0.43	0.32	0.31	0.66	0.38
	(e)	0.53	0.40	0.38	0.83	0.47
Per plot	1.85	1.37	1.31	2.86	1.64	
	%age	7.5	4.6	4.8	8.7	7.0

Barley

Salt applied in previous year cwt. per acre	Year						
	1943	1944	1945	1946	1947		
Muricate of potash applied in previous year K ₂ O cwt. per acre	0	27.9	28.9	31.7	36.5	21.2	
	2.5	25.7	28.9	33.0	36.6	22.0	
	5.0 (a)	25.8	27.3	32.7	36.3	21.1	
	7.5	24.6	28.4	30.2	37.5	20.9	
	Mean (b)	26.0	28.4	31.9	36.7	21.3	
	0	23.9	28.9	32.2	36.2	22.2	
	2.5	29.4	28.4	34.3	38.3	24.4	
	5.0 (a)	24.3	28.4	32.2	39.0	21.8	
	7.5	27.6	29.2	34.8	37.3	24.3	
	Mean (b)	26.3	28.7	33.4	37.7	23.2	
1.0	0	24.8	27.4	31.0	38.0	22.1	
	2.5	27.1	29.0	32.9	37.8	22.7	
	5.0 (a)	25.4	27.6	32.8	37.7	21.5	
	7.5	27.8	27.6	34.5	38.6	23.0	
	Mean (b)	26.3	27.9	32.8	38.0	22.3	
Salt in current year	2.5	25.7	29.1	32.6	37.9	22.2	
	None	5.0 (c)	24.5	28.3	32.4	37.9	22.0
	7.5	27.2	28.3	34.1	37.7	23.8	
	Mean (d)	25.8	28.6	33.0	37.8	22.7	
	2.5	29.1	28.5	34.1	37.3	23.8	
	5.0 (c)	25.8	27.3	32.7	37.4	21.0	
	7.5	26.2	28.5	32.2	37.8	21.6	
	Mean (d)	27.0	28.1	33.0	37.5	22.1	
	2.5	26.9	28.4	33.1	37.2	23.0	
	5.0 (c)	24.8	28.2	32.5	38.6	22.1	
Half rate	7.5	27.8	29.0	34.4	38.5	20.5	
	Mean (d)	26.5	28.5	33.4	38.1	21.9	
	2.5	27.9	29.2	33.7	37.9	23.1	
	5.0 (c)	25.5	27.4	32.6	36.7	20.8	
	7.5	25.6	27.8	31.9	37.1	24.9	
Salt applied Winter Winter	Mean (d)	26.4	28.1	32.7	37.2	22.9	
	0	25.5	28.4	31.6	36.9	21.8	
	2.5	27.4	28.8	33.4	37.6	23.0	
	5.0 (e)	25.2	27.8	32.6	37.7	21.5	
	7.5	26.7	28.4	33.2	37.8	22.7	
	Mean	26.2	28.4	32.7	37.5	22.3	
	Standard errors (22 d.f.)	(a)		0.80	1.25		
		(b)		0.39	0.62		
		(c)		0.61	0.95		
		(d)		0.37	0.58		
Mean		(e)		0.46	0.72		
	Per plot			1.61	2.50		
	%age			4.9	6.7		

THREE COURSE ROTATION EXPERIMENT

Long Hoos (begun in 1933)

The effects of straw compost, of straw ploughed in with fertilizers, and of magnesium sulphate

There are three series each of 24 plots, and the three crops of the rotation, potatoes, barley and sugar beet rotate on each series. The plots receive the same dressings every other year so that the direct and residual effects of manures can be measured. Area of each plot, 0.02 acre.

Treatments and rates of dressing (per acre)

	Straw tons	Compost tons	Sulphate of Ammonia cwt.	Super- phosphate of potash cwt.	Muriate of potash cwt.
No organic manure (Art.)	-	-	2.0	2.2	0.8
Compost in Autumn (Adco)	-	5.0	-	-	0.8
Straw, ploughed in in autumn, artificials in spring (St ₁)	2.5	-	2.0	2.2	0.8
Straw, ploughed in in autumn, artificials half in autumn, half in spring (St ₂)	2.5	-	2.0 ¹	2.2	0.8 ²

1. For sugar beet the half-dressing applied in spring is 1.3 cwt. Nitrate of Soda;
2. Until 1946 the spring half-dressing to potatoes was 0.4 cwt. Sulphate of potash.

These four treatments are applied to three plots of each crop each year.

Magnesium Sulphate: 2.5 cwt. per acre applied to one of three replicate plots of each treatment in each crop section.

In addition to the above dressings, the potatoes receive a basal dressing of 2 cwt. sulphate of ammonia, 2.2 cwt. superphosphate and 0.8 cwt. muriate of potash. The sugar beet receive 1.3 cwt. nitrate of soda, 1.1 cwt. superphosphate and 0.4 cwt. muriate of potash.

Details of the experiment are as given in the 1933 Station Report, pp. 118-119 except that no comparisons of winter green manuring crops are now made and that commencing in 1943, a yearly dressing of megnesium sulphate has been applied as detailed above. Results for 18 years are discussed in the Station Report for 1951, p. 135.

Crop Notes

Potatoes (after Sugar beet). Barley (after Potatoes). Sugar beet (after Barley)

	Variety	Planted	Lifted	Variety	Sown	Harvested	Variety	Sown	Lifted
1939	Ally	Apr. 17	Sept. 18	Plumage	Mar. 7	Aug. 18	Kuhn	Apr. 24	Oct. 25
				Archer	"				
1940	"	Apr. 26	Sept. 24	"	Mar. 9	Aug. 5	"	May 3	Oct. 17
1941	"	Apr. 28	Oct. 3	"	Mar. 25	Sept. 2	"	Apr. 30	Oct. 25
1942	Majestic	"	21 Oct. 3	"	Mar. 25	Aug. 17	Klein E	Apr. 24	Oct. 26
1943	"	Apr. 8	Sept. 17	"	Mar. 3	Aug. 7	"	Apr. 14	Oct. 14
1944	"	Mar. 28	Sept. 28	"	Mar. 7	Aug. 12	"	Apr. 22	Oct. 24
1945	"	Mar. 30	Sept. 24	"	Mar. 26	Aug. 11	"	Apr. 12	Oct. 19
1946	"	Apr. 1	Oct. 3	"	Mar. 21	Aug. 21	"	Apr. 15	Oct. 22
1947	"	Apr. 30	Sept. 27	"	Apr. 11	Aug. 11	"	Apr. 26	Oct. 17

Three-course Rotation Experiment

Ba/2.3

Potatoes. Total tubers: tons per acre

	Treatments applied same year					Treatments applied previous year				
	Art	Adco	St ₁	St ₂	Mean	Art	Adco	St ₁	St ₂	Mean
Mean Yields										
1939	11.02	10.14 ±0.439	12.41	11.03	11.15 ±0.219	8.87	8.55 ±0.439	10.32	10.51	9.56 ±0.219
1940	8.05	7.32 ±0.183	8.37	8.23	7.99 ±0.092	6.14	6.91 ±0.183	6.96	6.92	6.73 ±0.092
1941	11.08	9.16 ±0.652	12.33	11.16	10.93 ±0.326	8.68	10.31 ±0.652	9.48	10.50	9.74 ±0.326
1942	9.66	8.48 ±0.324	10.21	9.71	9.52 ±0.162	6.79	7.20 ±0.324	8.80	8.52	7.83 ±0.162
1943	8.20	7.05 ±0.188	8.53	8.93	8.18 ±0.094	5.56	5.79 ±0.188	6.93	6.80	6.27 ±0.094
1944	10.10	9.92 ±0.502	10.74	10.55	10.33 ±0.251	8.68	9.15 ±0.502	9.62	9.81	9.32 ±0.251
1945	11.21	10.05 ±0.467	11.67	11.69	11.16 ±0.234	9.14	8.91 ±0.467	10.83	10.24	9.78 ±0.234
1946	10.88	10.10 ±0.367	12.43	12.27	11.42 ±0.183	8.45	8.65 ±0.367	9.95	9.64	9.17 ±0.183
1947	7.56	6.07 ±0.562	7.33	7.07	7.01 ±0.281	5.51	6.45 ±0.562	5.78	6.29	6.01 ±0.281

Responses to Magnesium Sulphate

1943	-0.04 ±0.399	0.42	-1.31	0.17	-0.19 ±0.200	0.22	-1.03 ±0.399	-1.09	-0.34	-0.56 ±0.200
1944	1.70 ±1.06	-0.62 ±0.532	-0.22	0.68	0.38 ±0.532	-0.66	-1.28 ±1.06	1.20	1.00	0.06 ±0.532
1945	0.31 ±0.992	1.33 ±0.496	-0.06	0.51	0.52 ±0.496	-1.32	1.21 ±0.992	-0.38	0.13	0.09 ±0.496
1946	-0.65 ±0.778	-0.98 ±0.389	2.18	-0.19	0.09 ±0.389	0.11	0.44 ±0.778	-0.55	-0.22	-0.06 ±0.389
1947	0.20 ±1.19	0.34 ±0.596	0.38	-0.18	0.19 ±0.596	0.04	-0.23 ±1.19	-0.62	0.20	-0.15 ±0.596

Standard errors per plot

	d.f.
1939 0.760 tons per acre or 7.3%	16
1940 0.318 tons per acre or 4.3%	16
1941 1.13 tons per acre or 10.9%	16
1942 0.561 tons per acre or 6.5%	16
1943 0.326 tons per acre or 4.5%	8
1944 0.869 tons per acre or 8.8%	8
1945 0.810 tons per acre or 7.7%	8
1946 0.635 tons per acre or 6.2%	8
1947 0.974 tons per acre or 15.0%	8

σ^2 between plots
 σ^2_e plots + years

Barley Grain: cwt. per acre

	Treatments applied same year					Treatments applied previous year				
	Art	Adco	St ₁	St ₂	Mean	Art	Adco	St ₁	St ₂	Mean
Mean Yields										
1939	39.9	30.8 ±2.00	35.4	32.2	34.6 ±1.00	33.9	29.1 ±2.00	29.2	32.7	31.2 ±1.00
1940	29.0	29.4 ±1.28	28.5	29.5	29.1 ±0.640	28.3	24.6 ±1.28	26.5	29.0	27.1 ±0.640
1941	28.9	21.8 ±0.771	27.9	25.1	25.9 ±0.385	22.2	19.7 ±0.771	24.2	23.0	22.3 ±0.385
1942	32.1	26.0 ±1.82	27.4	28.6	28.5 ±0.909	26.9	26.2 ±1.92	27.3	24.3	26.2 ±0.909
1943	32.3	25.8 ±0.867	30.5	29.8	29.6 ±0.434	25.2	24.3 ±0.867	24.8	26.1	25.1 ±0.434
1944	35.5	31.4 ±1.54	33.0	34.7	33.6 ±0.769	31.3	32.2 ±1.54	29.9	31.0	31.1 ±0.769
1945	33.2	27.3 ±2.07	32.1	32.8	31.4 ±1.03	25.0	25.3 ±2.07	27.6	28.7	26.6 ±1.03
1946	34.6	33.5 ±1.54	29.7	32.2	32.5 ±0.768	29.2	28.8 ±1.54	27.9	29.9	29.0 ±0.768
1947	27.4	20.6 ±1.09	26.7	23.7	24.6 ±0.546	19.7	16.9 ±1.09	18.6	20.0	18.8 ±0.546
Responses to Magnesium Sulphate										
1943	1.1	4.8 ±1.84	0.1	0.2	1.6 ±0.920	-4.3	0.2 ±1.84	1.9	-2.5	-1.2 ±0.920
1944	0.6	1.2 ±3.26	-0.6	1.9	0.8 ±1.63	-3.3	-2.9 ±3.26	0.8	2.2	-0.8 ±1.63
1945	-2.2	-1.5 ±4.39	2.2	-2.2	-0.9 ±2.19	-0.5	-7.9 ±4.39	-0.8	-0.9	-2.5 ±2.19
1946	0.9	-0.4 ±3.26	1.2	-2.1	0.3 ±1.63	2.3	4.9 ±3.26	-1.2	-2.1	1.0 ±1.63
1947	-4.4	-2.3 ±2.31	-0.1	1.7	-1.3 ±1.16	2.4	1.4 ±2.31	-5.9	1.1	-0.3 ±1.16

Standard errors per plot

d.f.

1939	3.47 cwt. per acre or 10.6%	16
1940	2.22 cwt. per acre or 7.9%	15*
1941	1.34 cwt. per acre or 5.5%	16
1942	3.15 cwt. per acre or 11.5%	16
1943	1.50 cwt. per acre or 5.5%	8
1944	2.66 cwt. per acre or 8.2%	8
1945	3.58 cwt. per acre or 12.4%	8
1946	2.66 cwt. per acre or 8.7%	8
1947	1.89 cwt. per acre or 8.7%	8

*One missing plot.

Three Course Rotation Experiment

Ba/2.5

Barley. Straw: cwt. per acre

	Treatments applied same year					Treatments applied previous year				
	Art	Adco	St ₁	St ₂	Mean	Art	Adco	St ₁	St ₂	Mean
Mean Yields										
1939	50.4	36.5	44.4	40.1	42.8	37.5	35.2	33.9	37.2	36.0
1940	38.4	31.8	33.4	31.6	33.8	30.3	26.5	27.4	30.0	28.6
1941	37.3	26.0	34.9	29.2	31.8	27.4	24.4	27.6	28.5	27.0
1942	35.7	26.0	29.9	30.0	30.4	26.5	26.8	27.0	25.0	26.3
1943	34.4	24.4	31.9	29.7	30.1	24.2	24.7	24.8	26.6	25.1
1944	33.5	29.0	29.7	31.6	31.0	29.8	30.1	28.8	29.7	29.6
1945	37.4	27.2	36.1	34.1	33.7	26.5	29.0	29.3	30.5	28.8
1946	40.4	37.5	38.0	36.9	38.2	29.4	30.7	31.7	32.8	31.2
1947	26.5	22.3	27.2	22.8	24.7	20.0	18.5	20.0	21.0	19.9

Responses to Magnesium Sulphate

1943	-0.1	3.6	3.5	0.3	1.8	-4.3	2.1	1.1	-4.8	-1.5
1944	0.5	0.8	-7.5	2.4	-1.0	-4.3	-2.5	-0.7	-0.7	-2.0
1945	-2.0	0.0	2.0	-2.9	-0.7	-1.3	-9.1	-2.7	1.3	-3.0
1946	0.8	0.5	1.7	-5.4	-0.6	-2.6	3.3	5.2	0.4	1.6
1947	-4.4	-4.5	-1.0	-0.1	-2.5	-0.2	2.7	-3.2	0.5	-0.1

Ba/2.6

Sugar Beet. Roots (washed): tons per acre

	Treatments applied same year					Treatments applied previous year				
	Art	Adco	St ₁	St ₂	Mean	Art	Adco	St ₁	St ₂	Mean
Mean Yields										
1939	14.00	12.21	13.01	13.63	13.21	12.11	11.36	11.93	13.07	12.12
1940	10.47	9.45	9.95	11.41	10.32	10.14	9.43	9.59	9.49	9.66
1941	11.53	10.43	10.86	11.61	11.11	10.09	10.36	10.24	10.63	10.33
1942	14.24	10.02	13.52	13.08 ±0.714	12.72 ±0.357	11.87	10.49	11.37	11.09	11.20 ±0.357
1943	14.32	11.50	13.00	12.30 ±0.806	12.78 ±0.403	12.10	11.17	10.43	12.40	11.52 ±0.403
1944	8.87	8.43	7.48	7.90 ±1.05	8.17 ±0.525	7.73	7.81	7.91	8.36	7.95 ±0.525
1945	13.99	12.04	12.77	12.86 ±0.897	12.92 ±0.448	12.37	11.26	13.01	11.47	12.03 ±0.443
1946	14.50	12.16	12.51	13.36 ±0.311	13.13 ±0.156	12.33	11.50	12.17	12.42	12.10 ±0.156
1947	10.89	7.99	10.02	9.63 ±0.741	9.09 ±0.370	8.97	9.29	8.94	9.07	9.07 ±0.370
Responses to Magnesium Sulphate										
1943	-0.74	0.60	0.06	-1.14 ±1.71	-0.30 ±0.855	-0.17	0.26	-2.66	0.12	-0.61 ±0.855
1944	3.00	-3.07	0.10	0.38 ±2.23	0.10 ±1.11	1.91	-1.06	1.02	-1.60	0.06 ±1.11
1945	-0.46	-0.38	-0.23	-0.10 ±1.90	-0.29 ±0.951	-0.07	0.85	-0.22	2.11	0.24 ±0.951
1946	-0.93	0.68	-0.11	0.85 ±0.661	0.11 ±0.330	0.39	0.47	0.17	-0.68	0.09 ±0.330
1947	3.55	-2.61	-0.44	1.36 ±1.57	0.60 ±0.786	1.09	-0.97	-0.33	-0.48	-0.17 ±0.786

Standard errors per plot

d.f.

1942	1.24 tons per acre or 10.3%	16
1943	1.40 tons per acre or 11.5%	8
1944	1.82 tons per acre or 22.6%	8
1945	1.55 tons per acre or 12.5%	8
1946	0.539 tons per acre or 4.3%	8
1947	1.28 tons per acre or 13.7%	8

Sugar Beet. Total Sugar: cwt. per acre

	Treatments applied same year					Treatments applied previous year				
	Art	Adco	St ₁	St ₂	Mean	Art	Adco	St ₁	St ₂	Mean
Mean Yields										
1939	51.4	45.7 ±1.97	40.9	50.2	49.0 ±0.987	44.2	42.8 ±1.97	43.3	47.7	44.5 ±0.987
1940	44.2	40.1 ±2.47	41.9	48.9	43.8 ±1.23	42.6	39.9 ±2.47	41.3	40.1	41.0 ±1.23
1941	46.3	42.7 ±2.26	43.7	47.2	45.0 ±1.13	40.5	41.4 ±2.26	41.0	42.4	41.3 ±1.13
1942	51.5	36.4 ±2.54	49.2	48.0	46.3 ±1.27	42.1	38.1 ±2.54	42.0	40.7	40.7 ±1.27
1943	52.8	41.8 ±2.75	48.1	44.9	46.9 ±1.38	44.9	40.8 ±2.75	38.5	46.1	42.6 ±1.38
1944	29.6	28.0 ±3.57	24.2	26.1	27.0 ±1.79	26.0	26.5 ±3.57	26.7	27.9	26.3 ±1.79
1945	56.5	48.4 ±3.66	52.9	52.4	52.6 ±1.83	50.6	45.2 ±3.66	51.8	46.8	48.6 ±1.83
1946	52.5	45.0 ±1.13	46.3	49.1	48.2 ±0.566	45.3	42.2 ±1.13	44.0	46.4	44.5 ±0.566
1947	45.7	33.3 ±3.29	43.0	40.2	40.5 ±1.64	38.7	37.1 ±3.29	38.8	37.2	37.9 ±1.64
<i>Re</i> Responses to Magnesium Sulphate										
1943	-2.5	3.0 ±5.84	-0.9	-3.5	-1.0 ±2.92	-2.7	0.6 ±5.84	-9.4	0.5	-2.8 ±2.92
1944	10.6	-11.8 ±7.57	-0.3	2.3	0.02 ±3.79	6.3	-4.6 ±7.57	2.7	2.3	0.17 ±3.79
1945	-1.0	-0.1 ±7.77	-0.9	0.0	-0.5 ±3.88	1.0	6.7 ±7.77	-2.1	8.4	3.5 ±3.88
1946	-3.2	2.8 ±2.40	-0.9	4.1	0.7 ±1.20	1.7	3.5 ±2.40	0.4	-2.2	0.8 ±1.20
1947	14.9	-10.4 ±6.97	-1.8	7.8	2.6 ±3.48	6.0	-5.1 ±6.97	-3.7	-1.9	-1.2 ±3.48

Standard errors per plot d.f.

1939	3.42 cwt. per acre or	7.3%	16
1940	4.27 cwt. per acre or	10.1%	16
1941	3.91 cwt. per acre or	9.0%	16
1942	4.40 cwt. per acre or	10.1%	16
1943	4.77 cwt. per acre or	10.7%	8
1944	6.18 cwt. per acre or	23.0%	8
1945	6.34 cwt. per acre or	12.5%	8
1946	1.96 cwt. per acre or	4.2%	8
1947	5.69 cwt. per acre or	14.5%	8

Sugar Beet. Sugar Percentage

	Treatments applied same year					Treatments applied previous year				
	Art	Adco	St ₁	St ₂	Mean	Art	Adco	St ₁	St ₂	Mean
Mean Yields										
1939	18.35	18.71	18.78	18.38	18.56	18.27	18.83	18.12	18.25	18.37
1940	21.10	21.25	21.03	21.46	21.21 ±0.325	20.98	21.19	21.53	21.10	21.20 ±0.162
1941	20.10	20.48	20.11	20.32	20.25 ±0.227	20.08	19.96	20.00	19.99	20.01 ±0.113
1942	18.08	18.17	18.22	18.35	18.20 ±0.192	17.75	18.19	18.48	18.38	18.20 ±0.096
1943	18.43	18.18	18.52	18.28	18.35 ±0.171	18.52	18.30	18.47	18.60	18.47 ±0.086
1944	16.67	16.48	16.13	16.40	16.43 ±0.126	16.85	16.93	16.86	16.68	16.83 ±0.063
1945	20.21	20.11	20.74	20.39	20.36 ±0.174	20.47	20.05	19.91	20.38	20.20 ±0.087
1946	18.13	18.52	18.51	18.38	18.38 ±0.165	18.36	18.37	18.10	18.68	18.38 ±0.082
1947	20.97	20.91	21.44	20.86	21.04	21.26	20.63	20.88	20.76	20.88

Responses to Magnesium Sulphate

1943	0.09	0.30	-0.43 ±0.364	0.27	0.06 ±0.182	-0.83	-0.23 ±0.364	0.21	0.01	-0.21 ±0.182
1944	0.28	-1.14	-0.40 ±0.268	0.79	-0.12 ±0.134	-0.08	-0.70 ±0.268	-0.44	0.37	-0.21 ±0.134
1945	0.25	0.60	-0.01 ±0.368	0.14	0.24 ±0.184	0.51	1.42 ±0.368	-0.46	-0.06	0.35 ±0.184
1946	0.11	0.08	-0.20 ±0.349	0.39	0.10 ±0.175	0.12	0.80 ±0.349	-0.10	0.10	0.23 ±0.175
1947	0.00	0.36	0.04	0.05	0.11	0.69	-0.61 ±0.285	-1.24	0.14	-0.25

Standard errors per plot		d.f.
1940	0.562	16
1941	0.393	16
1942	0.333	16
1943	0.297	8
1944	0.219	8
1945	0.301	8
1946	0.285	8

Three Course Rotation Experiment

Ba/2.9

Sugar Beet. Tops: tons per acre

	Treatments applied same year					Treatments applied previous year				
	Art	Adco	St ₁	St ₂	Mean	Art	Adco	St ₁	St ₂	Mean
Mean Yields										
1939	15.09	11.40	12.83	12.52	12.96	12.33	10.24	12.08	12.22	11.72
1940	8.16	6.46	7.35 ±0.491	8.06	7.51 ±0.245	6.80	6.74	6.32 ±0.491	6.42	6.57 ±0.245
1941	11.91	8.64	9.32 ±0.675	10.87	10.18 ±0.337	8.26	9.57	8.47 ±0.675	9.15	8.86 ±0.337
1942	11.39	7.32	10.14 ±0.663	9.79	9.66 ±0.332	8.87	7.93	8.35 ±0.663	7.80	8.24 ±0.332
1943	11.52	8.43	8.74 ±1.02	8.74	9.36 ±0.512	7.72	8.27	6.25 ±1.02	8.15	7.60 ±0.512
1944	10.35	9.08	8.57 ±0.693	9.17	9.29 ±0.346	8.09	7.77	7.20 ±0.693	9.07	8.03 ±0.346
1945	9.42	6.47	7.70 ±0.315	6.64	7.56 ±0.158	6.31	6.13	7.30 ±0.315	6.98	6.68 ±0.158
1946	13.84	10.54	11.39 ±1.17	11.96	11.93 ±0.585	10.80	11.03	9.16 ±1.17	10.49	10.37 ±0.585
1947	6.12	4.38	5.56 ±0.280	5.83	5.47 ±0.140	4.44	4.96	4.87 ±0.280	5.05	4.83 ±0.140

Responses to Magnesium Sulphate

1943	-1.40	0.68	-0.83 ±2.17	-0.32	-0.47 ±1.09	-0.06	-0.15	-0.81 ±2.17	-0.07	-0.27 ±1.09
1944	1.22	-1.36	1.98 ±1.47	-1.44	0.10 ±0.735	2.05	0.16	0.16 ±1.47	-0.54	0.46 ±0.735
1945	-2.07	-0.93	-1.17 ±0.668	-0.18	-1.09 ±0.334	-0.49	-0.02	2.51 ±0.668	0.65	0.66 ±0.334
1946	-0.87	-1.00	-1.41 ±2.48	1.40	-0.47 ±1.24	-0.35	-0.47	-0.88 ±2.48	-0.60	-0.58 ±1.24
1947	1.14	-0.71	-0.47 ±0.594	-0.31	-0.09 ±0.297	-0.10	0.54	0.49 ±0.594	-0.95	0.04 ±0.297

Standard errors per plot

d.f.

1939	1.13 tons per acre or 1940	0.850 tons per acre or 1941	1.17 tons per acre or 1942	1.15 tons per acre or 1943	1.77 tons per acre or 1944	1.20 tons per acre or 1945	0.546 tons per acre or 1946	2.03 tons per acre or 1947	0.485 tons per acre or 9.2%	16
										16
										16
										16
										8
										8
										8
										8
										8

FOUR COURSE ROTATION EXPERIMENT

Great Hoos (begun in 1930)Residual Values of Humic and Phosphatic Fertilizers

The five treatments compared in this experiment are dung, Adco compost, straw with artificials, superphosphate and rock phosphate (Gafsa), the cropping following a Norfolk Rotation (potatoes, barley, ryegrass, wheat). There are four series (one for each crop in any particular year), each consisting of twenty-five plots.

Dung and Adco each supply 50 cwt. per acre of organic matter. The quantity of straw applied is equal to that used in making the Adco. The nutrient content of the three humic fertilizers is equalized by adding sulphate of ammonia, muriate of potash and superphosphate, to raise the applications to 1.8 cwt. N per acre, 1.2 cwt. P₂O₅ per acre and 3.0 cwt. K₂O per acre. The phosphatic fertilizers are applied at the rate of 1.2 cwt. P₂O₅ per acre, together with sulphate of ammonia and muriate of potash at the above rates.

Any given plot receives always the same treatment, but the treatment is applied to the plot only once in five years (except that the sulphate of ammonia and muriate of potash accompanying the phosphatic fertilizers are applied one fifth annually.) Thus each treatment is applied to one plot of each series every year, according to a Latin square scheme. In this way the residual effects of fertilizers are measured in every crop, and the period of the manurial cycle differs from that of the crop rotation.

Area of each plot, 0.024 acre.

Details of the experiment are as given in the 1932 Report, pp. 127-8, with the following alterations:-

1. From 1935 onwards, clover ryegrass ley was replaced by ryegrass alone, sown in autumn after ploughing barley stubble, with fertilizers applied as on wheat.
2. From 1942 onwards, each plot of the potato crop was split, a random half of each receiving an additional 0.4 cwt. N per acre as sulphate of ammonia each year.

Nutrients provided by organic fertilizers: cwt. per acre

Year	Manure as F.Y.M.			Manure as Adco			Manure as Straw			
	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	Organic Matter	N	P ₂ O ₅	K ₂ O
1939	1.782	1.344 [#]	4.503	1.646	1.520 [#]	1.189	123.75	0.862	0.480	2.652
1940	1.520	0.830	2.155	1.562	1.240 [#]	1.330	139.88	1.008	0.520	2.699
1941	1.775	0.471	1.802	1.260	0.877	0.538	106.74	0.573	0.183	1.526
1942	1.410	0.700	2.373	1.378	0.890	0.506	118.96	0.615	0.208	1.229
1943	1.526	0.645	2.193	1.282	0.840	0.474	133.23	0.832	0.270	1.506
1944	1.202	0.409	1.925	1.313	0.706	0.572	100.00	0.657	0.178	0.765
1945	1.524	0.492	2.712	0.995	0.555	0.622	82.00	0.297	0.088	0.684
1946	1.802	0.710	2.418	1.461	1.031	0.892	124.90	0.683	0.379	1.447
1947	1.588	0.748	1.560	1.585	1.169	0.911	123.00	0.849	0.261	1.546

[#] No application of artificials necessary.

Crop Notes

Crop year	1939	1940	1941	1942	1943	1944	1945	1946	1947

Potatoes. Variety: 1939-41, Ally; 1942-47, Majestic. Previous crop: Wheat

Planted	Apr. 17	Apr. 26	Apr. 30	Apr. 21	Apr. 8	Mar. 25	Mar. 30	Mar. 30	Apr. 29
Harvested	Sept. 19	Sept. 24	Oct. 3	Oct. 3	Sept. 15	Sept. 28	Sept. 25	Oct. 4	Sept. 30

Barley. Variety: Plumage Archer. Previous crop: Potatoes

Sown	Mar. 7	Mar. 11	Mar. 18	Mar. 25	Mar. 2	Mar. 6	Mar. 27	Mar. 20	Apr. 12
Harvested	Aug. 23	Aug. 6	Sept. 2	Aug. 17	Aug. 7	Aug. 12	Aug. 14	Aug. 21	Aug. 14

Ryegrass. Variety: Western Wolths. Previous crop: Barley

Sown	Sept. 13	Apr. 13 ¹	Oct. 30	Nov. 1	Sept. 24	Sept. 30	Sept. 13	Sept. 17	Apr. 12 ³
Harvested	June 6	June 22	Failed ²	June 20	June 14	June 14	June 13	June 14	Aug. 21

Wheat. Variety: 1939-45 Yeoman. 1946 Squareheads Master. Previous crop: Ryegrass

Sown	Oct. 24	Oct. 21	Oct. 29	Oct. 27	Oct. 10	Oct. 4	Oct. 26	Oct. 15	Oct. 24
Harvested	Aug. 14	Aug. 6	Aug. 22	Aug. 12	Aug. 3	Aug. 12	Aug. 13	Aug. 21	Aug. 18

(1) First sowing of ryegrass failed.

(2) Barley sown to replace ryegrass, which failed.

Sown: Apr 22. Cut for hay: July 23. Variety: Kenia.

(3) Wet season prevented autumn sowing of ryegrass.

Potatoes. Total tubers: tons per acre

Manure	Year of cycle	1939	1940	1941	1942	1943	1944	1945	1946	1947
Manure as F.Y.M.	1	5.28	4.95	3.02	9.08	6.90	10.68	9.60	7.29	6.13
	2	3.52	4.34	2.53	6.12	6.20	7.85	8.14	7.03	5.47
	3	4.03	3.01	2.69	6.56	5.48	11.69	7.27	5.54	5.84
	4	2.82	4.80	1.92	6.77	5.38	8.05	8.87	5.06	5.05
	5	4.16	3.79	2.42	3.82	5.46	8.97	6.11	6.38	4.30
Manure as Adco	1	4.14	4.87	1.86	7.77	8.34	12.49	11.82	9.24	6.05
	2	3.70	2.91	2.38	5.39	4.77	9.66	7.72	7.26	5.59
	3	2.25	2.91	2.04	4.28	4.72	8.55	6.80	6.60	4.56
	4	3.12	2.93	1.30	4.57	5.13	7.97	7.01	7.04	5.21
	5	2.46	2.05	1.75	5.15	5.26	9.63	7.08	6.18	5.13
Manure as Straw	1	3.59	7.18	4.42	10.22	7.13	12.60	12.83	9.13	6.57
	2	4.07	5.11	2.77	6.85	5.97	8.21	9.00	6.37	4.36
	3	4.23	3.45	3.00	7.93	6.62	6.88	8.43	7.20	4.55
	4	4.22	2.51	1.88	4.07	5.81	8.80	8.15	6.19	5.66
	5	3.48	3.50	2.76	6.73	4.92	10.02	7.79	6.64	5.07
Super- Phosphate	1	5.63	5.66	4.03	6.90	8.51	9.98	9.27	9.19	6.48
	2	5.72	4.58	4.14	4.46	6.29	10.86	9.45	7.58	5.63
	3	4.45	4.67	2.24	9.14	7.28	9.53	9.27	7.39	6.36
	4	3.98	3.56	2.76	7.23	8.40	11.06	9.84	8.41	5.98
	5	4.76	4.98	2.64	7.31	6.85	8.03	10.01	8.02	6.17
Gafsa Rock Phosphate	1	3.37	2.44	2.41	4.06	5.30	8.62	9.31	7.18	4.04
	2	2.22	3.37	2.26	5.57	6.58	9.60	7.67	6.41	5.41
	3	3.59	4.58	2.45	5.83	7.01	6.63	8.13	6.85	5.26
	4	3.81	2.29	2.85	4.13	5.88	8.72	7.93	6.24	5.47
	5	2.90	3.28	1.93	7.52	5.55	5.27	8.48	8.35	4.90

Potatoes. Total tubers: tons per acre. Effect of additional nitrogen

Manures	Year of Cycle	1942			N Effect	1943			N Effect
		Additional N Without	With	Mean		Additional N Without	With	Mean	
Manure as F.Y.M.	1	9.08	10.34	9.71	1.26	6.90	7.28	7.09	0.38
	2	6.12	8.31	7.22	2.19	6.20	7.99	7.10	1.79
	3	6.56	8.04	7.30	1.48	5.48	6.15	5.82	0.67
	4	6.77	7.08	6.92	0.31	5.38	7.99	6.68	2.61
	5	3.82	4.65	4.24	0.83	5.46	7.72	6.59	2.26
Manure as Adco	1	7.77	7.16	7.46	-0.61	8.34	9.20	8.77	0.86
	2	5.39	7.81	6.60	2.42	4.77	6.59	5.68	1.82
	3	4.28	4.61	4.44	0.33	4.72	6.62	5.67	1.90
	4	4.57	6.09	5.33	1.52	5.13	4.83	4.98	-0.30
	5	5.15	5.24	5.20	0.09	5.26	6.68	5.97	1.42
Manure as Straw	1	10.22	11.73	10.98	1.51	7.13	9.31	8.22	2.18
	2	6.85	9.67	8.26	2.82	5.97	7.63	6.80	1.66
	3	7.93	7.52	7.72	-0.41	6.62	8.03	7.32	1.41
	4	4.07	4.02	4.04	-0.05	5.81	7.95	6.88	2.14
	5	6.73	7.16	6.94	0.43	4.92	6.43	5.68	1.51
Super- Phosphate	1	6.90	8.31	7.60	1.41	8.51	10.41	9.46	1.90
	2	4.46	5.65	5.06	1.19	6.29	8.22	7.26	1.93
	3	9.14	10.04	9.59	0.90	7.28	8.58	7.93	1.30
	4	7.23	7.85	7.54	0.62	8.40	9.10	8.75	0.70
	5	7.31	7.19	7.25	-0.12	6.85	7.48	7.16	0.63
Gafsa Rock Phosphate	1	4.06	3.74	3.90	-0.32	5.30	5.63	5.46	0.33
	2	5.57	7.81	6.69	2.24	6.58	5.80	6.19	-0.78
	3	5.83	4.98	5.40	-0.85	7.01	8.19	7.60	1.18
	4	4.13	6.52	5.32	2.39	5.88	6.14	6.01	0.26
	5	7.52	5.64	6.58	-1.88	5.55	6.31	5.93	0.76

Four Course Rotation Experiment

Potatoes. Total tubers: tons per acre. Effect of additional nitrogen

Manures	Year of Cycle	1944			N Effect	1945			N Effect
		Additional N Without	With	Mean		Additional N Without	With	Mean	
Manure as F.Y.M.	1	10.68	12.72	11.70	2.04	9.60	12.77	11.18	3.17
	2	7.85	8.03	7.94	0.18	8.14	12.27	10.20	4.13
	3	11.69	12.09	11.89	0.40	7.27	10.50	8.88	3.23
	4	8.05	8.80	8.42	0.75	8.87	10.71	9.79	1.84
	5	8.97	11.90	10.44	2.93	6.11	8.03	7.07	1.92
Manure as Adco	1	12.49	10.96	11.72	-1.53	11.82	13.69	12.76	1.87
	2	9.66	11.10	10.38	1.44	7.72	9.62	8.67	1.90
	3	8.55	9.54	9.04	0.99	6.80	8.92	7.86	2.12
	4	7.07	8.69	8.33	0.72	7.01	9.27	8.14	2.26
	5	9.63	10.13	9.88	0.50	7.08	8.07	7.58	0.99
Manure as Straw	1	12.60	13.90	13.25	1.30	12.83	9.95	11.39	-2.88
	2	8.21	9.25	8.73	1.04	9.00	8.39	8.70	-0.61
	3	6.88	7.11	7.00	0.23	8.43	11.60	10.02	3.17
	4	8.80	9.14	8.97	0.34	8.15	9.52	8.84	1.37
	5	10.02	10.29	10.16	0.27	7.79	10.49	9.14	2.70
Super- Phosphate	1	9.98	10.99	10.48	1.01	9.27	9.82	9.54	0.55
	2	10.86	11.90	11.38	1.04	9.45	10.83	10.14	1.38
	3	9.53	10.80	10.16	1.27	9.27	10.49	9.88	1.22
	4	11.06	12.09	11.58	1.03	9.84	10.52	10.18	0.68
	5	8.03	9.03	8.53	1.00	10.01	10.56	10.28	0.55
Gafsa Rock Phosphate	1	8.62	9.34	8.98	0.72	9.31	8.46	8.88	-0.85
	2	9.60	12.05	10.82	2.45	7.67	7.88	7.78	0.21
	3	6.63	8.60	7.62	1.97	8.13	8.99	8.56	0.86
	4	8.72	8.13	8.42	-0.59	7.93	8.09	8.01	0.16
	5	5.27	7.42	6.34	2.15	8.48	6.98	7.73	-1.50

Potatoes, Total tubers: tons per acre. Effect of additional nitrogen

Manures	Year of Cycle	1946			1947			N Effect
		Additional N Without	With	Mean	Additional N Without	With	Mean	
Manure as F.Y.M.	1	7.29	8.65	7.97	1.36	6.13	8.06	7.10 1.93
	2	7.03	8.80	7.92	1.77	5.47	6.55	6.01 1.08
	3	5.54	8.00	6.77	2.46	5.84	6.36	6.10 0.52
	4	5.06	8.63	6.84	3.57	5.05	5.97	5.51 0.92
	5	6.38	6.79	6.58	0.41	4.30	5.39	4.84 1.09
Manure as Adco	1	9.24	9.18	9.21	-0.06	6.05	6.01	6.03 -0.04
	2	7.26	8.78	8.02	1.52	5.59	5.42	5.50 -0.17
	3	6.60	8.58	7.59	1.98	4.56	5.89	5.22 1.33
	4	7.04	7.09	7.06	0.05	5.21	5.94	5.58 0.73
	5	6.18	7.53	6.86	1.35	5.13	4.65	4.89 -0.48
Manure as Straw	1	9.13	9.93	9.53	0.80	6.57	6.35	6.46 -0.22
	2	6.37	10.66	8.52	4.29	4.36	5.80	5.08 1.44
	3	7.20	9.09	8.14	1.89	4.55	6.74	5.64 2.19
	4	6.19	7.26	6.72	1.07	5.66	6.43	6.04 0.77
	5	6.64	8.52	7.58	1.88	5.07	6.34	5.70 1.27
Super- Phosphate	1	9.19	10.08	9.64	0.89	6.48	7.41	6.94 0.93
	2	7.58	9.78	8.68	2.20	5.63	5.59	5.61 -0.04
	3	7.39	7.99	7.69	0.60	6.36	6.40	6.38 0.04
	4	8.41	8.18	8.30	-0.23	5.98	5.34	5.66 -0.64
	5	8.02	7.51	7.76	-0.51	6.17	5.62	5.90 -0.55
Gafsa Rock Phosphate	1	7.18	6.17	6.68	-1.01	4.04	4.70	4.37 0.66
	2	6.41	6.01	6.21	-0.40	5.41	5.65	5.53 0.24
	3	6.85	6.61	6.73	-0.24	5.26	5.21	5.24 -0.05
	4	6.24	6.20	6.22	-0.04	5.47	5.18	5.32 -0.29
	5	8.35	7.02	7.68	-1.33	4.90	4.79	4.84 -0.11

Four Course Rotation Experiment

Manure	Year of Cycle	Barley Grain: cwt. per acre								
		1939	1940	1941	1942	1943	1944	1945	1946	1947
Manure as F.Y.M	1	26.3	27.6	18.4	35.0	27.4	35.8	29.8	26.9	22.6
	2	25.9	32.5	14.6	24.1	23.3	29.4	21.8	24.2	16.2
	3	22.8	24.2	12.5	21.6	17.9	25.2	19.0	25.1	15.3
	4	23.7	20.7	15.3	22.0	19.6	23.0	19.2	22.5	13.3
	5	21.5	22.6	12.2	16.3	16.1	27.2	17.6	23.8	14.6
Manure as Adco	1	31.5	36.5	17.3	28.4	32.4	31.6	31.5	30.2	19.0
	2	26.1	28.1	13.8	18.6	22.7	30.3	22.4	25.2	13.8
	3	20.6	24.3	12.6	18.9	16.4	24.9	18.1	21.6	15.3
	4	20.5	18.5	10.4	17.0	20.9	27.6	16.6	24.5	13.4
	5	22.4	27.5	11.0	13.9	16.3	27.5	14.8	21.8	14.5
Manure as Straw	1	23.7	34.5	25.5	32.0	32.2	34.8	38.4	34.3	24.0
	2	24.5	27.3	15.1	26.6	23.1	28.2	24.8	27.2	16.5
	3	20.8	27.1	12.0	19.2	20.8	28.1	19.6	25.6	19.1
	4	23.8	26.0	12.6	17.2	21.6	26.0	16.8	24.3	14.1
	5	26.3	24.2	12.6	21.4	13.4	25.1	20.0	24.8	15.8
Super- phosphate	1	32.2	30.2	25.1	25.4	27.8	29.2	26.2	28.6	23.7
	2	32.3	30.8	24.8	27.2	25.3	28.6	26.3	22.8	22.6
	3	31.7	31.1	25.9	26.4	30.0	26.7	24.2	27.2	22.6
	4	31.7	31.3	22.0	28.6	28.3	27.5	22.7	29.1	22.0
	5	33.0	29.7	21.7	29.2	22.7	29.4	26.6	29.2	20.4
Gafsa Rock Phosphate	1	29.0	33.2	18.7	13.1	23.7	28.8	20.0	29.3	20.6
	2	29.8	34.5	15.0	27.7	24.6	28.0	21.4	28.9	17.4
	3	32.9	31.0	17.3	27.9	24.1	28.8	26.2	23.3	20.6
	4	31.5	30.4	22.3	22.8	20.9	28.1	21.2	30.3	19.9
	5	26.5	32.1	16.9	30.4	26.1	29.9	23.4	27.4	19.4

Ba/3.8

Barley Straw: cwt. per acre

Manure	Year of Cycle	1939	1940	1941	1942	1943	1944	1945	1946	1947
Manure as F.Y.M.	1	32.2	28.5	23.0	37.5	30.1	33.3	33.8	29.6	21.9
	2	28.8	25.3	23.2	22.2	23.8	29.1	23.7	28.4	17.5
	3	26.1	24.7	19.8	20.4	19.4	23.5	21.0	26.4	16.0
	4	27.6	22.0	20.8	20.7	19.0	22.8	20.9	28.4	13.5
	5	28.2	24.7	18.4	16.5	15.8	25.0	17.2	29.5	16.6
Manure as Adco	1	39.7	35.7	27.0	31.0	34.0	31.4	37.4	40.0	20.5
	2	32.0	29.2	21.2	23.9	21.8	26.4	25.6	29.4	14.3
	3	24.6	24.5	23.0	22.3	16.7	23.5	21.1	22.8	15.6
	4	26.5	27.7	17.0	23.6	21.8	26.4	21.8	33.2	14.3
	5	24.7	28.2	20.5	15.8	17.6	23.1	18.7	28.2	16.3
Manure as Straw	1	27.2	37.8	32.1	35.8	33.9	34.1	34.4	42.0	23.8
	2	29.0	27.7	21.6	25.6	22.6	26.8	27.8	34.0	17.7
	3	24.4	27.8	20.0	19.7	22.4	25.9	28.4	30.5	18.7
	4	27.8	26.4	18.9	21.1	20.7	23.6	25.4	30.0	15.7
	5	27.6	24.7	21.3	16.9	14.9	22.1	20.7	26.9	16.3
Super- phosphate	1	35.9	35.2	30.6	30.6	28.3	26.9	31.6	36.4	24.7
	2	39.6	32.3	30.8	26.5	28.0	28.2	25.4	28.2	22.6
	3	34.1	33.4	29.8	20.4	28.7	25.5	29.4	34.4	23.7
	4	33.7	32.9	29.7	28.0	27.6	24.0	29.4	32.5	22.7
	5	34.7	30.3	31.8	27.3	26.0	24.9	29.3	38.2	21.1
Gafsa Rock Phosphate	1	33.3	37.9	29.1	22.8	26.0	25.6	26.5	36.5	24.0
	2	36.8	39.8	29.2	31.3	27.4	26.2	25.1	34.7	18.9
	3	37.1	37.5	28.5	29.5	26.3	26.5	27.8	29.1	22.8
	4	37.3	32.3	*	23.4	24.1	25.1	29.4	35.5	22.6
	5	36.2	34.8	32.0	29.9	26.7	28.2	25.1	36.4	23.0

Recorded yield obviously incorrect.

Four Course Rotation Experiment

Ryegrass: cwt. per acre dry matter

Manure	Year of Cycle	1939	1940	1941	1942	1943	1944	1945	1946	1947
Manure as F.Y.M.	1	20.4	19.4	7.0	18.8	16.5	19.4	17.3	33.3	6.7
	2	10.9	11.4	9.7	9.5	12.7	10.2	13.1	22.8	3.2
	3	10.8	7.1	5.5	7.3	10.5	11.4	11.5	20.8	3.3
	4	8.5	9.1	6.5	9.0	8.9	8.2	11.8	12.3	3.2
	5	9.5	9.1	5.8	5.6	8.6	9.6	9.6	16.2	3.2
Manure as Adco	1	14.4	16.9	10.1	9.4	17.8	18.7	25.3	31.9	6.7
	2	10.2	9.9	6.7	5.8	10.6	13.3	14.0	27.6	3.5
	3	8.8	7.6	5.6	9.0	9.6	9.0	11.5	18.9	3.5
	4	7.2	8.9	5.4	6.1	8.3	9.1	8.1	16.5	1.2
	5	8.1	8.5	3.6	6.2	8.8	9.8	10.4	15.5	3.9
Manure as Straw	1	40.5	28.1	14.3	21.5	29.3	16.4	39.2	59.1	12.4
	2	11.0	8.8	4.4	8.6	9.4	10.7	13.2	21.7	3.7
	3	9.2	12.1	6.6	11.4	11.4	10.6	12.5	23.5	5.0
	4	9.3	9.8	7.0	9.6	8.3	11.8	10.6	15.5	4.4
	5	9.2	9.3	6.9	7.1	8.1	9.6	8.5	14.0	3.1
Super- phosphate	1	22.5	18.4	10.0	13.5	14.2	12.6	22.0	39.7	11.2
	2	23.3	12.9	10.0	15.6	18.1	12.0	20.0	33.7	7.8
	3	17.0	14.8	9.4	10.8	15.1	14.3	22.3	34.0	4.3
	4	20.2	10.9	11.6	7.1	13.7	15.4	19.2	36.7	6.9
	5	20.5	13.3	7.5	11.0	17.1	12.6	18.3	35.9	6.6
Gafsa Rock Phosphate	1	17.7	7.8	9.0	6.2	16.6	12.8	20.8	31.6	7.2
	2	16.1	12.0	9.3	9.7	17.7	8.5	18.7	31.4	5.8
	3	16.1	10.7	8.4	3.6	14.8	10.4	19.2	32.3	8.4
	4	18.5	11.5	6.7	10.2	15.4	11.2	16.9	37.1	5.4
	5	24.9	10.4	5.4	15.6	11.8	8.9	18.3	29.0	7.2

In 1941 the ryegrass failed and the plots were resown with barley, which was cut for hay.

Wheat Grain: cwt. per acre

Manure	Year of Cycle	1939	1940	1941	1942	1943	1944	1945	1946	1947
Manure as F.Y.M.	1	23.9	26.0	18.0	19.9	24.9	14.1	24.6	22.5	12.7
	2	19.3	20.1	16.4	17.8	20.0	7.8	22.2	17.3	11.2
	3	15.4	16.3	15.4	19.0	15.6	12.1	17.9	12.0	10.4
	4	16.2	17.4	12.5	14.6	15.0	8.2	19.4	14.6	7.8
	5	14.6	18.1	14.1	14.9	14.0	7.9	18.1	17.3	8.9
Manure as Adco	1	19.3	26.3	17.7	22.9	24.0	14.1	27.5	22.7	12.8
	2	17.0	18.9	15.1	18.5	14.9	10.4	19.4	19.1	11.1
	3	16.4	16.3	15.6	17.4	15.9	9.2	19.3	14.4	9.1
	4	12.6	16.3	13.7	23.4	15.8	11.6	16.9	12.2	9.2
	5	17.4	16.8	12.1	15.1	12.7	9.9	15.6	12.6	7.3
Manure as Straw	1	23.9	28.7	24.4	16.9	28.4	18.9	30.3	28.7	11.3
	2	17.4	19.1	13.7	13.9	17.0	6.4	20.0	14.1	11.2
	3	15.8	18.2	14.0	18.2	16.1	12.1	20.8	16.4	10.6
	4	17.4	25.7	16.2	18.1	16.2	7.3	17.7	12.6	10.3
	5	14.3	18.8	11.9	18.4	14.0	7.6	17.4	14.7	8.9
Super- Phosphate	1	22.7	18.8	16.5	20.6	21.4	9.4	22.1	15.9	8.6
	2	21.4	19.3	17.4	17.8	17.9	10.2	21.0	12.0	11.3
	3	19.4	21.8	17.9	20.1	22.2	11.0	22.1	16.0	9.0
	4	19.2	21.9	18.1	19.5	21.6	11.3	22.6	16.2	10.9
	5	21.3	21.4	18.9	21.0	21.4	9.5	23.4	17.9	10.2
Gafsa Rock	1	19.6	24.7	16.1	18.9	20.4	12.8	19.2	15.7	9.0
	2	22.5	21.6	14.5	19.9	20.2	17.1	21.2	17.7	7.5
	3	16.9	23.6	19.1	19.9	19.6	15.5	21.1	20.5	6.7
	4	21.7	21.6	18.0	22.0	21.0	15.9	21.9	16.1	2.4
	5	19.8	22.2	18.2	18.8	18.2	12.6	16.9	15.8	9.2

Four Course Rotation Experiment

Wheat Straw: cwt. per acre

Manure	Year of Cycle	1939	1940	1941	1942	1943	1944	1945	1946	1947
Manure as F.Y.M.	1	33.6	36.7	22.2	32.7	38.0	27.5	35.9	43.1	16.8
	2	26.9	25.7	20.1	24.6	28.5	12.1	31.2	35.3	13.4
	3	20.9	21.5	17.3	26.5	20.3	16.5	24.7	27.4	15.8
	4	26.3	24.9	14.4	20.9	26.1	11.7	25.7	34.3	10.0
	5	27.2	21.0	15.8	23.2	18.3	10.8	25.7	32.4	10.8
Manure as Adco	1	36.4	36.2	24.4	32.1	36.6	23.6	41.4	43.5	22.2
	2	23.3	22.4	17.4	25.0	25.3	14.7	27.1	36.4	17.7
	3	25.0	21.4	19.5	22.6	29.7	13.2	26.2	28.7	12.2
	4	26.6	21.3	16.9	32.5	22.2	13.7	22.1	24.9	13.3
	5	25.8	21.1	13.9	21.9	22.5	13.9	20.2	25.5	10.0
Manure as Straw	1	35.8	40.9	34.7	25.1	44.6	29.8	49.8	50.7	24.6
	2	23.7	21.6	16.9	20.4	29.9	8.8	27.4	28.2	13.2
	3	35.6	23.2	17.0	24.9	22.4	16.5	29.7	32.3	13.4
	4	24.4	31.7	18.1	24.4	22.5	10.3	26.1	27.0	14.5
	5	20.9	22.0	15.2	24.5	19.3	10.8	23.9	28.9	11.3
Super- phosphate	1	33.0	24.3	20.4	27.8	33.2	12.9	30.3	30.1	15.1
	2	33.2	25.8	22.2	26.0	31.0	12.7	30.3	23.9	16.4
	3	34.5	28.2	21.1	32.3	30.4	14.7	30.6	31.3	13.0
	4	27.7	30.5	22.0	26.6	32.7	13.9	33.2	30.7	14.4
	5	32.3	27.1	22.6	31.8	28.7	12.7	32.1	37.2	14.9
Gafsa Rock Phosphate	1	27.3	31.3	20.7	23.6	30.3	16.9	30.1	30.5	16.1
	2	29.9	27.3	18.0	28.1	35.7	20.8	29.6	35.4	9.5
	3	35.9	29.5	22.8	26.3	28.7	16.7	31.3	39.3	9.8
	4	30.3	28.7	21.4	28.8	33.4	18.4	32.4	32.5	4.5
	5	32.6	31.7	21.2	25.8	26.7	14.7	24.0	31.1	12.8

SIX COURSE ROTATION EXPERIMENT

ROTHAMSTED, Long Hoos IV, and WOBURN, Stackyard Field, Series B.

Experiment begun in 1930 (Rothamsted) and 1929 (Woburn)

Seasonal effects of N, P₂O₅ and K₂O

The six crops of the rotation are sugar beet, barley, clover hay, wheat, potatoes and rye. Both at Rothamsted and Woburn there are fifteen plots under each crop in any year, and each plot receives one of the three common fertilizers, sulphate of ammonia, superphosphate and muriate of potash, at one of five levels. The rates of application of fertilizers are

N and P₂O₅: 0.00, 0.15, 0.30, 0.45, 0.60 cwt. per acreK₂O: 0.00, 0.25, 0.50, 0.75, 1.00 cwt. per acre

Plots do not receive the same treatment throughout, but on each plot the fifteen treatments follow one another in a definite order in successive years, and in this way cumulative effects of treatments are avoided.

0.027

Area of each plot: Rothamsted 0.025 acre Woburn 0.0271 acre. X

Details of the experiment are as given in the 1932 Report, p.131, except that since 1934 the forage crop has been replaced by rye, harvested as a mature crop, and that the green manure crops are now omitted. Since 1934 lime has been applied at the rate of 10 cwt. per acre at two stages in the rotation: immediately after the removal of the potato crop and before sowing barley.

The increments given in the following tables are linear regressions of yield on quantity of fertilizer.

Crop Notes

	ROTHAMSTED			WOBURN		
	Variety	Sown	Harvested	Variety	Sown	Harvested
Sugar beet						
1939	Kuhn	Apr. 22	Oct. 24	Kuhn	May 4	Nov. 14
1940	"	May 3	Nov. 26	"	" 6	" 19
1941	"	Apr. 30	" 3	"	Apr. 29	Dec. 12
1942	Klein E.	" 24	Oct. 23	"	" 22	" 9
1943	" "	" 14	" 14	Klein E.	" 17	" 17
1944	" "	" 21	" 20	" "	May 4	" 18
1945	" "	" 12	" 18	" "	" 3	Nov. 28
1946	" "	" 13	" 22	" "	Apr. 16	Oct. 12
1947	" "	" 26	" 16	" "	May 3	" 17

ROTHAMSTED

WOBURN

	Variety	Sown	Harvested	Variety	Sown	Harvested
Barley						
1939	Flumage Archer	Mar. 7	Aug. 16	Flumage	Mar. 1	Aug. 30
1940	" "	" 9	" 3	Archer	" 9	" 30
1941	" "	" 25	Sep. 2	"	" 17	" 27
1942	" "	" 25	Aug. 17	"	Apr. 1	" 13
1943	" "	" 2	" 7	"	Mar. 4	Jul. 29
1944	" "	" 6	" 8	"	" 14	Aug. 28
1945	" "	" 26	" 6	"	" 5	" 2
1946	" "	" 20	" 15	"	" 19	" 22
1947	" "	Apr. 12	" 8	"	Apr. 17	" 6
Clover						
1939	Crimson Clover	Apr. 24 [#]	Jul. 28	Broad Red	31.3.38,	Jun. 9
1940	Montgomery Red	23.4.39	Jun. 17	" "	14.9.38	
1941	" "	May 3 [#]	Aug. 25	" "	17.4.39	" 20
1942	" "	9.5.41	Jun. 27	" "	6.5.40	Jul. 10
1943	" "	7.5.42	Jul. 15	" "	5.5.41	Jun. 26
1944	" "	Mar. 27	" 28	" "	20.5.42	" 7
1945	Crimson Clover	12.9.44	Jun. 13	" "	16.4.43	" 5
1946	Montgomery Red	28.3.45	" 22	Montgomery	Mar. 7 [#]	Aug. 3
1947	" "	15.5.46	" 18	Red	23.4.45	Jul. 22
	# Second sowing; first sowing		failed		2.5.46	Jun. 17
Wheat						
1939	Yeoman	22.10.38	Aug. 12	Yeoman	31.10.38	Aug. 14
1940	"	21.10.39	" 6	"	1.11.39	" 12
1941	"	29.10.40	" 22	"	28.10.40	" 14
1942	"	27.10.41	" 12	"	21.11.41	" 19
1943	"	10.10.42	" 2	Eaten by rats		
1944	"	4.10.43	" 3	Yeoman	11.10.43	Aug. 10
1945	"	26.10.44	" 6	Crop failed		
1946	"	17.10.45	" 9	Yeoman	15.10.45	Aug. 21
1947	"	18.10.46	" 5	Squareheads	5.11.46	" 6
				Master		
Potatoes						
1939	Ally	Apr. 17	Sep. 18	Ally	Apr. 12	Sep. 8
1940	"	" 26	" 23	"	Mar. 30	Sep. 17
1941	"	" 29	Oct. 2	"	Apr. 17	Oct. 3
1942	Majestic	" 21	" 2	Majestic	" 16	" 1
1943	"	" 9	Sep. 17	"	" 12	Sep. 20
1944	"	Mar. 28	" 29	"	May 5	Oct. 3
1945	"	" 29	" 22	"	Apr. 24	" 4
1946	"	" 29	Oct. 2	"	" 24	" 3
1947	"	May 2	Sep. 29	"	May 12	" 2
Rye						
1939		8.11.38	Aug. 12		31.10.38	Aug. 11
1940		23.10.39	" 3		1.11.39	" 2
1941		29.10.40	" 14		22.10.40	" 7
1942		28.10.41	" 4		21.11.41	" 1
1943		13.10.42	Jul. 24		10.11.42	Jul. 29
1944		2.10.43	" 22		12.10.43	" 30
1945		2.11.44	" 30		10.11.44	Aug. 3
1946		16.10.45	Aug. 1		5.10.45	" 19
1947		18.10.46	Jul. 29		6.11.46	Jul. 29

Six Course Rotation Experiment

Ba/4.3

Mean yields and increments in yield per cwt. of N, P₂O₅ and K₂O

	Sugar Beet							
	ROTHAMSTED			WOBURN				
	Mean Yield	N	P ₂ O ₅	K ₂ O	Mean Yield	N	P ₂ O ₅	K ₂ O
<u>Roots (Washed): tons per acre</u>								
1939	9.98	2.16	-1.07	2.03	8.76	3.75	-1.71	0.44
1940	8.25	3.64	-2.67	-0.24	5.84	2.40	-0.39	0.79
1941	9.30	4.17	0.47	0.83	7.87	3.11	-1.19	0.81
1942	12.48	0.76	-0.35	0.39	6.78	-0.24	-5.33	1.53
1943	12.21	7.36	-1.29	-1.05	7.54	9.52	-0.98	-1.52
1944	6.45	-1.25	2.75	2.26	11.56	4.53	1.60	-1.24
1945	12.44	5.06	-2.01	0.38	12.75	6.47	-2.47	3.12
1946	11.18	3.71	-0.99	-0.16	5.11	2.97	-2.83	0.33
1947	6.80	-1.88	2.55	0.62	8.06	-1.23	1.93	0.26
<u>Sugar Percentage</u>								
1939	17.86	1.00	-0.20	1.33	18.25	10.05	0.18	-0.10
1940	19.25	-0.25	0.75	0.04	18.66	-1.52	-0.13	0.07
1941	19.52	-0.01	0.81	0.77	18.35	-0.71	-0.52	0.20
1942	17.92	-2.87	1.46	0.90	17.88	-0.59	-0.97	-0.04
1943	18.56	-1.16	0.45	-0.03	17.67	-1.27	-0.40	-0.44
1944	16.58	-1.79	2.63	1.32	17.76	-0.35	-0.09	0.26
1945	19.24	1.07	-0.27	0.42	17.98	-0.41	0.57	0.56
1946	18.10	-0.85	-0.16	0.36	19.50	0.11	0.82	0.80
1947	19.22	-0.65	-1.76	0.26	20.06	0.10	0.53	1.01
<u>Total Sugar: cwt. per acre</u>								
1939	35.7	9.6	-4.1	9.9	32.0	13.5	-5.9	1.4
1940	31.8	13.3	-8.9	-0.9	21.7	7.0	-1.6	3.0
1941	36.3	16.3	3.3	4.8	28.9	10.3	-5.2	3.3
1942	44.7	-4.4	2.6	3.5	24.2	-1.5	-20.0	5.2
1943	45.3	24.4	-4.0	-4.0	26.6	31.5	-4.1	-6.0
1944	21.4	-6.3	12.5	9.4	41.1	15.2	5.6	-3.9
1945	47.9	21.6	-8.5	2.6	45.9	22.0	-7.6	12.4
1946	40.4	11.4	-4.0	0.2	19.9	11.7	-10.3	2.2
1947	26.2	-8.4	7.4	2.8	32.3	-4.7	8.6	2.6
<u>Tops: tons per acre</u>								
1939	11.01	3.40	-1.00	-0.32	5.73	2.42	-0.52	-0.62
1940	4.81	2.25	-0.98	-0.25	2.93	0.98	-0.65	0.42
1941	8.92	8.18	-0.22	-0.86	6.03	4.47	-1.41	0.52
1942	12.73	9.99	-3.08	-1.28	2.78	-0.51	-1.66	0.49
1943	6.94	8.75	1.82	-0.93	3.62	3.71	1.84	-0.28
1944	9.31	3.95	0.35	1.84	5.95	1.37	3.61	-2.70
1945	10.46	7.20	1.22	1.06	4.82	3.49	-0.73	1.94
1946	9.98	8.97	2.33	0.10	3.27	2.80	-1.65	-0.46
1947	4.28	1.37	1.63	0.04	3.56	0.93	1.00	0.21

Ba/4.4

Mean yields and increments in yield per cwt. of N, P₂O₅ and K₂O

ROTHAMSTED

WOBURN

	Mean Yield	N	P ₂ O ₅	K ₂ O	Mean Yield	N	P ₂ O ₅	K ₂ O
<u>Barley Grain: cwt. per acre</u>								
1939	38.3	18.5	7.1	-1.9	21.6	8.1	-6.7	7.1
1940	30.4	19.6	-3.9	-0.2	22.3	13.9	9.9	-0.3
1941	28.5	11.7	6.0	-1.6	16.4	5.4	2.1	-4.9
1942	31.0	21.3	-3.7	1.4	25.0	22.4	8.5	-1.0
1943	32.4	10.6	2.3	0.6	15.1	15.7	0.5	2.2
1944	30.8	13.7	3.8	-0.9	19.4	9.5	-5.4	7.1
1945	30.7	10.9	2.7	2.2	16.0	12.7	-3.5	-5.4
1946	33.0	7.1	-6.3	0.4	24.0	8.9	-1.1	-1.3
1947	27.5	14.7	-5.2	-3.8	18.4	26.7	5.1	-1.0
<u>Barley Straw: cwt. per acre</u>								
1939	41.8	17.1	9.5	0.6	32.7	20.1	-3.6	-8.4
1940	31.6	24.5	2.7	0.6	21.7	33.9	8.5	0.1
1941	30.4	16.0	10.9	-2.6	21.7	8.7	-2.8	-0.8
1942	31.8	26.9	-2.5	4.4	34.0	38.5	6.0	-3.6
1943	33.6	17.4	1.7	-1.0	30.7	31.7	-4.9	8.4
1944	29.7	20.2	8.0	-0.9	19.0	16.4	-6.1	4.8
1945	33.1	6.6	0.7	0.3	21.0	20.5	1.3	-4.1
1946	43.5	22.6	-7.3	0.5	24.6	15.4	-2.9	-4.4
1947	26.1	19.9	-4.3	-2.0	16.9	19.9	-2.7	-5.2
<u>Clover Hay. Dry matter: cwt. per acre</u>								
1939	22.3	9.4	4.7	2.9	25.1	-12.3	13.7	-1.7
1940	39.6	9.5	5.1	2.3	42.5	1.5	4.8	4.1
1941	19.8	6.4	-0.3	-0.4	21.0	-7.7	3.6	-4.5
1942	23.5	4.7	-4.3	-4.7	21.6	-7.4	11.5	3.7
1943	40.4	9.3	-14.5	6.6	15.6	20.7	-2.9	-4.4
1944	15.6	6.2	0.3	6.2	6.6	11.6	-3.9	1.8
1945	36.3	4.9	-1.4	-0.9	27.5	12.5	1.3	5.9
1946	31.9	21.3	13.8	-6.6	55.0	4.7	-8.7	-2.3
1947	32.0	1.8	-2.5	0.7	23.9	12.1	-2.9	1.7
<u>Potatoes: Total tubers, tons per acre</u>								
1939	8.10	1.31	2.60	2.35	6.86	6.39	0.57	-0.03
1940	6.72	1.38	3.01	2.12	7.07	5.20	2.68	1.00
1941	9.24	5.79	0.63	0.64	8.52	9.63	-4.29	-0.83
1942	9.47	1.95	-0.33	3.71	8.46	4.72	4.03	-0.25
1943	8.50	4.29	0.15	2.50	5.90	3.55	1.07	-1.15
1944	10.07	6.38	2.05	0.84	9.98	2.07	4.97	-2.10
1945	9.55	3.79	-0.95	3.59	10.61	5.46	-2.05	0.90
1946	10.75	7.97	-1.73	2.49	6.60	6.35	-1.79	-2.51
1947	5.69	1.52	1.61	1.29	6.51	1.42	-1.02	2.90

Six Course Rotation Experiment

ROTHAMSTED					WOBURN				
	Mean Yield	N	P ₂ O ₅	K ₂ O	Mean Yield	N	P ₂ O ₅	K ₂ O	
<u>Wheat.</u> <u>Grain:</u> <u>cwt. per acre</u>									
1939	27.7	0.7	-5.8	0.2	23.3	-1.1	9.7	-3.1	
1940	27.7	11.7	-1.5	-1.6	18.7	16.6	-7.1	0.2	
1941	19.3	6.1	-10.2	-4.3	19.5	11.4	10.7	1.4	
1942	37.2	10.5	2.8	-1.4	10.8	10.5	13.4	-3.6	
1943	36.8	3.9	-0.5	3.8	(Crop eaten by rats in barn)				
1944	41.1	11.9	-1.5	-1.6	9.9	6.3	0.6	1.2	
1945	25.8	12.3	2.1	1.2	(Crop failed)				
1946	29.2	13.8	12.7	-2.4	13.4	16.5	1.7	2.7	
1947	25.1	10.1	-0.4	0.5	9.5	8.2	-7.3	5.3	
<u>Wheat Straw:</u> <u>cwt. per acre</u>									
1939	48.0	22.7	-1.1	-0.3	35.5	8.7	20.2	-6.3	
1940	34.1	23.9	-1.9	-1.6	32.3	28.5	-5.7	-2.2	
1941	24.3	9.5	-11.3	-3.3	31.3	24.2	17.6	3.3	
1942	44.8	14.8	3.5	-1.0	20.7	13.9	11.6	0.0	
1943	49.0	5.1	-1.7	3.9	(Crop eaten by rats in barn)				
1944	52.8	15.4	-14.1	-1.5	29.7	10.5	-3.2	6.5	
1945	36.8	22.9	0.7	1.8	(Crop failed)				
1946	47.3	19.3	18.7	-3.5	22.4	25.9	-2.1	6.6	
1947	27.5	15.0	0.9	-1.6	15.4	14.8	-7.3	6.5	
<u>Rye Grain:</u> <u>cwt. per acre</u>									
1939	22.1	19.9	1.2	0.0	15.7	11.2	0.5	0.7	
1940	26.8	11.1	-0.3	0.1	20.3	13.6	1.7	-0.3	
1941	16.7	10.5	0.3	-4.0	18.9	5.1	-3.7	1.0	
1942	27.8	24.4	-0.3	-2.0	23.7	33.8	-4.5	-3.6	
1943	32.2	27.5	-7.1	1.5	8.7	13.3	2.7	0.0	
1944	31.0	15.9	-1.5	-2.6	17.1	17.6	3.1	0.4	
1945	25.7	12.9	-0.9	2.3	14.2	11.5	-1.1	-1.7	
1946	29.4	10.4	-3.1	0.4	14.8	9.1	-2.9	-3.5	
1947	15.8	9.8	1.5	1.2	14.9	21.3	2.7	-2.0	
<u>Rye Straw:</u> <u>cwt. per acre</u>									
1939	42.8	32.9	7.7	-2.1	33.3	28.6	-6.1	3.5	
1940	36.7	19.5	-0.6	-1.3	37.7	30.3	-1.9	3.2	
1941	25.3	17.1	2.2	-1.9	36.3	15.1	-0.5	5.0	
1942	36.5	24.3	2.3	1.6	42.2	53.2	-6.3	-5.9	
1943	47.0	32.7	-9.9	2.9	25.5	28.9	7.4	-6.8	
1944	62.3	11.6	2.8	-3.0	37.8	31.3	2.3	0.7	
1945	32.7	14.7	0.7	1.6	29.1	26.5	5.1	-0.2	
1946	55.7	17.5	-5.9	-2.5	26.3	16.3	1.7	-3.3	
1947	21.8	11.4	1.6	-2.4	18.3	26.9	-8.9	3.3	

LONG PERIOD CULTIVATION EXPERIMENT

Long Hoos V, 1934-39

This experiment was completed in 1939. The results were reported by E.W.Russell and B.A.Keen in "Studies in Soil Cultivation - X. The results of a six-year cultivation experiment". J.Agric.Sci., 31, (1941), 326.

The experiment was designed to compare the effects on crop yield and weed infestation of continued ploughing, rotary cultivation with the Simar implement and stirring the soil with a cultivator, and also to investigate the value of cyanamide as a weed-killer. The crops grown in rotation were wheat, mangolds, barley, there being four blocks of twelve plots each under each crop. The twelve treatments were made up of all combinations of:-

Ploughing (P), Simaring (S) or Cultivating (C)
 Shallow cultivation (4 ins.) (Sh) or deep cultivation (8 ins.) (D)
 Cyanamide (Cy) or Nitrochalk (N)

On two of the four blocks the treatments were continued on the same plots in successive years, while on the other two blocks cycles of cultivations and manuring were used.

Area of each plot, 0.015 acre.

A basal dressing of 0.75 cwt. P₂O₅ per acre (as superphosphate) and 1.0 cwt. K₂O per acre (as muriate of potash) was given to all plots growing mangolds.

Details of the experiment are as given in the 1934 Report, p. 175.

Crop Notes

	Wheat	Mangolds	Barley
Variety	Wilhelmina	Yellow Globe	Plumage Archer
Previous crop	Barley	Wheat	Mangolds
Date sown	24/10/38	April 25	March 20
Harvested	Aug. 18	Oct. 25	Aug. 29

Standard errors per plot:

Wheat Grain, 0.950 cwt. per acre or 3.6%, 11 d.f.
 Mangolds Roots, 1.539 tons per acre or 6.6%, 11 d.f.
 Barley Grain, 1.313 cwt. per acre or 6.9%, 11 d.f.

1939

	Continuous			Cycle A			Cycle B			Mean		
	1938 P	1938 S	1938 C	1939 Mean	1939 P	1939 S	1939 C	1939 P	1939 S			
Wheat Grain: cwt. per acre												
N D	31.3	25.0	24.3	±0.67	26.9	22.5	27.3	21.4	29.3	26.6	22.8	±0.39
Sh	28.6	26.5	22.8	28.6	26.0	27.6	26.6	24.7	27.5	25.0	22.3	25.6
Cy D	29.8	23.8	23.6	29.8	25.7	27.6	25.9	26.6	29.5	23.7	25.8	26.5
Sh	25.4	24.5	26.6	25.4	25.5	29.3	26.2	24.6	23.7	21.0	23.9	24.8
Wheat Straw: cwt. per acre												
N D	45.4	36.5	34.0	38.6	52.9	51.6	41.9	46.7	40.7	34.6	44.7	
Sh	42.5	39.4	34.0	38.6	43.8	44.7	43.8	42.1	39.5	33.4	41.2	
Cy D	46.3	36.6	35.0	39.3	41.5	45.4	40.7	44.8	34.3	38.0	40.8	
Sh	43.7	35.6	40.7	40.0	45.0	39.3	42.7	34.3	31.3	38.8	38.6	
Mangolds Roots: tons per acre												
N D	24.98	22.29	23.56	±1.09	23.61	25.30	25.19	23.85	28.26	23.21	27.16	±0.63
Sh	26.20	23.21	21.88	26.20	23.76	26.46	23.50	24.72	24.20	21.94	26.99	24.64
Cy D	24.46	23.30	23.07	24.46	23.61	28.03	23.74	23.91	29.48	25.30	22.63	25.52
Sh	24.93	21.13	19.01	24.93	21.69	24.72	26.99	24.96	22.87	21.12	24.61	24.21
Mangolds Tops: tons per acre												
N D	5.72	5.37	6.15	5.72	6.27	5.57	6.04	5.80	5.28	6.62	5.93	
Sh	5.86	5.98	5.69	5.86	5.75	6.04	5.80	5.57	4.87	6.44	5.74	
Cy D	6.09	5.92	5.69	5.90	5.75	5.80	5.80	7.08	5.69	5.11	5.87	
Sh	6.56	5.51	5.40	5.82	5.46	6.04	5.98	5.80	4.99	5.57	5.64	
Barley Grain: cwt. per acre												
N D	19.8	22.6	17.1	±0.93	19.8	23.6	21.5	19.2	23.6	22.6	18.8	±0.54
Sh	22.6	18.6	15.7	22.6	19.0	22.1	24.1	19.9	24.7	16.4	20.4	21.3
Cy D	23.6	20.6	16.4	23.6	20.2	23.1	21.7	20.5	25.2	20.1	21.7	22.0
Sh	21.5	16.2	14.1	21.5	17.3	22.5	19.2	16.8	19.3	18.0	20.8	19.4
Barley Straw: cwt. per acre												
N D	24.3	26.2	22.0	24.2	25.1	23.7	23.8	28.6	25.0	24.1	25.0	
Sh	27.2	23.2	16.8	22.4	22.1	25.8	27.7	28.7	21.9	23.7	25.0	
Cy D	25.7	24.6	21.6	24.0	27.4	24.1	25.3	28.1	22.8	25.3	25.5	
Sh	25.2	20.0	17.5	20.9	25.1	22.6	21.5	24.8	18.6	22.8	22.6	

Standard errors are based on 11 d.f.

Long Period Cultivation ExperimentMean of Nitrochalk and Cyanamide
1939

	Continuous			Cycle A			Cycle B			Mean
	1938 P	S	C	1939 P	S	C	1938 S	C	P	
Wheat Grain: cwt. per acre										
D	30.6	24.4	24.0	26.3	25.0	26.6	24.0	29.4	25.2	24.3
Sh	27.0	25.5	24.7	25.7	28.4	26.4	24.6	25.6	23.0	23.1
Mean	28.8	25.0	24.4	26.1	26.7	26.5	24.3	27.5	24.1	23.7
	± 0.34		± 0.48		± 0.48		± 0.48		± 0.48	
Wheat Straw: cwt. per acre										
D	45.8	36.6	34.5	39.0	47.2	48.5	41.3	45.8	37.5	36.3
Sh	43.1	37.5	37.4	39.3	44.4	42.0	43.2	38.2	35.4	36.1
Mean	44.4	37.0	36.0	39.2	45.8	45.2	42.2	42.0	36.4	36.2
Mangolds: Roots: tons per acre										
D	24.72	22.80	23.32	23.61	26.66	24.46	23.88	28.87	24.26	24.90
Sh	25.56	22.17	20.44	22.72	25.59	25.24	24.84	23.54	21.53	25.80
Mean	25.14	22.48	21.88	23.17	26.12	24.85	24.36	26.20	22.90	25.35
	± 0.54		± 0.77		± 0.77		± 0.77		± 0.77	
Mangolds Tops: tons per acre										
D	5.90	5.64	5.92	5.82	6.01	5.68	5.92	6.44	5.48	5.86
Sh	6.21	5.74	5.54	5.83	5.60	6.04	5.89	5.68	4.93	6.00
Mean	6.06	5.69	5.73	5.83	5.80	5.86	5.90	6.06	5.20	5.93
Barley Grain: cwt. per acre										
D	21.7	21.6	16.8	20.0	23.4	21.6	19.8	24.4	21.4	20.2
Sh	22.0	17.4	14.9	18.1	22.3	21.6	18.4	22.0	17.2	20.6
Mean	21.8	19.5	15.8	19.0	22.8	21.6	19.1	23.2	19.3	20.4
	± 0.46		± 0.66		± 0.66		± 0.66		± 0.66	
Barley Straw: cwt. per acre										
D	25.0	25.4	21.8	24.1	26.2	23.9	24.6	28.4	23.9	24.7
Sh	26.2	21.6	17.2	21.7	23.6	24.2	24.6	26.8	20.2	23.2
Mean	25.6	23.5	19.5	22.9	24.9	24.0	24.6	27.6	22.0	24.0
	± 0.46		± 0.66		± 0.66		± 0.66		± 0.66	

Standard errors are based on 11 d.f.

DEEP CULTIVATION ROTATION EXPERIMENT
Long Hoos I and II (begun in 1944)

- (1) To compare deep ploughing (about 12 inches) with shallow ploughing (about 6 inches).
- (2) A test of dung ploughed in at different depths.
- (3) To compare ploughing in mineral fertilizers (superphosphate and muriate of potash) at different depths with applying them in the seed bed.

The crops of the six course rotation are sugar beet, barley, one year seeds, wheat, potatoes, oats. There are six series, one for each crop of the rotation. Each series has 16 main plots for the ploughing, dung, P and K treatments. The main plots are divided into two to test the depth of application of mineral fertilizers to sugar beet and potatoes. Each plot receives the same treatment throughout the experiment.

System of replication: 2 blocks of 8 plots each in each series, the four-factor interaction of main plot treatments being confounded with block differences.

Area of each whole plot: 0.0132 acre, before rejecting edge rows.

Treatments

Whole plots: All combinations of:-

- (1) Shallow (6") v. Deep (12") ploughing. Ploughing treatments on stubbles in Autumn for sugar beet and potatoes, on seeds aftermath for wheat.

	Sugar Beet	Potatoes
(2) No dung v. dung ploughed in	10 tons	20 tons
(3) No phosphate v. superphosphate	0.6 cwt. P_2O_5	0.8 cwt. P_2O_5
(4) No potash v. muriate of potash	0.6 cwt. K_2O	1.0 cwt. K_2O

(all amounts per acre)

Half plots (Sugar beet and potatoes only):-

Minerals (PK combinations) ploughed in v. minerals in seed bed.

Basal manuring: Applied in the ridges for potatoes, as a top dressing to wheat and in the seed bed for other crops.

Basal Manuring (continued)

Bc/1.2
Spring
Oats

	Sugar beet	Barley	Ley	Wheat	Potatoes	
Sulphate of Ammonia (cwt. N per acre)	0.8	0.3	-	0.5	0.6	0.2
Basic Slag (cwt. P ₂ O ₅ per acre)	-	0.6	-	-	-	-

Non-experimental Cultivations: These are carried out over the whole of any series, with the proviso that they must not be deeper than 6 inches except that deep ploughed plots may be worked to a depth of below 6 inches for the root crops.

Series	Preliminary years		Years of complete cropping	
	1944	1945	1946	1947
1	Potatoes	Spring oats	Sugar beet	Barley
2	Sugar beet	Barley	Ley	Wheat
3	(Barley)	(Ley)	Wheat	Potatoes
4	(Wheat)	(Barley)	Potatoes	Spring oats
5	(Wheat)	Potatoes	Spring oats	Sugar beet
6	(Wheat)	Sugar beet	Barley	Ley

(Non-experimental crops in brackets)

The Physics Department has made observations on crop development from 1944 onwards.

Crop Notes

	Sugar beet	Barley	Ley	Wheat	Potatoes	Spring oats
1944						
Sown	Apr. 20				Apr. 5	
Harvested	Oct. 19				Sep. 27	
1945						
Sown	Apr. 13	Mar. 26			Apr. 4	Mar. 22
Harvested	Oct. 22	Aug. 11			Sep. 25	Aug. 16
1946						
Sown	Apr. 13	Mar. 21	27.3.45	Oct. 16	May 1	Mar. 21
Harvested	Oct. 24	Aug. 13	Jun. 24	Aug. 12	Oct. 1	Aug. 6
1947						
Sown	Apr. 25	Apr. 12	16.4.46	Oct. 18	May 2	Apr. 11
Harvested	Nov. 1	Aug. 11	Jun. 10	Aug. 1	Sep. 30	Aug. 5
Variety	Klein E	Plumage Archer	Mixture given below	Yeoman	Majestic	Star

Seeds mixture (all amounts per acre): 18 lb. Perennial ryegrass, 6 lb. late flowering red clover, 2 lb. Alsike clover.

Sugar Beet

Roots (washed): tons per acre

Standard errors

	Mean yield	per whole plot		per split plot	
1944	10.83	0.890 or 8.2, .9	d.f.	1.70 or 15.3%, 7	d.f.
1945	15.20	1.03 or 6.8%, 6	d.f.	0.781 or 5.1%, 7	d.f.
1946	16.21	1.19 or 7.4%, 10	d.f.	1.61 or 9.7%, 7	d.f.
1947	10.08	0.658 or 6.5%, 4	d.f.	0.418 or 4.1%, 7	d.f.

Responses to treatments

	Mean	Ploughing	Dung	Phosphate	Potash
		Shallow Deep	Abs. Pres.	Abs. Pres.	Abs. Pres.
<u>1944</u>				±0.63	
Ploughing	±0.44				
deep-					
shallow	0.39	- -	0.45 0.33	0.59 0.19	0.90 -0.12
Dung	1.00	1.06 0.94	- -	0.68 1.32	1.48 0.52
Phosphate	0.51	0.71 0.31	0.19 0.83	- -	0.47 0.55
Potash	0.98	1.49 0.47	1.46 0.50	0.94 1.02	- -
<u>1945</u>				±0.73	
Ploughing	±0.52				
deep-					
shallow	0.53	- -	0.80 0.26	1.04 0.02	0.65 0.41
Dung	1.00	1.27 0.73	- -	1.23 0.77	1.50 0.50
Phosphate	0.61	1.12 0.10	0.84 0.38	- -	0.34 0.88
Potash	0.21	0.33 0.09	0.71 -0.29	-0.06 0.48	- -
<u>1946</u>				±0.85	
Ploughing	±0.60				
deep-					
shallow	0.64	- -	0.72 0.56	0.13 1.15	0.73 0.55
Dung	0.91	0.99 0.83	- -	2.06 -0.24	1.49 0.33
Phosphate	0.44	-0.07 0.95	1.59 -0.71	- -	0.13 0.75
Potash	0.79	0.88 0.70	1.37 0.21	0.48 1.10	- -
<u>1947</u>				±0.47	
Ploughing	±0.33				
deep-					
shallow	3.24	- -	3.20 3.28	3.45 3.03	3.51 3.17
Dung	2.23	2.19 2.27	- -	2.19 2.27	2.65 1.81
Phosphate	0.01	0.22 -0.20	-0.03 0.05	- -	0.65 -0.63
Potash	0.20	0.27 0.13	0.62 -0.22	0.84 -0.44	- -

Deep Cultivation Rotation Experiment

Ploughing: The plough used for deep cultivations was a Ransome Solotrac giving a depth of 12 inches at least, except in 1944 when a Massey Harris Grub Breaker was used. This could not be made to cut a 12 inch deep furrow at all satisfactorily, the actual depth in that year being 9-12 inches. Until 1947 the whole of the seeds area was ploughed 6" deep after the hay was carted, the deep ploughing being carried out subsequently at the same time as the stubbles were deep ploughed for roots. In autumn 1946 the second ploughing of the seeds could not be carried out owing to wet conditions, so there was no test of deep ploughing on wheat in 1947. In summer 1947 the deep and shallow ploughing treatments were carried out directly on the hay stubble and this will be continued.

Deep Cultivation Rotation Experiment

Sugar Beet

Roots (washed): tons per acre

Time of application of mineral fertilizers

		Superphosphate			Potash			Mean
		None	Ploughed in	In s e d b e d	None	Ploughed in	In s e d b e d	
	Standard errors: (a)		(b) & (c)		(a)		(b) & (c)	
<u>1944</u>	Shallow	10.28	10.73	11.25	9.89	11.37	11.41	10.64
	Deep	10.87	11.97	10.39	10.79	12.05	10.46	11.03
	No dung	16.23	10.20	10.61	9.60	11.22	10.88	10.32
	Dung	10.92	12.50	10.99	11.08	12.16	10.99	11.33
	Mean	10.58	11.35	10.81	10.34	11.70	10.94	10.83
<u>1945</u>	Shallow	14.37	15.54	15.43	14.76	15.28	14.91	14.93
	Deep	15.41	15.89	15.13	15.42	15.38	15.64	15.46
	No Dung	14.28	15.35	14.89	14.34	15.13	14.98	14.70
	Dung	15.50	16.09	15.68	15.84	15.53	15.57	15.69
	Mean	14.89	15.72	15.28	15.09	15.33	15.28	15.21
<u>1946</u>	Shallow	15.93	15.93	15.78	15.45	16.46	16.20	15.89
	Deep	16.05	17.03	16.97	16.18	16.51	17.23	16.53
	No dung	14.96	17.12	15.99	15.07	16.14	16.74	15.75
	Dung	17.02	15.85	16.77	16.56	16.83	16.69	16.66
	Mean	15.99	16.48	16.38	15.82	16.49	16.72	16.21
<u>1947</u>	Shallow	8.35	8.86	8.27	8.32	9.17	8.01	8.46
	Deep	11.79	11.50	11.70	11.63	12.04	11.48	11.69
	No dung	8.97	9.12	8.78	8.65	9.99	8.55	8.96
	Dung	11.16	11.24	11.19	11.30	11.23	10.93	11.19
	Mean	10.07	10.18	9.98	9.97	10.61	9.74	10.08
	Standard errors:		(a)		(b)		(c)	
	1944		0.44		0.85		0.74	
	1945		0.52		0.39		0.58	
	1946		0.60		0.80		0.83	
	1947		0.33		0.21		0.36	

Standard errors (b) are for use in horizontal comparisons only, (a) and (c) for all other comparisons.

N

Sugar Beet
Sugar Percentage

	Mean	Standard errors		
		per whole plot	per split plot	
1944	15.73	0.371, 9 d.f.	0.398, 7 d.f.	
1945	19.09	0.728, 7 d.f.	1.12, 7 d.f.	
1946	15.59	0.480, 10 d.f.	0.500, 7 d.f.	

		Responses to treatments							
		Ploughing		Dung		Phosphate		Potash	
		Shallow	Deep	Abs.	Pres.	Aba.	Pres.	Abs.	Pres.
<u>1944</u>									
Ploughing	±0.19					±0.26			
deep shallow	0.05	-	-	-0.24	0.34	-0.11	0.21	0.11	-0.01
Dung	-0.25	-0.54	0.04	-	-	-0.22	-0.28	-0.23	-0.27
Phosphate	0.13	-0.03	0.29	0.16	0.10	-	-	0.28	-0.02
Potash	0.38	0.44	0.32	0.40	0.36	0.53	0.23	-	-
<u>1945</u>						±0.51			
Ploughing	±0.36								
deep-shallow	0.34	-	-	-0.17	0.85	0.18	0.50	0.40	0.28
Dung	-0.19	-0.70	0.32	-	-	-0.25	-0.13	0.02	-0.40
Phosphate	-0.31	-0.47	-0.15	-0.37	-0.25	-	-	-0.45	-0.17
Potash	0.39	0.45	0.33	0.60	0.18	0.25	0.53	-	-
<u>1946</u>						±0.34			
Ploughing	±0.24								
deep shallow	0.16	-	-	-0.11	0.43	0.16	0.16	-0.24	0.56
Dung	-0.28	-0.55	-0.01	-	-	0.08	-0.64	-0.06	-0.50
Phosphate	0.33	0.33	0.33	0.69	-0.03	-	-	0.20	0.46
Potash	0.26	-0.14	0.66	0.48	0.04	0.13	0.39	-	-
<u>1947</u>									
Ploughing									
deep shallow	-0.31	-	-	-0.24	-0.38	-0.53	-0.09	-0.15	-0.47
Dung	-0.01	0.06	-0.08	-	-	-0.10	0.08	-0.23	0.21
Phosphate	-0.01	-0.23	0.21	-0.10	0.08	-	-	-0.10	0.08
Potash	0.32	0.48	0.16	0.10	0.54	0.23	0.41	-	-

Deep Cultivation Rotation ExperimentSugar Beet
Sugar percentage

Time of application of mineral fertilizers

	Superphosphate			Potash			Mean
	None	Ploughed	In seed bed	None	Ploughed	In seed bed	
<u>Standard errors:</u>	(a)	(b) &	(c)	(a)	(b) &	(c)	
<u>1944</u>							
Shallow	15.72	15.70	15.67	15.48	15.93	15.92	15.70
Deep	15.61	15.88	15.92	15.60	15.78	16.04	15.76
No dung	15.77	15.92	15.94	15.65	16.07	16.03	15.85
Dung	15.55	15.67	15.64	15.42	15.64	15.93	15.60
Mean	15.66	15.79	15.79	15.54	15.86	15.98	15.73
<u>1945</u>							
Shallow	19.16	18.62	18.76	18.70	18.67	19.62	18.92
Deep	19.34	19.39	18.97	19.10	19.77	19.08	19.26
No dung	19.38	18.87	19.13	18.89	19.01	19.96	19.19
Dung	19.12	19.14	18.60	18.90	19.43	18.74	19.00
Mean	19.25	19.01	18.86	18.90	19.22	19.35	19.09
<u>1946</u>							
Shallow	15.34	15.50	15.84	15.58	15.52	15.35	15.51
Deep	15.51	16.06	15.60	15.34	15.99	16.00	15.67
No dung	15.38	16.04	16.10	15.49	16.05	15.88	15.73
Dung	15.47	15.52	15.34	15.43	15.46	15.47	15.45
Mean	15.42	15.78	15.72	15.46	15.76	15.68	15.59
<u>1947</u>							
Shallow	20.47	20.03	20.45	20.11	20.38	20.82	20.36
Deep	19.94	20.14	20.17	19.97	20.10	20.16	20.05
No dung	20.26	20.14	20.17	20.16	20.07	20.44	20.21
Dung	20.16	20.02	20.45	19.92	20.40	20.54	20.20
Mean	20.21	20.08	20.31	20.04	20.24	20.49	20.20

Standard errors:	(a)	(b)	(c)
1944	±0.19	±0.20	±0.23
1945	±0.36	±0.56	±0.54
1946	±0.24	±0.25	±0.30

Standard errors (b) are for use in horizontal comparisons only, (a) and (c) for all other comparisons.

Sugar Beet

Total sugar: cwt. per acre

Standard errors

	Mean yield	per whole plot	per sub-plot
1944	34.1	3.32 or 9.8%, 9 d.f.	6.23 or 17.7%, 7 d.f.
1945	58.0	3.87 or 6.7%, 8 d.f.	4.37 or 7.5%, 7 d.f.
1946	50.5	4.54 or 9.0%, 10 d.f.	5.54 or 10.8%, 7 d.f.
1947	40.7	2.37 or 5.8%, 4 d.f.	1.26 or 3.1%, 7 d.f.

Responses to treatments

	Mean	Ploughing		Dung		Phosphate		Potash	
		Shallow	Deep	Absent	Present	Absent	Present	Absent	Present
<u>1944</u>	<u>±1.66</u>					<u>±2.35</u>			
Ploughing		-	-	1.0	1.8	1.6	1.2	3.1	-0.3
deep-shallow	1.4	-	-						
Dung	2.6	2.2	3.0	-	-	1.7	3.5	4.2	1.0
Phosphate	2.0	2.2	1.8	1.1	2.9	-	-	2.2	1.8
Potash	3.9	5.6	2.2	5.5	2.3	4.1	3.7	-	-
<u>1945</u>	<u>±1.94</u>					<u>±2.74</u>			
Ploughing		-	-	2.7	3.7	4.7	1.7	3.7	2.7
deep-shallow	3.2	-	-						
Dung	3.4	2.9	3.9	-	-	4.2	2.6	5.7	1.1
Phosphate	1.5	3.0	0.0	2.3	0.7	-	-	-0.2	3.2
Potash	1.9	2.4	1.4	4.2	-0.4	0.2	3.6	-	-
<u>1946</u>	<u>±2.27</u>					<u>±3.21</u>			
Ploughing		-	-	2.0	3.0	0.9	4.1	1.4	3.6
deep-shallow	2.5	-	-						
Dung	1.8	1.3	2.3	-	-	6.4	-2.8	4.4	-0.8
Phosphate	2.5	0.9	4.1	7.1	-2.1	-	-	1.0	4.0
Potash	3.3	2.2	4.4	5.9	0.7	1.8	4.8	-	-
<u>1947</u>	<u>±1.19</u>					<u>±1.68</u>			
Ploughing		-	-	12.5	12.3	12.8	12.0	12.9	11.9
deep-shallow	12.4	-	-						
Dung	9.0	9.1	8.9	-	-	8.6	9.4	10.1	7.9
Phosphate	0.1	0.5	-0.3	-0.3	0.5	-	-	2.5	-2.3
Potash	1.4	1.9	0.9	2.5	0.3	3.8	-1.0	-	-

Deep Cultivation Rotation Experiment

Sugar Beet

Total sugar: cwt. per acre

	Time of application of mineral fertilizers						Mean	
	Superphosphate			Potash				
	None	Ploughed in bed	In seed bed	None	Ploughed in bed	In seed bed		
<u>1944</u>	Standard error (a)	(b) and (c)		(a)	(b) and (c)		Mean	
	Shallow	32.3	33.7	35.3	30.6	36.2	33.4	
	Deep	33.9	38.3	33.1	33.7	38.1	34.8	
	No dung	32.3	32.7	34.0	30.1	36.1	32.8	
	Dung	34.0	39.3	34.4	34.3	38.1	35.4	
	Mean	33.1	36.0	34.2	32.2	37.2	34.1	
<u>1945</u>	Shallow	54.8	57.9	57.9	55.2	57.1	56.4	
	Deep	59.6	61.7	57.4	58.9	60.9	59.6	
	No dung	55.1	57.9	57.0	54.2	57.5	56.3	
	Dung	59.3	61.7	58.3	59.9	60.5	59.6	
	Mean	57.2	59.8	57.6	57.0	59.0	58.0	
<u>1946</u>	Shallow	48.8	49.6	49.9	48.1	51.0	49.3	
	Deep	49.7	54.6	53.1	49.6	52.8	51.8	
	No dung	46.1	54.8	51.6	46.7	51.8	49.6	
	Dung	52.5	49.3	51.5	51.1	52.0	51.4	
	Mean	49.3	52.1	51.5	48.9	51.9	50.6	
<u>1947</u>	Shallow	34.2	35.5	33.9	34.5	37.4	34.4	
	Deep	47.0	46.3	47.2	46.4	48.4	46.9	
	No dung	36.3	36.7	35.3	34.9	40.0	36.2	
	Dung	44.9	45.0	45.7	45.0	45.8	45.1	
	Mean	40.6	40.9	40.5	39.9	42.9	40.7	
	Standard errors:			(a)	(b)	(c)		
	1944			1.66	3.12	2.76		
	1945			1.94	2.18	2.48		
	1946			2.27	2.77	3.00		
	1947			1.19	0.63	1.27		

Standard errors (b) are for use in horizontal comparisons only, (a) and (c) for all other comparisons.

X Sugar Beet

Tops: tons per acre

Mean Yield		Standard errors							
		per whole plot				per sub-plot			
1944	16.85	1.32	or 7.9%, 9 d.f.			1.75	or 10.2%, 7 d.f.		
1945	16.05	1.27	or 7.9%, 8 d.f.			1.32	or 8.2%, 7 d.f.		
1946	18.89	1.08	or 5.7%, 7 d.f.			1.37	or 7.3%, 7 d.f.		
1947	7.00	0.281	or 4.0%, 4 d.f.			0.612	or 8.6%, 7 d.f.		

Responses to treatments

	Mean	Ploughing		Dung		Phosphate		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Ploughing	±0.66					±0.93			
deep-shallow	-0.18	-	-	-0.70	0.34	-0.50	0.14	0.03	-0.39
Dung	1.82	1.30	2.34	-	-	1.43	2.21	3.01	0.63
Phosphate	-0.01	-0.33	0.31	-0.40	0.38	-	-	-0.25	0.23
Potash	1.92	2.13	1.71	3.11	0.73	1.68	2.16	-	-
<u>1945</u>						±0.90			
Ploughing	±0.63								
deep-shallow	1.66	-	-	1.20	2.12	1.02	2.30	2.28	1.04
Dung	1.05	0.59	1.51	-	-	1.36	0.74	1.16	0.94
Phosphate	0.44	-0.20	1.08	0.75	0.13	-	-	0.03	0.85
Potash	0.13	0.75	-0.49	0.24	0.02	-0.28	0.54	-	-
<u>1946</u>						±0.76			
Ploughing	±0.54								
deep-shallow	0.04	-	-	1.74	-1.66	0.44	-0.36	1.07	-0.99
Dung	1.60	3.30	-0.10	-	-	0.25	2.95	1.43	1.77
Phosphate	-0.40	0.00	-0.80	-1.75	0.95	-	-	-0.16	-0.64
Potash	0.07	1.10	-0.96	-0.10	0.24	0.31	-0.17	-	-
<u>1947</u>						±0.20			
Ploughing	±0.14								
deep-shallow	1.73	-	-	1.84	1.62	2.21	1.25	1.61	1.85
Dung	1.69	1.80	1.58	-	-	1.90	1.48	2.22	1.16
Phosphate	0.16	0.64	-0.32	0.37	-0.05	-	-	0.68	-0.36
Potash	0.19	0.07	0.31	0.72	-0.34	0.71	-0.33	-	-

(a) *above ground* (b) *below ground* (c) *above ground* (d) *below ground*
(e) *above ground* (f) *below ground* (g) *above ground* (h) *below ground*

Deep Cultivation Rotation Experiment

Bc/1.11

Sugar Beet

Tops: tons per acre

Time of application of mineral fertilizers

	Superphosphate			Potash			Mean
	None	Ploughed in	In seed bed	None	Ploughed in	In seed bed	
<u>Standard errors:</u>	(a)	(b) and (c)		(a)	(b) and (c)		
<u>1944</u>							
Shallow	17.11	16.00	17.56	15.88	17.28	18.74	16.94
Deep	16.61	16.77	17.07	15.91	17.44	17.80	16.76
No dung	16.14	14.94	16.55	14.39	16.68	18.32	15.94
Dung	17.57	17.82	18.08	17.40	18.04	18.22	17.76
Mean	16.86	16.38	17.32	15.90	17.36	18.27	16.85
<u>1945</u>							
Shallow	15.32	15.21	15.03	14.84	15.79	15.39	15.22
Deep	16.34	17.31	17.52	17.12	16.14	17.13	16.88
No dung	15.15	15.99	15.80	15.40	15.46	15.83	15.52
Dung	16.51	16.53	16.75	16.57	16.47	16.70	16.58
Mean	15.83	16.26	16.28	15.98	15.96	16.26	16.05
<u>1946</u>							
Shallow	18.87	18.34	19.38	18.32	19.37	19.45	18.87
Deep	19.30	19.30	17.71	19.38	19.09	17.76	18.90
No dung	18.96	17.61	16.80	18.14	18.73	17.33	18.08
Dung	19.21	20.02	20.30	19.57	19.74	19.87	19.69
Mean	19.09	18.82	18.55	18.85	19.23	18.60	18.88
<u>1947</u>							
Shallow	5.82	6.72	6.20	6.10	6.59	5.76	6.14
Deep	8.03	7.20	8.22	7.72	7.89	8.16	7.87
No dung	5.97	6.04	6.65	5.80	6.77	6.26	6.16
Dung	7.87	7.88	7.77	8.02	7.70	7.66	7.85
Mean	6.92	6.96	7.21	6.91	7.24	6.96	7.00
<u>Standard errors:</u>		(a)		(b)		(c)	
1944		0.66		0.88		0.90	
1945		0.63		0.66		0.79	
1946		0.54		0.68		0.73	
1947		0.14		0.31		0.26	

Standard errors (b) are for use in horizontal comparisons only, (a) and (c) for all other comparisons.

Sugar Beet

Plant number: thousands per acre

Standard errors

	Mean yield	per whole plot				per split plot			
1944	26.4	1.59	or 6.0%	9 d.f.	0.990	or 3.7%	7 d.f.		
1945	26.4	1.55	or 5.9%	9 d.f.	1.25	or 4.7%	7 d.f.		
1946	24.4	0.912	or 3.7%	7 d.f.	0.760	or 3.1%	7 d.f.		
1947	23.1								
		Responses to treatments							
		Ploughing		Dung	Phosphate	Potash	Abs.	Pres.	Abs.
		Shallow	Deep	Abs.	Pres.	Abs.			Pres.
<u>1944</u>	Mean						±1.12		
Ploughing	±0.80								
deep-shallow	1.2	-	-	0.7	1.7	1.0	1.4	2.7	-0.3
Dung	-0.8	-1.3	-0.3	-	-	-1.4	-0.2	-0.9	-0.7
Phosphate	-0.2	-0.4	0.0	-0.3	0.4	-	-	0.2	-0.6
Potash	0.5	2.0	-1.0	0.4	0.6	0.9	0.1	-	-
<u>1945</u>	±0.78						±1.10		
Ploughing									
deep-shallow	0.3	-	-	1.0	-0.4	-0.1	0.7	0.4	0.2
Dung	-0.2	0.5	-0.9	+0.9	-	-0.1	-0.3	-0.4	0.0
Phosphate	1.2	0.8	1.6	1.3	1.1	-	-	0.6	1.8
Potash	-0.4	-0.3	-0.5	-0.6	-0.2	-1.0	0.2	-	-
<u>1946</u>	±0.46						±0.65		
Ploughing									
deep-shallow	-0.2	-	-	0.3	-0.7	-0.4	0.0	-0.6	0.2
Dung	0.2	0.7	-0.3	-	-	1.2	-0.8	-0.7	1.1
Phosphate	-0.1	-0.3	0.1	0.9	-1.1	-	-	-0.2	0.0
Potash	1.0	0.6	1.4	0.1	1.9	0.9	1.1	-	-
<u>1947</u>									
Ploughing									
deep-shallow	0.2	-	-	-0.6	1.0	0.5	-0.1	-1.0	1.4
Dung	1.2	0.4	2.0	-	-	0.9	1.5	1.1	1.3
Phosphate	0.5	0.8	0.2	0.2	0.8	-	-	0.7	0.3
Potash	1.7	0.5	2.9	1.6	1.8	1.9	1.5	-	-

Deep Cultivation Rotation Experiment

Sugar Beet

Plant number: thousands per acre
 Time of application of mineral fertilizers

	Superphosphate			Potash			Mean
	None	Ploughed	In seed bed	None	Ploughed	In seed bed	
Standard errors:	(a)	(b) & (c)		(a)	(b) & (c)		
<u>1944</u>							
Shallow	26.0	25.5	25.6	24.8	26.6	27.0	25.8
Deep	27.0	26.9	27.0	27.4	27.2	25.8	27.0
No dung	27.2	26.3	26.5	26.6	27.1	26.9	26.8
Dung	25.8	26.1	26.2	25.6	26.7	25.9	26.0
Mean	26.5	26.2	26.3	26.1	26.9	26.4	26.4
<u>1945</u>							
Shallow	25.9	26.4	27.0	26.4	25.9	26.4	26.3
Deep	25.7	27.5	27.3	26.8	25.7	26.9	26.6
No dung	25.9	26.9	27.5	26.9	26.2	26.3	26.6
Dung	25.7	27.0	26.8	26.4	25.5	27.0	26.3
Mean	25.8	26.9	27.2	26.6	25.8	26.6	26.4
<u>1946</u>							
Shallow	24.6	24.6	24.0	24.1	25.1	24.5	24.4
Deep	24.2	24.5	24.2	23.5	24.8	25.2	24.3
No dung	23.8	25.2	24.3	24.2	24.7	23.9	24.3
Dung	25.0	24.0	23.9	23.5	25.1	25.8	24.5
Mean	24.4	24.6	24.1	23.8	24.9	24.6	24.4
<u>1947</u>							
Shallow	22.6	23.3	23.5	22.8	23.7	22.7	23.0
Deep	23.1	21.8	24.9	21.8	25.1	24.3	23.2
No dung	22.4	21.5	23.7	21.8	23.7	22.8	22.5
Dung	23.3	23.5	24.7	22.8	25.1	24.1	23.7
Mean	22.9	22.5	24.2	22.3	24.4	23.5	23.1
Standard errors:		(a)		(b)		(c)	
1944		0.80		0.50		0.87	
1945		0.78		0.62		0.90	
1946		0.46		0.38		0.53	

Standard errors (b) are for use in horizontal comparisons only, (a) and (c) for all other comparisons.

Barley - Residual Effects

Grain: cwt. per acre

	Mean yield	Standard error per plot
1945	29.2	1.43 or 4.9%, 7 d.f.
1946	36.4	1.79 or 4.9%, 7 d.f.
1947	31.2	0.903 or 2.9%, 4 d.f.

Responses to treatments

	Mean	Ploughing Shallow	Ploughing Deep	Dung Abs.	Dung Pres.	Phosphate Abs.	Phosphate Pres.	Potash Abs.	Potash Pres.
Grain: cwt. per acre									
<u>1945</u>									
Ploughing	+0.71					+1.01			
deep-shallow	-0.6	-	-	-0.9	-0.3	0.1	-1.3	-1.5	0.3
Dung	1.3	1.0	1.6	-	0.3	0.3	2.3	2.4	0.2
Phosphate	0.8	1.5	0.1	-0.2	1.8	-	-	0.3	1.3
Potash	0.7	-0.2	1.6	1.8	-0.4	0.2	1.2	-	-
<u>1946</u>									
Ploughing	+0.90					+1.27			
deep-shallow	0.4	-	-	0.6	0.2	1.2	-0.4	1.0	-0.2
Dung	1.3	1.5	1.1	-	-	1.0	1.6	2.4	0.2
Phosphate	0.7	1.5	-0.1	0.4	1.0	-	-	0.6	0.8
Potash	0.0	0.6	-0.6	1.1	-1.1	-0.1	0.1	-	-
<u>1947</u>									
Ploughing	+0.45					+0.64			
deep-shallow	2.4	-	-	2.4	2.4	2.3	2.5	2.4	2.4
Dung	0.9	0.9	0.9	-	-	0.3	1.5	0.7	1.1
Phosphate	0.0	-0.1	0.1	-0.6	0.6	-	-	0.4	-0.4
Potash	0.4	0.4	0.4	0.2	0.6	0.8	0.0	-	-

Deep Cultivation Rotation Experiment

Barley - Residual Effects

Straw: cwt. per acre

Mean Yield

Standard error per plot

1945	32.7	2.36 or 7.2%, 7 d.f.
1946	43.4	2.33 or 5.4%, 7 d.f.
1947	31.2	

Mean	Ploughing		Dung		Phosphate		Potash	
	Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
	Straw: cwt. per acre							
Ploughing	+1.18				+1.67			
deep-shallow	-0.7	-	-	-1.2	-0.2	0.4	-1.8	-3.1 1.1.7
Dung	3.3	2.8	3.8	-	-	2.8	3.8	4.3 2.3
Phosphate	-0.1	1.0	-1.2	-0.6	0.4	-	-	-0.6 0.4
Potash	1.7	-0.7	4.1	2.7	0.7	1.2	2.2	- -
<u>1946</u>					+1.64			
Ploughing	+1.16							
deep-shallow	0.8	-	-	3.2	-1.6	2.5	-0.9	2.6 -1.0
Dung	5.1	7.5	2.7	-	-	6.1	4.1	10.4 -0.2
Phosphate	0.9	2.6	-0.8	1.9	-0.1	-	-	0.4 1.4
Potash	1.1	2.9	-0.7	6.4	-4.2	0.6	1.6	- -
<u>1947</u>								
Ploughing								
deep-shallow	2.4	-	-	2.3	2.5	1.8	3.0	3.4 1.4
Dung	1.7	1.6	1.8	-	-	0.8	2.6	1.2 2.2
Phosphate	-0.8	-1.4	-0.2	-1.7	0.1	-	-	-0.4 -1.2
Potash	0.8	1.8	-0.2	0.3	1.3	1.2	0.4	- -

N

Wheat
Straw: cwt. per acre

Hay: cwt. per acre - Residual effects

	1946	1947
Mean yield	82.2	54.2
Standard error per plot	3.54 or 4.3% (6 d.f.)	2.82 or 5.2% (4d.f.)

Responses to treatments

Mean	Ploughing	Dung		Phosphate		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.
<u>1946</u>							
Ploughing	+1.77	-	-	2.9	1.1	+2.519	-3.7
deep-shallow	-0.9	-	-	-2.9	1.1	1.9	-3.7
Dung	3.8	1.8	5.8	-	-	3.4	4.2
Phosphate	1.3	4.1	-1.5	0.9	1.7	-	-
Potash	1.5	0.5	2.5	0.8	2.2	-1.3	4.3
<u>1947</u>							
Ploughing	+1.41	-	-	-1.7	2.7	-1.4	+1.99
deep-shallow	0.5	-	-	-	-	-	2.4
Dung	-1.0	-3.2	1.2	-	-	-1.3	-0.7
Phosphate	0.7	-1.2	2.6	0.4	1.0	-	1.2
Potash	1.7	3.2	0.2	1.6	1.8	2.2	1.2

Potatoes

Ware tubers: tons per acre

Standard errors

	Mean yield	per whole plot	per split plot
1944	10.82	0.863 or 8.0%, 9 d.f.	1.10 or 9.9%, 7 d.f.
1945	11.01	0.917 or 8.3%, 7 d.f.	0.531 or 4.8%, 7 d.f.
1946	11.30	0.640 or 5.7%, 6 d.f.	1.08 or 9.4%, 7 d.f.
1947	5.66	0.538 or 9.5%, 4 d.f.	0.397 or 6.8%, 7 d.f.

Responses to treatments

	Mean	Ploughing	Dung	Phosphate	Potash
		Shallow	Abs. Pres.	Abs. Pres.	Abs. Pres.
<u>1944</u>				±0.61	
Ploughing	±0.43				
deep-shallow	0.17	-	-0.01	0.35 -0.69	1.03 0.26 0.08
Dung	1.27	1.09	1.45	- -	1.29 1.25 1.29 1.25
Phosphate	0.38	-0.48	1.24	0.40 0.36	- - 0.33 0.43
Potash	1.16	1.25	1.07	1.18 1.14	1.11 1.21 - -
<u>1945</u>				±0.65	
Ploughing	±0.46				
deep-shallow	1.44	-	-	1.72 1.16	2.30 0.58 1.70 1.18
Dung	3.29	3.57	3.01	- -	3.18 3.40 3.99 2.59
Phosphate	0.49	1.35	-0.37	0.38 0.60	- - -0.10 1.08
Potash	1.09	1.35	0.83	1.79 0.39	0.50 1.68 - -
<u>1946</u>				±0.45	
Ploughing	±0.32				
deep-shallow	0.86	-	-	0.41 1.31	1.17 0.55 0.27 1.45
Dung	1.46	1.01	1.91	- -	1.64 1.28 1.12 1.80
Phosphate	0.32	0.63	0.01	0.50 0.14	- - -0.23 0.87
Potash	0.69	0.10	1.28	0.35 1.03	0.14 1.24 - -
<u>1947</u>				±0.38	
Ploughing	±0.27				
deep-shallow	0.04	-	-	0.01 0.07	0.34 -0.26 -0.46 0.54
Dung	1.64	1.61	1.67	- -	1.61 1.67 2.41 0.87
Phosphate	0.16	0.46	-0.14	0.13 0.19	- - 0.11 0.21
Potash	1.05	0.55	1.55	1.82 0.28	1.00 1.10 - -

Deep Cultivation Rotation Experiment

Wheat

	Mean yield	Standard error per plot
Grain: cwt. per acre		
1946	29.0	3.56 or 12.3%, 13 d.f.
1947	26.8	2.25 or 8.4%, 14 d.f.
Straw: cwt. per acre		
1946	48.0	7.43 or 15.5%, 13 d.f.
1947	32.3	

Responses to treatments

<u>1946</u>		Grain		Straw	
Response to deep ploughing (deep-shallow)		2.2(± 1.78)		1.1(± 3.71)	
<u>1947</u>		Dung		Phosphate	
Residual Effects	Mean	Abs.	Pres.	Abs.	Pres.
Grain: cwt. per acre					
	± 1.12			± 1.59	
Dung	1.5	-	-	2.1	0.9
Phosphate	-0.9	-0.3	-1.5	-	-
Potash	-0.1	0.1	-0.3	-2.1	1.9
Straw: cwt. per acre					
Dung	3.2	-	-	3.7	2.7
Phosphate	-0.9	-0.4	-1.4	-	-
Potash	-0.5	-0.1	-0.9	-3.5	2.5

There was no test of deep ploughing in this year.

Potatoes

Percentage Ware

	Mean	Standard errors					
		per whole plot	per split plot				
1944	78.1	1.07, 9 d.f.	1.43, 7 d.f.				
1945	96.3	0.404, 5 d.f.	0.310, 7 d.f.				
1946	95.3	1.17, 7 d.f.	1.87, 7 d.f.				

Responses to treatments

	Mean	Ploughing		Dung		Phosphate		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
<u>1944</u>									
Ploughing									
deep-shallow	0.6	-	-	0.7	0.5	-0.9	2.1	1.6	-0.4
Dung	2.6	2.7	2.5	-	-	1.4	3.8	3.0	2.2
Phosphate	-3.0	-4.5	-1.5	-4.2	-1.8	-	-	-6.1	0.1
Potash	-2.4	-1.4	-3.4	-2.0	-2.8	-5.5	0.7	-	-
<u>1945</u>									
Ploughing	+0.20					+0.29			
deep-shallow	1.0	-	-	1.2	0.8	0.9	1.1	1.0	1.0
Dung	0.5	0.7	0.3	-	-	0.6	0.4	0.7	0.3
Phosphate	-0.5	-0.6	-0.4	-0.4	-0.6	-	-	-0.4	-0.6
Potash	0.6	0.6	0.6	0.8	0.4	0.7	0.5	-	-
<u>1946</u>									
Ploughing	+0.58					+0.83			
deep-shallow	0.5	-	-	0.5	0.5	0.8	0.2	1.0	0.0
Dung	-0.4	-0.4	-0.4	-	-	0.3	-1.1	-0.2	-0.6
Phosphate	-1.6	-1.3	-1.9	-0.9	-2.3	-	-	-1.7	-1.5
Potash	0.7	1.2	0.2	0.9	0.5	0.6	0.8	-	-
<u>1947</u>									
Ploughing									
deep-shallow	-0.3	-	-	-0.8	0.2	-0.4	-0.2	-0.7	0.1
Dung	2.2	1.7	2.7	-	-	2.1	2.3	3.6	0.8
Phosphate	-0.8	-0.9	-0.7	-0.9	-0.7	-	-	-1.0	-0.6
Potash	+2.2	+1.8	+2.6	+3.6	+0.8	2.0	2.4	-	-

alluvial

Deep Cultivation Rotation Experiment

Potatoes

Ware tubers: tons per acre

Time of application of mineral fertilizers

Standard errors	Superphosphate			Potash			Mean
	None	Ploughed in bed	In seed	None	Ploughed in bed	In seed	
<u>1944</u>							
Shallow	10.98	11.01	9.97	10.11	11.33	11.38	10.73
Deep	10.29	10.89	12.16	10.37	10.98	11.90	10.91
No dung	9.99	10.24	10.52	9.59	10.56	11.00	10.18
Dung	11.28	11.66	11.61	10.89	11.76	12.29	11.46
Mean	10.64	10.95	11.06	10.24	11.16	11.64	10.82
<u>1945</u>							
Shallow	9.62	10.76	11.17	9.61	10.86	11.07	10.29
Deep	11.92	11.16	11.93	11.32	11.47	12.82	11.73
No dung	9.18	9.22	9.88	8.47	9.94	10.58	9.37
Dung	12.36	12.69	13.22	12.46	12.39	13.32	12.66
Mean	10.77	10.96	11.55	10.47	11.16	11.95	11.01
<u>1946</u>							
Shallow	10.56	10.87	11.51	10.83	10.10	11.75	10.88
Deep	11.73	11.21	12.27	11.10	12.22	12.53	11.73
No dung	10.33	10.06	11.59	10.40	10.04	11.46	10.57
Dung	11.96	12.03	12.18	11.52	12.28	12.81	12.03
Mean	11.14	11.04	11.89	10.96	11.16	12.14	11.30
<u>1947</u>							
Shallow	5.40	5.47	6.26	5.36	5.36	6.46	5.63
Deep	5.75	5.20	6.01	4.90	5.71	7.19	5.68
No dung	4.77	4.57	5.23	3.93	5.04	6.45	4.84
Dung	6.38	6.11	7.04	6.34	6.03	7.20	6.48
Mean	5.58	5.34	6.13	5.13	5.54	6.82	5.66
Standard errors:	(a)			(b)		(c)	
1944	0.43			0.55		0.58	
1945	0.46			0.27		0.50	
1946	0.32			0.54		0.50	
1947	0.27			0.20		0.30	

Standard errors (b) are for use in horizontal comparisons only, (a) and (c) for all other comparisons.

Deep Cultivation Rotation Experiment
 Potatoes
 Percentage ware

Time of application of mineral fertilizers

	Superphosphate			Potash			Mean
	None	Ploughed in	In seed bed	None	Ploughed in	In seed bed	
Standard errors:	(a)	(b) & (c)		(a)	(b) & (c)		
<u>1944</u>							
Shallow	80.0	76.3	74.8	78.5	76.9	77.48	77.8
Deep	79.2	77.4	78.1	80.2	76.2	77.4	78.5
No dung	77.8	76.4	75.7	77.8	75.7	76.1	76.8
Dung	81.5	77.3	77.2	80.8	77.4	78.6	79.4
Mean	79.6	76.8	76.4	79.3	76.6	77.4	78.2
<u>1945</u>							
Shallow	96.0	95.3	95.6	95.4	95.6	96.4	95.7
Deep	97.0	96.8	96.4	96.5	96.8	97.3	96.8
No dung	96.2	95.8	95.9	95.6	96.0	96.9	96.1
Dung	96.8	96.2	96.2	96.3	96.5	96.8	96.5
Mean	96.5	96.0	96.0	96.0	96.2	96.8	96.3
<u>1946</u>							
Shallow	95.7	94.5	94.3	94.5	95.9	95.3	95.0
Deep	96.5	95.6	93.6	95.5	95.9	95.4	95.6
No dung	95.9	96.0	94.2	95.0	95.9	96.0	95.5
Dung	96.2	94.1	93.8	94.9	96.0	94.6	95.1
Mean	96.0	95.0	94.0	95.0	96.0	95.3	95.3
<u>1947</u>							
Shallow	93.8	93.2	92.5	92.4	93.9	94.6	93.3
Deep	93.4	92.6	92.7	91.7	94.0	94.6	93.0
No dung	92.5	91.8	91.4	90.2	93.3	94.4	92.1
Dung	94.6	94.0	93.8	93.9	94.6	94.8	94.3
Mean	93.6	92.9	92.6	92.1	93.9	94.6	93.2

Standard errors:	(a)	(b)	(c)
1944	0.54	0.72	0.74
1945	0.20	0.16	0.23
1946	0.58	0.94	0.88

Standard errors (b) are for use in horizontal comparisons only, (a) and (c) for all other comparisons.

Spring Oats - Residual Effects

1945	16.8	1.12 or 6.6%, 6 d.f.	28.0	1.21 or 4.3%, 7 d.f.
1946	41.9	1.86 or 4.4%, 5 d.f.	57.3	2.72 or 4.8%, 5 d.f.
1947	28.9	1.97 or 6.8%, 4 d.f.	35.3	

Responses to treatments

Mean	Shallow	Deep	Ploughing	Dung	Phosphate	Potash	
			Abs.	Pres.	Abs.	Pres.	
Grain: cwt. per acre							
<u>1945</u>				±0.79			
Ploughing	+0.56						
deep-shallow	-0.9	-	-0.3	-1.5	-1.8	0.0	-0.9
Dung	1.6	2.2	1.0	-	1.4	1.8	0.8
Phosphate	±0.6	-0.3	1.5	0.4	0.8	-	1.2
Potash	0.3	0.3	0.3	-0.5	1.1	0.9	-0.3
<u>1946</u>				±1.32			
Ploughing	+0.93						
deep-shallow	0.6	-	-	2.0	-0.8	1.1	0.1
Dung	2.5	3.9	1.1	-	-	3.4	1.6
Phosphate	1.8	2.3	1.3	2.7	0.9	-	-
Potash	-0.3	-1.8	1.2	1.6	-2.2	0.5	-1.1
<u>1947</u>				±1.39			
Ploughing	+0.98						
deep-shallow	-1.5	-	-	-0.6	-2.4	-1.8	-1.2
Dung	80.8	1.7	-0.1	-	-	2.5	-0.9
Phosphate	1.0	0.7	1.3	2.7	-0.7	-	-
Potash	-0.6	0.1	-1.3	0.8	-2.0	-0.8	-0.4
Straw: cwt. per acre							
<u>1945</u>				±0.86			
Ploughing	+0.60						
deep-shallow	-0.1	-	-	0.8	-1.0	0.4	-0.6
Dung	1.5	2.4	0.6	-	-	0.3	2.7
Phosphate	-1.6	-1.1	-2.1	-2.8	-0.4	-	-0.1
Potash	1.2	1.1	1.3	1.1	1.3	2.7	-0.3
<u>1946</u>				±1.92			
Ploughing	+1.36						
deep-shallow	0.2	-	-	0.3	0.1	-1.5	1.9
Dung	6.5	6.6	6.4	-	-	9.0	4.0
Phosphate	2.6	0.9	4.3	5.1	0.1	-	-
Potash	-0.3	0.3	-0.9	1.1	-1.7	-0.9	0.3
<u>1947</u>							
Ploughing							
deep-shallow	-2.4	-	-	-1.7	-3.1	-2.3	-2.5
Dung	1.4	2.1	0.7	-	-	2.1	0.7
Phosphate	-0.6	-0.5	-0.7	0.1	-1.3	-	-0.7
Potash	0.5	0.1	0.9	1.9	-0.9	0.4	0.6

GRAZING EXPERIMENT

Highfield (begun in 1937)The residual manurial value of feeding stuffs consumed on grassland

The experiment consisted of 3 blocks each of 3 plots. Each year one block began a new three-year cycle. In the first year of each cycle all three plots of the block were grazed with cattle only, additional feeding stuffs being fed on one plot. In the following winter or early spring another plot of each block received fertilizers estimated to be equivalent to the residual manurial value of the feeding stuffs consumed on the first plot in the previous year. The third plot of each block received neither feeding stuff nor fertilizer. In the second and third years of the cycle each of the plots was grazed by cattle and sheep which were weighed regularly.

Area of each plot, 5 acres.

Details of the design of the experiment are as given in the 1937 Station Report, pp. 24-27.

In order to obtain a single measure of the yield of a plot, the amount of starch equivalent produced on each plot in each year has been calculated. In this way it is possible to include in a single figure the energy value of the food required to produce the observed live-weight increases, together with the energy value required to maintain the animals. The calculations are based upon the tables of average composition given by Woodman, H.E., in "Rations for Livestock", Min. of Agric. and Fish. Bull. No. 48. H.M.S.O., 1948.

A full report of the experiment is given by Boyd, D.A., Crowther, E.M., Moffatt, J.R. and Yates, F. in a paper entitled "A grazing experiment on residual manurial value of feeding stuffs consumed on grass". J.R.A.S.E., 110, (1949), 104-114.

Herbage analyses were made by the Botany Department.

Amounts of Feeding Stuffs and Fertilizers applied

Year	Plot	Feeding Stuffs cwt. per acre			Year	Plot	Fertilizers cwt. per acre		
		1st Period	2nd Period	Total			N	P ₂ O ₅	K ₂ O
1938	4	3.00	5.98	8.98					
1938	2	3.38	6.74*	10.12	1939	6	0.17	0.06	0.07
1939	7	3.06	3.78*	3.24	1939	3	0.19	0.07	0.07
1940	4	3.24	4.80	8.04	1940	8	0.17	0.05	0.05
1941	2	4.00	8.91	12.91	1941	6	0.16	0.05	0.06
1942	7	2.89	5.11	8.00	1942	3	0.25	0.09	0.09
1943	4	3.80	4.31	8.11	1943	8	0.16	0.06	0.05
1944	2	3.35	7.40	10.75	1944	6	0.17	0.07	0.07
1945	7	4.52	4.52	9.03	1945	3	0.25	0.09	0.10
1946	4	3.65	5.54	9.19	1946	8	0.18	0.07	0.06
					1947	6	0.19	0.08	0.07

*Plus 1.41 second-period ration fed during first period.
Feeding stuffs; equal parts of flaked maize and undecorticated cotton cake (1st period) or decorticated groundnut cake (2nd period).

Starch Equivalent: lb. per acre

Plot	No Manure	1		2		3		4		5		6		7		8	
		Cake 44	1938,41, Equivalent Manures	No Manure	Cake 44	1938,41, Equivalent Manures	No Manure	Cake 43,46	1938,40, Equivalent Manures	No Manure	Cake 43,46	1939,42, Equivalent Manures	No Manure	Cake 45	1939,42, Equivalent Manures	No Manure	Cake 45
1937	Cattle	779	763	710	544	576	594	672	629	616							
	Sheep	116	142	117	134	112	101	86	119	79							
1938	Cattle							639	648	613							
	Sheep							653	670	664							
1939	Cattle	864	1033	985	917	1022	1059	769	993	963							
	Sheep	746	724	807	746	774	848	646	650	625							
1940	Cattle	680	847	719	747	1108	792	1027	976	950							
	Sheep	601	560	631	598	713	629	702	655	652							
1941	Cattle				775	813	783										
	Sheep				822	656	838										
1942	Cattle	780	849	768	747	1108	792	882	1195	1163							
	Sheep	716	852	811	598	713	629	908	999	863							
1943	Cattle	870	872	1086	775	813	783	1074	1399	945							
	Sheep	689	728	936	822	656	838	622	680	591							
1944	Cattle				1182	1017	1408										
	Sheep				716	551	674										
1945	Cattle				1079	1006	1174										
	Sheep				696	624	731										
1946	Cattle	811	1231	1037	1079	1006	1174	1254	1799	1489							
	Sheep	445	680	743	696	624	731	712	689	690							
1947	Cattle	1296	1627	1312	1312	1254	1625	1397	1694	1290							
	Sheep	646	631	568	1549	1042	582	383	420	556							

Highfield Grazing Experiment

Bd/1.3

Live weight increase: lb. per acre

Plot	1		2		3		4		5		6		7		8	
	No Manure	Cake 1938, 41, Equivalent Manures	No Manure	Cake 1938, 40, Equivalent Manures	No Manure	Cake 1938, 40, Equivalent Manures	No Manure	Cake 1938, 40, Equivalent Manures	No Manure	Cake 1939, 42, Equivalent Manures	No Manure	Cake 1939, 42, Equivalent Manures	No Manure	Cake 1939, 42, Equivalent Manures	No Manure	Cake 1939, 42, Equivalent Manures
1937	Cattle	181	176	151	6	105	118	126	4	140	118	9	114	-1		
	Sheep	3	13	1	6	14	6	4	4	0	0	9	114	-1		
1938	Cattle															
	Sheep															
1939	Cattle	193	274	231	123	205	237	242	130	148	235	96	229	96	229	96
	Sheep	114	100	133	133	115	107	130	144	84	111	192	180	180	180	180
1940	Cattle	135	178	74	95	128	214	147	217	91	74	72	72	72	72	72
	Sheep	91														
- 1941	Cattle															
	Sheep															
1942	Cattle	154	177	133	127	160	190	144	124	237	312	300	300	300	300	300
	Sheep	104	145	127	135	96	135	144	124	315	113	79	79	79	79	79
1943	Cattle	222	221	263	108	332	296	404	219	52	398	236	236	236	236	236
	Sheep	69	84	84	108	54	30	19	53	52	31	29	29	29	29	29
1944	Cattle															
	Sheep															
1945	Cattle	175	276	237	68	209	210	219	53	67	278	388	328	328	328	328
	Sheep	38	54	339	264	74	53	67	48	67	319	63	56	56	56	56
1946	Cattle	276	276	339	59	343	215	373	81	73	38	378	290	290	290	290
	Sheep	62	62	59	48	46						48	90	90	90	90
1947	Cattle															
	Sheep															

- 2 -

Plot	Grazing days per acre							
	1 No Manure	2 Cake 1938, 41, 44	3 Equivalent Manures	4 No Manure	5 1938, 40, 43, 46	6 Cake Equivalent Manures	7 No Manure	8 1939, 42 Equivalent Manures
1937	Cattle Sheep	74 74	74 74	62 65	63 65	63 65	66 62	66 62
1938	Cattle Sheep	107 344	110 356	114 342	113 343	120 362	126 379	83 253
1939	Cattle Sheep	96 288	122 308	112 314				253
1940	Cattle Sheep							
1941	Cattle Sheep							
1942	Cattle Sheep	115 339	121 355	121 366	119 358	153 457	147 358	121 361
1943	Cattle Sheep	96 288	125 288	125 373	106 338	102 316	117 374	145 434
1944	Cattle Sheep							
1945	Cattle Sheep	92 198	133 293	112 303	113 346	92 285	127 389	126 301
1946	Cattle Sheep	122 259	155 259	132 267	131 268	115 257	149 301	145 308
1947	Cattle Sheep							

GREEN MANURING EXPERIMENT

Woburn Stackyard (begun in 1936)

The experiment was designed to test the effects on kale of leys and green manures ploughed-in in mid-season, and also the effects of dung, straw and sulphate of ammonia. A barley crop tests the residual effects of these treatments.

The rotation was 1st year, green manure crops followed by kale, 2nd year barley. The leys and green manures were clover, ryegrass, tares and mustard: the tares and mustard were grown on the same plots every second year, but the clover and ryegrass alternately. A crop of hay was taken from the ley plots each year until 1940 and again in 1942; in other years all green material was buried.

There are 40 plots under each of the test crops, and until 1943 the experiment consisted of a single replicate (for each crop) of a 5×2^2 factorial design with all combinations of the following treatments, applied to kale;

Leys and green manures: Fallow (F), clover (C), ryegrass (R), tares (T) and mustard (M). The clover and ryegrass are undersown in the barley and the other green manures are sown after the barley stubble has been ploughed.

Dung: None, 10 tons per acre (D)

Straw: None, 1.5 tons per acre (S)

Sulphate of ammonia: 0.4, 0.8 cwt. N per acre (N)

A basal dressing of 3 cwt. superphosphate and 1 cwt. muriate of potash per acre is applied.

In 1944 and succeeding years a top dressing of sulphate of ammonia (0.3 cwt. N per acre in 1944, 0.4 cwt. N per acre thereafter) has been applied to half the plots under barley so that the experiment is now in half replicate according to the identity $I \equiv (R+C-T-M-F) DSNA$, where A represents the top dressing of sulphate of ammonia.

From 1946 onwards, a further dressing of sulphate of ammonia (0.4 cwt. N per acre) has been applied to the fallow and all ley and green manure crops, on those plots which receive the top dressing when under barley. Also from 1946 onwards, cabbages replaced the kale (which had failed in several years), and the green manures were changed, tares being replaced by lupins and mustard by rape.

Full details of the original design are given in the 1936 Report, p. 203.

Owing to an error in the chain used, the plot area has been given previously as 0.0367 acre. The correct value is 0.0379 acre. Consequently the yields given in the Station Reports for 1936-38 should be multiplied by 0.968.

Crop Notes

Test Year

	1939	1940	1941	1942
<u>Leys and Green Manures</u>				
Clover and Ryegrass				
undersown	31.3.38	19.4.39	3.5.40	5.5.41
cut	June 8	June 3		June 2
Tares sown	6.10.38	Apr. 23 ⁽¹⁾	16.10.40	Apr. 20 ⁽²⁾
Mustard sown	6.10.38	Apr. 23 ⁽¹⁾	Mar 24 ⁽¹⁾	Apr. 15 ⁽¹⁾
Manures ploughed in	June 16	June 20	May 23	June 13
<u>Kale (Thousand Head)</u>				
Sown	June 21	June 26		June 16
Cut	Feb. 1940	Feb. 1941	Failed ⁽³⁾	Feb and Mar. 1943
<u>Cabbages (January King)</u>				
Transplanted				
Cut				
<u>Barley (Plumage Archer)</u>				
Sown	Mar. 1	Mar. 4	Mar. 14	Apr. 1
Harvested	Aug. 29	Aug. 26	Aug. 26	Aug. 19

(1) Second sowing. First sowing failed and was ploughed in

(2) Maple peas, replacing tare crop which failed

(3) First sowing destroyed by flea beetle; second sowing by pigeons

Green Manuring Experiment

Be/1.3

Crop Notes

	Test Year				
	1943	1944	1945	1946	1947
Clover and Ryegrass					
undersown				23.4.45	2.5.46
	(Leys and green manures were ploughed in after failure of the kale.)				
Lupins sown				Apr.15	Apr.16
Rape sown				Apr.15 ⁽¹⁾	Apr.16
Manures ploughed in				June 18	June 28
Kale (Thousand Head)					
Sown				(Fallow)	
Cut	Failed ⁽⁴⁾	Failed ⁽⁴⁾			
Cabbages (January King)					
Transplanted				July 22 ⁽⁵⁾	July 12 ⁽⁵⁾
Cut				Jan. & Mar. 1947	Dec. Feb. & Mar. 1948
Barley (Plumage Archer)					
Sown	Mar. 17	Mar. 9	Mar. 6	Mar. 18	Apr. 18
Harvested	Aug. 20	Sept. 9	Aug. 3	Aug. 23	Aug. 12
(1) Second sowing. First sowing failed and was ploughed in.					
(4) Crop failed due to attack of flea beetle.					
(5) Gaps were filled in by further transplanting during subsequent two months.					

Kale

Be/1.4

Leys and green manures:	None	Tares	Clover	Mustard	Rye-grass	Mean
Manures						
Total weight: tons per acre						
					<u>1939</u>	
					<u>±0.418</u>	<u>±0.187</u>
No dung	4.54	4.62	6.31	4.57	3.61	4.73
Dung	5.87	6.27	8.27	6.11	4.38	6.18
No straw	5.49	5.92	7.57	5.27	4.21	5.69
Straw	4.92	4.96	7.02	5.41	3.77	5.22
Sulphate of ammonia						
0.4 cwt.N per acre	4.56	4.69	6.58	4.79	2.44	4.61
0.8 cwt.N per acre	5.85	6.20	8.01	5.89	5.55	6.30
Mean <u>±0.296</u>	5.20	5.44	7.29	5.34	3.99	5.46
					<u>1940</u>	
					<u>±0.562</u>	<u>±0.251</u>
No dung	6.16	5.54	5.62	4.59	3.43	5.07
Dung	7.09	7.24	8.04	6.85	5.52	6.95
No straw	6.73	6.53	6.86	6.08	5.07	6.26
Straw	6.52	6.24	6.80	5.35	3.88	5.76
Sulphate of ammonia						
0.4 cwt.N per acre	5.46	5.25	6.95	5.07	4.18	5.38
0.8 cwt.N per acre	7.79	7.52	6.71	6.36	4.77	6.63
Mean <u>±0.397</u>	6.62	6.39	6.83	5.72	4.47	6.01
					<u>1942</u>	
					<u>±0.725</u>	<u>±0.324</u>
No dung	8.73	9.11	11.18	9.08	5.88	8.80
Dung	11.46	11.92	13.61	11.54	10.02	11.71
No straw	9.74	9.53	11.92	10.18	8.28	9.93
Straw	10.45	11.49	12.87	10.43	7.62	10.57
Sulphate of ammonia						
0.4 cwt.N per acre	9.01	9.63	12.67	8.52	5.63	9.09
0.8 cwt.N per acre	11.17	11.40	12.12	12.10	10.27	11.41
Mean <u>±0.513</u>	10.09	10.51	12.40	10.31	7.95	10.25

Note: The kale failed in 1941, 1943 and 1944. The land was fallowed in 1945.

Standard errors per plot

1939 0.84 or 15.3%

1940 1.12 or 18.7%

1942 1.45 or 14.1%

These are based on 16 d.f.

Green Manuring Experiment

Be/1.5

Cabbages

Leys and green manures:	None	Lupins	Clover	Rape	Rye-grass	Means	Mean
Manures		Total weight: tons per acre					
			1946 ±0.721				±0.322
No dung	2.96	3.27	4.68	3.20	2.46	3.31	
Dung	4.80	4.09	6.67	4.44	3.78	4.76	
No straw	3.52	4.29	5.90	4.27	2.84	4.16	
Straw	4.24	3.06	5.45	3.37	3.41	3.91	
Sulphate of ammonia							
0.4 cwt.N per acre	4.08	2.92	5.22	2.93	2.58	3.55	
0.8 cwt.N per acre	3.68	4.44	6.12	4.71	3.66	4.52	
Sulphate of ammonia to green manures.							
None	3.10	3.87	6.24	2.43	3.46	3.82	
0.4 cwt.N per acre	4.66	3.48	5.10	5.20	2.79	4.25	
Mean ±0.540	3.88	3.68	5.67	3.82	3.12	4.03	
			1947 ±0.430				±0.192
No dung	3.76	2.36	1.20	1.83	1.87	2.20	
Dung	4.40	3.11	1.11	2.36	2.04	2.60	
No straw	4.32	2.42	1.21	2.05	2.35	2.47	
Straw	3.83	3.06	1.10	2.14	1.56	2.34	
Sulphate of ammonia							
0.4 cwt.N per acre	3.38	2.43	0.97	1.96	1.94	2.14	
0.8 cwt.N per acre	4.77	3.04	1.34	2.22	1.98	2.67	
Sulphate of ammonia to green manures.							
None	3.95	2.92	1.12	2.07	1.85	2.38	
0.4 cwt.N per acre	4.20	2.56	1.19	2.12	2.06	2.43	
Mean ±0.304	4.08	2.74	1.16	2.09	1.96	2.40	

Standard errors per plot

1946 1.44 or 35.7%
1947 0.86 or 35.8%

These are based on 9 d.f.

Cabbages

Be/1.6

Means

Leys and green manures:	None	Lupins	Clover	Rape	Rye-grass	Mean
Manures						
	Plant number:	thousands per acre				
		<u>1946</u>				
			<u>±0.80</u>			<u>±0.36</u>
No dung	16.8	16.0	16.0	16.4	15.6	16.2
Dung	16.2	15.7	16.0	16.9	16.2	16.2
No straw	16.1	17.0	15.8	16.9	15.7	16.3
Straw	16.8	14.7	16.2	16.5	16.1	16.1
Sulphate of ammonia						
0.4 cwt.N per acre	16.7	16.0	16.5	16.5	16.6	16.5
0.8 cwt.N per acre	16.3	15.7	15.5	16.8	15.3	15.9
Sulphate of ammonia to green manures.						
None	16.8	16.4	15.6	16.6	16.8	16.4
0.4 cwt.N per acre	16.1	15.4	16.4	16.7	15.1	15.9
Mean <u>±0.61</u>	16.5	15.9	16.0	16.7	15.9	16.2
<u>1947</u>						
		<u>±0.76</u>				<u>±0.34</u>
No dung	17.1	16.9	15.6	15.8	15.5	16.2
Dung	17.4	17.6	14.9	15.1	16.0	16.2
No straw	17.3	17.4	14.7	16.1	16.9	16.5
Straw	17.2	17.2	15.7	14.9	14.6	15.9
Sulphate of ammonia						
0.4 cwt.N per acre	16.4	17.6	15.7	16.4	16.0	16.4
0.8 cwt.N per acre	18.1	16.9	14.7	14.6	15.6	16.0
Sulphate of ammonia to green manures.						
None	16.7	17.7	14.4	14.9	15.8	15.9
0.4 cwt.N per acre	17.8	16.9	16.0	16.1	15.7	16.5
Mean <u>±0.54</u>	17.3	17.3	15.2	15.5	15.8	16.2

Standard errors per plot.

1946 1.60 or 9.9%
 1947c 1.52 or 9.4%

These are based on 9 d.f.

Green Manuring Experiment

Be/1.7

Differential Responses

Kale. Total weight: tons per acre

Response to	Mean	Dung		Straw		per acre		Sulphate of Ammonia to Kale cwt.N		to Barley and Green crops	
		Abs.	Pres.	Abs.	Pres.	0.4	0.8	Abs.	Pres.	Abs.	Pres.
<u>1939</u>	± 0.264					± 0.374					
Dung	1.45	-	-	0.90	2.00	1.88	1.02				
Straw	-0.48	-1.03	0.07	-	-	-0.70	-0.26				
Sulph.of Amm.	1.69	2.12	1.26	1.47	1.91	-	-				
<u>1940</u>	± 0.355					± 0.502					
Dung	1.88	-	-	1.76	2.00	2.43	1.33				
Straw	-0.50	-0.62	-0.38	-	-	-1.45	0.45				
Sulph.of Amm.	1.25	1.80	0.70	0.30	2.20	-	-				
<u>1942</u>	± 0.458					± 0.648					
Dung	2.91	-	-	3.06	2.76	3.34	2.48				
Straw	0.64	0.79	0.49	-	-	0.33	0.95				
Sulph.of Amm.	2.32	2.75	1.89	2.01	2.63	-	-				

Cabbages. Total weight: tons per acre

<u>1946</u>	± 0.456			± 0.651							
Dung	1.45	-	-	1.76	1.14	2.54	0.36	1.43	1.47		
Straw	-0.26	0.05	-0.57	-	-	-0.12	-0.40	-1.40	0.88		
Sulph.of Amm.	0.98	2.07	-0.11	1.12	0.84	-	-	1.60	0.36		
Sulph.of Amm. to Green crops	0.43	0.41	0.45	-0.71	1.57	1.05	-0.19	-	-		

<u>1947</u>	± 0.272			± 0.389							
Dung	0.40	-	-	0.24	0.56	0.39	0.41	0.53	0.27		
Straw	-0.13	-0.29	0.03	-	-	0.14	-0.40	0.03	-0.29		
Sulph.of Amm.	0.53	0.52	0.54	0.80	0.26	-	-	0.44	0.62		
Sulph.of Amm. to Green crops	0.04	0.17	-0.09	-0.20	-0.12	-0.05	0.13	-	-		
				$+0.20$							

Cabbages. Plant Number: thousands per acre

<u>1946</u>	± 0.51			± 0.72							
Dung	0.0	-	-	-0.1	0.1	0.7	-0.7	-0.4	0.4		
Straw	-0.2	-0.3	-0.1	-	-	0.1	-0.5	-1.2	0.8		
Sulph.of Amm. to Cabbages	-0.5	0.2	-1.2	-0.2	-0.8	-	-	-1.4	0.4		
Sulph.of Amm. to Green crops	-0.5	-0.9	-0.1	-1.5	$+0.5$	-1.4	0.4	-	-		
<u>1947</u>	± 0.48			± 0.69							

Dung	0.0	-	-	-0.6	0.6	0.6	-0.6	0.4	-0.4		
Straw	-0.5	-1.1	0.0	-	-	-1.1	0.0	-0.6	-0.5		
Sulph.of Amm. to Cabbages	-0.4	0.1	-1.0	-1.0	0.1	-	-	-0.8	-0.1		
Sulph.of Amm. to Green crops	0.6	1.0	0.2	0.5	0.7	0.3	0.9	-	-		

Barley

Grain: cwt. per acre

Means

Leys and green manures before kale:	None	Tares	Clover	Mustard	Rye- grass	Mean
Manures applied to previous kale			<u>1939</u>			
			<u>±1.16</u>			<u>±0.52</u>
No dung	12.8	15.1	18.7	13.8	14.1	14.9
Dung	15.8	15.9	20.6	17.0	18.4	17.5
No straw	12.4	15.1	19.9	16.2	16.2	16.0
Straw	16.2	15.9	19.5	14.6	16.5	16.6
Sulphate of Ammonia						
0.4 cwt.N per acre	15.0	15.0	20.9	14.2	13.5	15.7
0.8 cwt.N per acre	13.6	16.0	18.4	16.7	19.1	16.8
Mean <u>±0.82</u>	14.3	15.5	19.7	15.4	16.3	16.3
			<u>1940</u>			
			<u>±1.53</u>			<u>±0.68</u>
No dung	10.5	11.0	13.4	10.6	7.7	10.7
Dung	13.7	15.0	16.1	11.9	11.9	13.8
No straw	10.8	12.4	13.7	12.1	9.3	11.6
Straw	13.3	13.7	15.8	10.4	10.4	12.7
Sulphate of Ammonia						
0.4 cwt. N per acre	9.6	11.8	15.3	11.1	9.1	11.4
0.8 cwt. N per acre	14.5	14.2	14.1	11.3	10.6	13.0
Mean <u>±1.08</u>	12.0	13.0	14.7	11.2	9.8	12.2
			<u>1941</u>			
			<u>±0.86</u>			<u>±0.38</u>
No dung	11.0	11.5	12.5	10.1	11.4	11.3
Dung	12.9	12.6	14.7	11.7	14.4	13.3
No straw	12.3	11.5	13.8	11.0	12.8	12.3
Straw	11.6	12.6	13.6	10.7	13.0	12.3
Sulphate of Ammonia						
0.4 cwt.N per acre	11.7	10.7	12.0	9.5	12.5	11.3
0.8 cwt.N per acre	12.2	13.4	15.3	12.3	13.3	13.3
Mean <u>±0.61</u>	12.0	12.0	13.7	10.8	12.9	12.3

Barley

Grain: cwt. per acre

Means

Leys and green manures before kale:	None	Tares	Clover	Mustard	Rye- grass	Mean
Manures applied to previous kale			<u>1942</u>			
No dung	9.0	8.3	7.1	8.1	9.8	8.5
Dung	11.2	12.1	10.7	10.3	12.0	11.3
No straw	9.6	8.9	8.1	9.4	9.8	9.2
Straw	10.6	11.4	9.7	9.0	12.0	10.5
Sulphate of Ammonia						
0.4 cwt.N per acre	9.0	10.4	8.3	9.6	11.1	9.7
0.8 cwt.N per acre	11.3	10.0	9.5	8.8	10.7	10.1
Mean ± 0.59	10.1	10.2	8.9	9.2	10.9	9.9
			<u>1943</u>			
			± 1.76			± 0.79
No dung	7.4	11.0	9.9	7.1	6.6	8.4
Dung	13.1	11.4	11.2	13.2	11.2	12.0
No straw	9.1	12.3	11.6	8.7	8.8	10.1
Straw	11.5	10.1	9.5	11.6	9.0	10.3
Sulphate of Ammonia						
0.4 cwt.N per acre	10.2	12.0	10.4	8.9	9.6	10.2
0.8 cwt.N per acre	10.4	10.4	10.7	11.3	8.2	10.2
Mean ± 1.24	10.3	11.2	10.6	10.1	8.9	10.2
			<u>1944</u>			
			± 1.06			± 0.48
No dung	9.9	10.1	8.6	8.2	7.0	8.8
Dung	8.6	10.5	8.1	8.7	12.0	9.6
No straw	9.6	10.4	7.4	8.6	7.9	8.8
Straw	8.9	10.2	9.3	8.4	11.1	9.6
Sulphate of Ammonia						
0.4 cwt.N per acre	10.0	10.6	7.5	9.3	9.2	9.3
0.8 cwt.N per acre	8.5	10.0	9.3	7.7	9.8	9.1
Sulphate of Ammonia applied to Barley						
None	7.3	9.4	6.5	7.9	9.8	8.2
0.3 cwt.N per acre	11.2	11.2	10.3	9.1	9.2	10.2
Mean ± 0.75	9.2	10.3	8.4	8.5	9.5	9.2

Be/1.10

Barley

Grain: cwt. per acre

Means

Leys and green manures before kale	None	Tares	Clover	Mustard	Rye- grass	Mean
			<u>1945</u>			
			±0.60			±0.27
No dung	13.4	15.4	13.2	13.6	13.2	13.8
Dung to kale	16.7	15.2	15.8	18.2	14.2	16.0
No straw	16.3	14.3	14.1	15.5	13.7	14.8
Straw to kale	13.8	16.3	14.8	16.3	13.7	15.0
Sulph. of Amm. to kale						
0.4 cwt.N per acre	16.2	15.4	14.1	15.8	13.2	14.9
0.8 cwt.N per acre	13.9	15.2	14.9	16.0	14.2	14.8
Sulph. of Amm. to Barley						
None	10.9	10.7	10.3	9.6	9.8	10.3
0.4 cwt.N per acre	19.2	19.9	18.6	22.2	17.6	19.5
Mean	±0.42	15.0	15.3	14.5	15.9	13.7
			<u>1946</u>			
			±0.90			±0.40
No dung	10.5	9.4	9.8	10.6	9.9	10.0
Dung to kale in 1945 [#]	9.6	10.5	10.2	9.8	9.8	10.0
No straw	10.5	9.5	11.4	9.6	8.4	9.9
Straw to kale in 1945 [#]	9.6	10.4	8.6	10.8	11.3	10.1
Sulph. of Amm. to kale in 1945 [#]						
0.4 cwt.N per acre	10.3	10.4	8.4	11.6	7.8	9.7
0.8 cwt.N per acre	9.8	9.5	11.6	8.8	11.8	10.3
Sulph. of Amm. to Barley						
None	5.5	6.4	4.4	6.8	5.8	5.8
0.4 cwt.N per acre	14.6	13.5	15.7	13.6	13.8	14.2
Mean	±0.63	10.0	10.0	10.0	10.2	9.8
						10.0

^KThe Kale crop of 1945 failed and therefore no manures were applied.

Green Manuring Experiment

Be/1.11

Barley

Grain: cwt. per acre

Means

Leys and green manures before cabbages	None	Lupins	Clover	Rape	Rye- grass	Mean
<u>1947</u>						
±0.96						±0.43
No dung	14.2	15.2	12.0	14.2	13.1	13.7
Dung to cabbages	18.3	17.8	16.2	18.0	16.6	17.4
No straw	16.5	16.1	14.0	16.1	14.8	15.5
Straw to cabbages	16.1	16.8	14.2	16.1	14.9	15.6
Sulphate of ammonia to Cabbages						
0.4 cwt.N per acre	16.2	16.0	13.4	16.3	15.9	15.6
0.8 cwt.N per acre	16.4	16.9	14.9	15.9	13.9	15.6
Sulphate of ammonia to Barley						
None	14.3	14.3	13.1	13.8	13.2	13.7
0.4 cwt.N per acre	18.3	18.6	15.1	18.5	16.6	17.4
Mean ±0.68	16.3	16.5	14.1	16.1	14.9	15.6

Barley

Straw: cwt. per acre

Means

Leys and green manures before kale:	None	Tares	Clover	Mustard	Rye- grass	Mean
<u>1943</u>						
No dung	8.9	12.5	14.2	8.8	8.9	10.7
Dung to kale	15.8	14.2	17.2	17.4	15.0	15.9
No straw	11.1	14.5	17.1	11.9	11.6	13.2
Straw to kale	13.6	12.2	14.3	14.4	12.3	13.4
Sulphate of ammonia to kale						
0.4 cwt.N per acre	12.4	13.5	15.2	12.2	11.7	13.0
0.8 cwt.N per acre	12.2	13.1	16.2	14.1	12.2	13.6
Mean	12.3	13.3	15.7	13.1	12.0	13.3

Barley

Straw: cwt. per acre

Means

Leys and green manures before kale:	None	Tares	Clover	Mustard	Rye- grass	Mean
	<u>1944</u>					
	<u>±0.97</u>					
No dung	12.3	12.8	11.0	10.6	8.4	11.0
Dung to kale	12.4	15.4	10.8	10.1	14.7	12.7
No straw	12.5	13.5	10.2	10.3	9.9	11.3
Straw to kale	12.1	14.8	11.6	10.3	13.2	12.4
Sulphate of ammonia to kale						
0.4 cwt. N per acre	12.6	15.1	10.5	11.1	11.4	12.1
0.8 cwt. N per acre	12.1	13.1	11.3	9.6	11.7	11.6
Sulphate of ammonia to barley						
None	9.8	13.2	8.3	9.7	12.2	10.6
0.3 cwt.N per acre	14.8	15.1	13.5	10.9	10.9	13.0
Mean <u>±0.69</u>	12.3	14.1	10.9	10.3	11.6	11.3
	<u>1945</u>					
	<u>±1.18</u>					
No dung	15.3	16.2	14.6	14.5	13.8	14.9
Dung to kale	20.8	18.7	18.8	19.5	16.1	18.8
No straw	18.6	16.8	16.3	16.8	14.8	16.7
Straw to kale	17.5	18.1	17.1	17.2	15.1	17.9
Sulphate of ammonia to kale						
0.4 cwt.N per acre	19.0	18.6	15.9	17.3	14.1	17.0
0.8 cwt.N per acre	17.1	16.3	17.5	16.7	15.8	16.7
Sulphate of ammonia to barley						
None	13.4	12.1	12.3	10.7	11.7	12.0
0.4 cwt.N per acre	22.7	22.8	21.1	23.3	18.2	21.6
Mean <u>±0.84</u>	18.0	17.4	16.7	17.0	15.0	16.8

Green Manuring Experiment

Be/1.13

Barley

Straw: cwt. per acre

Means

Leys and green manures before kale(1)	None	Tares (2)	Clover	Mustard (3)	Rye-grass	Mean
<u>1946</u>						
No dung	14.1	12.9	13.6	14.2	11.4	13.2
Dung to kale in 1943 ^x	16.0	17.8	15.2	14.0	12.9	15.2
No straw	14.6	14.1	15.1	14.0	11.2	13.8
Straw to kale in 1943 ^x	15.4	16.6	13.8	14.2	13.1	14.6
Sulphate of ammonia to kale 1943 ^x						
0.4 cwt.N per acre	14.4	15.6	13.8	15.4	11.2	14.1
0.8 cwt.N per acre	15.7	15.1	15.0	12.7	13.1	14.3
Sulphate of ammonia to barley						
None	8.6	10.4	9.4	10.2	7.5	9.2
0.4 cwt.N per acre	21.4	20.3	19.4	17.9	16.8	19.2
Mean	15.0	15.3	14.4	14.1	12.2	14.2
<u>1947</u>						
			±0.78			±0.35
No dung	14.4	15.6	12.4	14.8	14.4	14.3
Dung to cabbages	19.1	18.2	16.4	18.7	18.1	18.1
No straw	16.7	16.9	13.7	17.3	16.0	16.1
Straw to cabbages	16.8	16.9	15.1	16.2	16.5	16.3
Sulphate of ammonia to cabbages						
0.4 cwt.N per acre	16.6	16.3	13.7	16.3	17.0	16.0
0.8 cwt.N per acre	16.9	17.5	15.0	17.1	15.5	16.4
Sulphate of ammonia to barley						
None	14.3	15.2	13.3	14.4	15.3	14.5
0.4 cwt.N per acre	19.2	18.6	15.5	19.1	17.2	17.9
Mean ±0.55	16.8	16.9	14.4	16.7	16.2	16.2

^x The kale crop of 1945 failed and therefore no manures were applied.

Note. The following changes in headings are applicable to the 1947 barley results.

1. Cabbages replaced Kale. 2. Lupins replaced Tares. 3. Rape replaced Mustard.

Standard errors per plot.

Grain:	cwt. per acre	Straw:	cwt. per acre
1939	2.32 or 14.3%	1945	1.21 or 8.1%
1940	3.06 or 25.1%	1946	1.79 or 17.9%
1941	1.72 or 14.0%	1947	1.93 or 12.4%
1942	1.67 or 16.9%		
1943	3.51 or 34.3%		
1944	2.13 or 23.2%		

These are based on 16. d.f., 1939-1943

9 d.f., 1944-1947

Differential Responses

Barley. Grain: cwt. per acre

Response to	Mean	Dung		Straw		Sulphate of Ammonia cwt. N to Barley	
		Abs.	Pres.	Abs.	Pres.	per acre 0.4.	0.8 Abs. Pres.
<u>1939</u>	± 0.74	± 1.04					
Dung	2.6	-	-	3.2	2.0	3.6	1.6
Straw	0.6	1.2	0.0	-	-	0.5	0.7
Sulph. of Amm.	1.0	2.0	0.0	0.9	1.1	-	-
<u>1940</u>	± 0.97	± 1.37					
Dung	3.1	-	-	3.6	2.6	3.8	2.4
Straw	1.1	1.6	0.6	-	-	0.6	1.6
Sulph. of Amm.	1.6	2.3	0.9	1.1	2.1	-	-
<u>1941</u>	± 0.54	± 0.77					
Dung	2.0	-	-	1.9	2.1	2.7	1.3
Straw	0.0	-0.1	0.1	-	-	0.3	-0.3
Sulph. of Amm.	2.0	2.7	1.3	2.3	1.7	-	-
<u>1942</u>	± 0.53	± 0.75					
Dung	2.8	-	-	3.2	2.4	3.4	2.2
Straw	1.3	1.7	0.9	-	-	1.0	1.6
Sulph. of Amm.	0.4	0.9	-0.1	0.1	0.7	-	-
<u>1943</u>	± 0.98	± 1.39					
Dung	3.6	-	-	5.1	2.1	4.7	2.5
Straw	0.2	1.7	-1.3	-	-	2.2	-1.8
Sulph. of Amm.	0.0	1.1	-1.1	2.0	-2.0	-	-
<u>1944</u>	± 0.67	± 0.95					
Dung	0.8	-	-	2.1	-0.5	0.7	0.9
Straw	0.8	2.1	-0.5	-	-	1.5	0.0
Sulph. of Amm.	-0.3	-0.4	-0.2	0.5	-1.0	-	-0.6
Sulph. of Amm. to Barley	2.0	2.0	2.1	2.0	2.1	1.8	2.4
<u>1945</u>	± 0.38	± 0.54					
Dung	2.3	-	-	2.7	1.9	3.1	1.5
Straw	0.2	0.6	-0.2	-	-	0.1	0.3
Sulph. of Amm.	-0.1	0.7	-0.9	-0.2	0.0	-	-1.0
Sulph. of Amm. to Barley	9.2	8.6	9.8	9.2	9.2	8.3	10.1
<u>1946</u>	± 0.57	± 0.81					
Dung	-0.1	-	-	0.1	-0.2	0.5	-0.6
Straw	0.2	0.4	0.1	-	-	-0.7	1.2
Sulph. of Amm.	0.6	1.2	0.1	-0.3	1.5	-	1.7
Sulph. of Amm. to Barley	8.5	8.4	8.6	9.2	7.8	9.6	7.4
<u>1947</u>	± 0.61	± 0.87					
Dung	3.6	-	-	3.9	3.3	5.0	2.2
Straw	0.1	0.4	-0.2	-	-	0.3	-0.1
Sulph. of Amm.	0.0	1.4	-1.4	0.2	-0.2	-	0.4
Sulph. of Amm. to Barley	3.7	2.6	4.8	2.1	5.3	4.1	3.3

All treatments were applied to the previous Kale (or Cabbage) unless otherwise stated.

Differential Responses

Barley. Straw: cwt. per acre

Response to	Mean	Dung				Straw		Sulphate of Ammonia cwt. N to Barley per acre		
		Abs.	Pres.	Abs.	Pres.	0.4	0.8	Abs.	Pres.	
<u>1943</u>										
Dung	5.2	-	-	7.2	3.2	5.6	4.8			
Straw	0.1	2.1	-1.9	-	-	1.7	-1.5			
Sulph. of Amm.	0.6	1.0	0.2	2.2	-1.0	--	-			
	<u>±0.61</u>				<u>±0.87</u>					
<u>1944</u>										
Dung	1.6	-	-	3.3	-0.1	1.5	1.8	0.8	2.4	
Straw	1.1	2.8	-0.6	-	-	2.2	0.0	1.1	1.2	
Sulph. of Amm.	-0.6	-0.8	-0.4	0.5	-1.7	-	-	-1.3	0.1	
Sulph. of Amm. to Barley	2.4	1.6	3.2	2.4	2.4	1.7	3.1	-	-	
	<u>±0.75</u>				<u>±1.07</u>					
<u>1945</u>										
Dung	3.9	-	-	4.7	3.1	5.1	2.7	3.8	4.0	
Straw	0.3	1.1	-0.5	-	-	0.2	0.4	-0.5	1.1	
Sulph. of Amm.	-0.3	0.9	-1.5	-0.4	-0.2	-	-	-1.3	0.7	
Sulph. of Amm. to Barley	9.6	9.5	9.7	8.8	10.4	8.6	10.6	-	-	
	<u>±</u>									
<u>1946</u>										
Dung	2.0	-	-	2.1	1.8	2.6	1.3	1.3	2.6	
Straw	0.8	1.0	0.7	-	-	0.5	1.1	1.7	0.0	
Sulph. of Amm.	0.2	0.9	-0.4	-0.1	0.5	-	-	1.1	-0.7	
Sulph. of Amm. to Barley	9.9	9.3	10.6	10.8	9.1	10.8	9.1	-	-	
	<u>±0.49</u>				<u>±0.71</u>					
<u>1947</u>										
Dung	3.8	-	-	3.0	4.6	4.9	2.7	3.1	4.5	
Straw	0.2	-0.6	1.0	-	-	0.0	0.4	-0.7	1.1	
Sulph. of Amm.	0.4	1.5	-0.7	0.2	0.6	-	-	2.0	-1.2	
Sulph. of Amm. to Barley	3.4	2.7	4.1	2.5	4.3	5.0	1.8	-	-	

All treatments were applied to the previous Kale (or Cabbage) unless otherwise stated.

N

No. replications within years

3 yrs on each series: from each
plot take mean
in Regr

Rem = within-plot errors

Treatment totals \times Series \rightarrow between plot errors

But also replicate within years for first 5 years

LEY AND ARABLE ROTATIONS EXPERIMENT

Woburn, Stackyard, Series D (begun in 1938)

The purpose of the experiment is to test the value of a three year ley, three years of lucerne and an arable rotation with a one year ley, as means of building up soil fertility, in comparison with a rotation without leys. The effects of these crop sequences are measured by the yields of two following crops of potatoes and barley, which may be termed the indicator crops. Each rotation therefore has five courses. The rotations compared are:-

- (1) Three year ley
 - (2) Three years of lucerne
 - (3) Potatoes, wheat, one year ley[#]
 - (4) Potatoes, wheat, kale
- } Potatoes, barley

There are five series, one for each course of the five year rotation, so that all courses of every rotation are represented every year. Each series has eight main plots, on four of which the same rotation continues throughout the experiment. On the other four plots, ley and arable rotations alternate.

Each main plot is divided into two sub-plots, one of which receives dung at the rate of 15 tons per acre applied to the indicator crop of potatoes only. The same sub-plots receive dung throughout the experiment. All plots are liberally manured with inorganic fertilizers.

Details are as given in the 1938 Report pp. 135-137, except that owing to the unsatisfactory crops obtained on the kale plots in the years 1938-44 it was decided to substitute sugar beet for kale as from 1945.

Owing to an error in the chain used, the sub-plot area has been given previously as 0.0390 acre. The correct value is 0.0403 acre. Consequently the yields given in the Station Report for 1938 should be multiplied by 0.968. The correct values for 1938 are included in this Report.

[#] The one year ley will be referred to as "hay", to distinguish from the three year ley. The geological term "series" is used to denote the area under each crop in a rotation experiment. However, in this experiment the word "block" is used with the same meaning.

Crop Notes

Bf/1.2

First three years of each Rotation

Previous crop - Barley

Rotation 1. Ley

Grass Mixture

		lb. per acre	lb. per acre
Perennial Ryegrass		14	Wild White Clover 2
Cocksfoot		8	Italian Ryegrass (1941 10
Late Flowering Red Clover		4	and onwards only)

First year	Block	Sown	First year Grazed by sheep	Second year Grazed by sheep	Third year Grazed by sheep
1938	3	May 20	Aug. 3 - Sept. 14 2 grazings = 16 days	Apr. 13 - Oct. 17 9 grazings = 108 days	Apr. 27 - Oct. 16 4 grazings = 55 days
1939	5	May 4	Aug. 16 - Oct. 20 3 grazings = 28 days	Apr. 17 - Oct. 25 5 grazings = 119 days	May 5 - Oct. 26 4 grazings = 94 days
1940	4	May 6	Aug. 26 - Nov. 2 2 grazings = 30 days	May 5 - Oct. 18 4 grazings = 65 days	May 27 - Sept. 19 4 grazings = 38 days
1941	2	May 14	No grazing ¹	May 19 - Sept. 14 4 grazings = 46 days	Aug. 31 - Oct. 1 3 grazings = 17 days ²
1942	1	May 15	Aug. 24 - Sept. 8 1 grazing, on half plots	Sept. 8 ³ - Oct. 20 3 grazings = 21 days	June 21 ³ - Oct. 27 4 grazings = 46 days
1943	3	Apr. 28	Oct. 20 1 grazing ⁴	July 12 - Nov. 5 4 grazings = 44 days	May 26 - Sept. 29 4 grazings = 70 days
1944	5	May 8	July 28 - Nov. 13 4 grazings = 34 days	June 11 - Oct. 28 4 grazings = 62 days	May 9 - Nov. 19 6 grazings = 95 days
1945	4	May 7	Aug. 10 - Oct. 15 2 grazings = 41 days	May 25 - Nov. 3 5 grazings = 85 days	May 21 - Oct. 19 4 grazings = 48 days
1946	2	May 1	Aug. 15 - Nov. 10 3 grazings = 24 days	May 5 - Oct. 15 4 grazings = 51 days	
1947	1	May 1	Oct. 3 - 11 1 grazing = 9 days		

- Owing to an unfavourable season, and because the wrong quantity of manure was applied.
- Prior to the grazings there was one cut of hay on June 22.
- The sheep were put on late owing to war conditions.
- There was one cut, on July 15, to get rid of annual weeds.

Ley and Arable Rotations Experiment

Bf/1.3

Rotation 2. Lucerne

Variety: Provence 1939-1944, Grim 1945, Argentine 1946.

First year	Block	First year			Second year			Third year		
		1st Sown	2nd cut	1st cut	2nd cut	3rd cut	1st cut	2nd cut	3rd cut	
1938	3	May 20	Sept. 14	-	July 6	Aug. 18	Nov. 13	July 10	Sept. 4	Dec. 22
1939	5	May 4	Aug. 19	Nov. 13	July 10	Sept. 4	-	June 17	Aug. 12	Dec. 22
1940	4	May 6 ^x	Sept. 4	-	June 24	Aug. 12	Dec. 22	June 26	Sept. 16	-
1941	2	May 14	Aug. 12	-	June 26	Sept. 16	-	June 7	Aug. 31	-
1942	1	May 15	Sept. 16	-	June 7	Aug. 31	-	June 6	Aug. 8	-
1943	3	May 5	-	-	June 6	Sept. 4	-	June 8	Aug. 13	Nov. 12
1944	5	May 9	Sept. 12	-	June 8	Aug. 13	Nov. 12	June 25	Sept. 16	Dec. 10
1945	4	May 7	Aug. 13	Nov. 12	June 26	Sept. 16	-	June 11	July 31	Oct. 3
1946	2	May 1	Sept. 16	-	June 26	July 31	-			
1947	1	May 7	Oct. 3	-						

~~H~~ Bad patches reseeded June 11.

Rotations 3 and 4. First year - Potatoes, second year - Wheat, third year -
Hay (Rot.3) and Kale or Sugar Beet (Rot.4).

Varieties:

Potatoes - Majestic

Wheat = Red Standard

Kale - Thousand Head

Sugar Beet- Kleinwanzleben E.

First year	Block	1st year		Second year		Third year			Fourth year	
		Potatoes		Wheat		Hay (Rot.3)	1st	2nd	Kale (Rot.4)	Harvest-
		Harvest-		Harvest-		Sown	cut		Sown	ed
		Sown	ed	Sown	ed	Sown			Sown	ed
1938	3	Apr. 23	Sept. 22	31.10.38	Aug. 14	27.4.39	June 20		May 6	Dec. 30
1939	5	Apr. 25	Sept. 12	2.11.39	Aug. 12	3.5.40	June 17	Aug. 12	Apr. 29	Dec. 16
1940	4	Apr. 26	Sept. 20	16.10.40	Aug. 14	5.5.41	June 25	Sept. 16	Apr. 22	Dec. 29
1941	2	Apr. 24	Oct. 15	24.3.42*	Sept. 15	20.5.42	1 cut, date unknown		Crop failed	
1942	1	Apr. 21	Oct. 5	(Eaten by rats)		16.4.43	June 6	Sept. 12	May 9*	7.2.45
1943	3	May 13	Sept. 22	11.10.43	Aug. 10	9.5.44	June 8	Aug. 13	Sugar beet	
1944	5	Apr. 28	Sept. 20	May 10*	Aug. 10	10.5.45	June 24	Sept. 16	May 3	Nov. 26
1945	4	Apr. 27	Oct. 2	2.10.45	Aug. 22	19.5.46	June 11	July 31	May 2	Oct. 12
1946	2	Apr. 26	Oct. 8	6.11.46	Aug. 6				May 2	Oct. 17
1947	1	May 12	Oct. 10							

*Second sowing, first sowing failed.

Indicator Crops - Potatoes and Barley

Fourth and fifth years of rotations

Varieties: Potatoes - Majestic; Barley - Plumage Archer

Year	Block	Potatoes			Block	Barley	
		Sown	Harvested	Sown		Harvested	
1938	4	Prior to rotations	Apr. 22	Sept. 30	5	Mar. 9	Aug. 25 Aug. 25
1939	2		Apr. 24	Sept. 20	4	Mar. 2	Aug. 30
1940	1		Apr. 12	Sept. 25	2	Prior to rotations	Mar. 21 Aug. 30
1941	3	Fourth year of Cycle 1	Apr. 18	Nov. 7	1	Mar. 17	Aug. 27
1942	5		Apr. 17	Oct. 9	3	Fifth year of Cycle 1	Apr. 13 Aug. 19
1943	4		Apr. 14	Sept. 23	5	Mar. 4	Aug. 25
1944	2	rotations	Apr. 27	Oct. 2	4	Mar. 14	Aug. 30
1945	1		Apr. 26	Oct. 11	2	rotations	Mar. 2 Aug. 10
1946	3	Fourth year of Cycle 2	Apr. 29	Oct. 6	1	Mar. 20	Aug. 23
1947	5		Apr. 12	Oct. 2	3	Fifth year of Cycle 2	Apr. 17 Aug. 12

Rotation 1. Ley

Sheep-days of grazing per acre

First Cycle

Block 3			Block 5			Block 4		
1938	1939	1940	1939	1940	1941	1940	1941	1942
596	3667	2103	1135	1898	2062	645	1347	1344
Block 2			Block 1					
1941	1942	1943	1942	1943	1944			
None	1833	453	721	429	1050			

Second Cycle

Block 3			Block 5			Block 4		
1943	1944	1945	1944	1945	1946	1945	1946	1947
143	955	1910	764	1695	3101	1146	2787	1840
Block 2			Block 1					
1946	1947		1947					
811	1840		387					

Ley and Arable Rotations ExperimentRotation 2. Lucerne

Yields obtained during first three years of each rotation

Hay (85% dry matter), tons per acre

First Cycle

First Block year	First year	Second year				Third year						
		Crops:	1st	2nd	Total	1st	2nd	3rd	Total	1st	2nd	3rd
1938	3	0.61	-	0.61	0.81	0.54	0.14	1.49	1.86	0.78	-	2.64
1939	5	0.34	0.08	0.42	1.36	0.73	-	2.09	1.48	1.26	0.12	2.87
1940	4	1.05	-	1.05	1.51	0.97	0.11	2.59	2.03	0.83	-	2.86
1941	2	0.39	-	0.39	2.05	0.53	-	2.58	1.67	0.64	-	2.31
1942	1	0.32	-	0.32	1.29	0.77	-	2.06	1.96	1.58	-	3.54

Second Cycle

First year	Block	First year	Second year				Third year						
			1st	2nd	Total	1st	2nd	3rd	Total	1st	2nd	3rd	Total
1943	3	1st	2nd	Total	1st	2nd	3rd	Total	1st	2nd	3rd	Total	
No dung	Not cut				-	1.58	0.86	-	2.44	1.84	1.36	0.10	3.30
Dung (1941)					-	1.48	0.87	-	2.35	1.76	1.10	0.06	2.92
Mean Increase					-	1.53	0.86	-	2.39	1.80	1.23	0.08	3.11
Previous rotation:					-	-0.10	0.01	-	-0.09	-0.08	-0.26	-0.04	-0.38
Lucerne					-	1.42	0.77	-	2.19	1.68	1.22	0.09	2.99
Arable					-	1.64	0.96	-	2.60	1.92	1.23	0.07	3.22
1944	5												
No dung	0.94	-	0.94	1.43	1.24	0.20	2.87	2.26	2.08	0.04	4.38		
Dung (1942)	1.00	-	1.00	1.50	1.28	0.24	3.02	2.54	1.90	0.10	4.54		
Mean Increase	0.97	-	0.97	1.46	1.26	0.22	2.94	2.40	1.99	0.07	4.46		
0.06	-	0.06	0.07	0.04	0.04	0.15	0.28	-0.18	0.06	0.16			
Previous rotation:													
Lucerne	1.08	-	1.08	1.87	1.38	0.26	3.51	2.28	2.04	0.06	4.38		
Arable	0.85	-	0.85	1.06	1.14	0.18	2.38	2.52	1.94	0.08	4.54		
1945	4												
No dung	0.74	0.05	0.79	1.44	1.18	-	2.62	1.96	0.77	0.06	2.78		
Dung (1943)	0.70	0.04	0.74	1.76	1.25	-	3.01	2.04	1.02	0.04	3.10		
Mean Increase	0.72	0.04	0.76	1.60	1.22	-	2.82	2.00	0.89	0.05	2.94		
-0.04	-0.01	-0.05	0.32	0.07	-	0.39	0.08	0.26	-0.02	0.32			
Previous rotation:													
Lucerne	0.84	0.04	0.88	1.76	1.20	-	2.96	2.12	0.96	0.05	3.13		
Arable	0.60	0.06	0.66	1.44	1.24	-	2.68	1.87	0.83	0.06	2.75		

Lucerne Hay, tons per acre

Bf/1.6

First

Second Cycle (continued)

year Block

First year

Second year

Third year

1946	2	1st	2nd	Total	1st	2nd	3rd	Total	1st	2nd	3rd	Total
No dung		0.53	-	0.53	0.64	0.25	-	0.89				
Dung (1944)		0.51	-	0.51	0.72	0.34	-	1.06				
Mean		0.52	-	0.52	0.68	0.29	-	0.97				
Increase		-0.02	-	0.02	0.08	0.09	-	0.17				
Previous rotation:												
Lucerne		0.62	-	0.62	0.48	0.31	-	0.79				
Arable		0.42	-	0.42	0.88	0.28	-	1.16				
1947	1											
No dung		0.10	-	0.10								
Dung (1945)		0.07	-	0.07								
Mean		0.09	-	0.09								
Increase		-0.03	-	-0.03								
Previous rotation:												
Lucerne		0.09	-	0.09								
Arable		0.08	-	0.08								

Ley and Arable Rotations ExperimentRotations 3 and 4First year - Potatoes, total tubers in tons per acre; percentage ware

	Total tubers	% ware	Total tubers	% ware	Total tubers	% ware	Total tubers	% ware
	<u>1938 - Block 3</u>		<u>1939 - Block 5</u>		<u>1940 - Block 4</u>		<u>1941 - Block 2</u>	
No dung					9.71	87.0	12.06	82.3
Dung [#]					11.50	89.1	12.19	80.6
Mean Increase	12.77	92.2	11.13	85.4	10.61	88.0	12.12	81.4
	-	-	-	-	1.79	2.1	0.13	-1.7
(Second cycle)								
	<u>1942 - Block 1</u>		<u>1943 - Block 3</u>		<u>1944 - Block 5</u>		<u>1945 - Block 4</u>	
No dung	9.24	75.8	7.76	75.7	12.18	86.0	11.71	86.4
Dung [#]	9.59	75.5	9.53	80.7	13.33	86.0	13.28	88.9
Mean Increase	9.41	75.6	8.64	78.2	12.75	86.0	12.50	87.6
	0.35	-0.3	1.77	5.0	1.15	0.0	1.57	2.5
Previous rotation:								
Ley			10.48	82.6	12.54	86.8	11.66	88.6
Lucerne			8.28	76.8	13.26	86.1	13.82	87.6
Arable			8.74	82.4	12.75	85.1	12.56	89.5
With hay								
Arable			7.08	70.9	12.46	86.2	11.94	85.2
With kale								
	<u>1946 - Block 2</u>		<u>1947 - Block 1</u>					
No dung	8.04	87.2	6.11	91.9				
Dung [#]	9.13	87.6	6.24	90.4				
Mean Increase	8.58	87.4	6.18	91.2				
	1.09	0.4	0.13	-1.5				
Previous rotation:								
Ley	10.21	89.2	6.90	91.6				
Lucerne	8.69	89.2	6.67	92.0				
Arable								
With hay	7.32	86.8	5.32	91.0				
Arable								
With kale	8.12	84.3	5.81	90.1				

[#]Dung was applied two years previously, e.g. in 1938 on Block 4.

Rotations 3 and 4

Second year - Wheat, grain and straw in cwt. per acre

	Grain	Straw	Grain	Straw	Grain	Straw	Grain	Straw
First Cycle								
	<u>1939 - Block 3</u>		<u>1940 - Block 5</u>		<u>1941 - Block 4</u>		<u>1942 - Block 2</u>	
No dung					8.7	14.3		
Dung					9.5	16.2	Crop complete failure	
Mean Increase	12.9	22.5	15.2	24.2	9.1	15.2		
	-	-	-	-	0.8	1.9		
Second cycle								
	<u>1944 - Block 3</u>		<u>1945 - Block 5</u>		<u>1946 - Block 4</u>		<u>1947 - Block 2</u>	
No dung	9.1	23.3	13.7	30.2	10.8	18.0	7.7	14.8
Dung	11.3	27.8	14.0	25.8	12.6	24.2	8.8	14.8
Mean Increase	10.2	25.6	13.8	28.0	11.7	21.1	8.2	14.8
	2.2	4.5	0.3	-4.4	1.8	6.2	1.1	0.0
Previous rotation:								
Ley	14.1	32.1	15.2	33.4	12.1	21.5	9.6	18.4
Lucerne	10.6	24.3	15.0	26.0	12.7	21.9	8.6	13.5
Arable with hay	11.2	27.6	11.0	22.2	8.6	14.2	5.5	10.5
Arable with kale	5.0	18.1	14.2	30.4	13.3	26.8	9.3	16.8

* The dung was applied three years previously, e.g. in 1938 on block 4, etc.

Ley and Arable Rotations ExperimentRotation 3Third year - Hay, yield (85% dry matter) in tons per acre

	First Crop	Second Crop	Total	First Crop	Second Crop	Total	First Crop	Second Crop	Total
First cycle									
	<u>1940 - Block 3</u>			<u>1941 - Block 5</u>					
	0.78	-	0.78	1.31	0.33	1.64			
Second cycle									
	<u>1942 - Block 4</u>			<u>1943 - Block 2</u>			<u>1944 - Block 1</u>		
No dung	1.85	0.22	2.07	1.04	-	1.04	1.02	0.37	1.38
Dung	2.10	0.21	2.31	1.12	-	1.12	1.02	0.34	1.36
Mean Increase	1.98	0.21	2.19	1.08	-	1.08	1.02	0.35	1.37
	0.25	-0.01	0.24	0.08	-	0.08	-	-0.03	-0.02
Previous rotation:									
Ley	1.27	0.30	1.57				2.18	0.40	2.59
Lucerne				2.86	1.20	4.06			
Arable				0.16	0.04	0.20	0.13	-0.09	0.05
with hay	1.68	0.42	2.10	2.58	1.07	3.65	1.46	0.14	1.60

¹⁵Dung was applied four years previously, e.g. in 1938 on Block 4 etc.

Rotation 4Third year - Kale, in tons per acre, Cycle 1 or

Sugar Beet, clean beet and tops in tons
per acre, total sugar in cwt.
per acre and sugar percentage,
Cycle 2.

Cycle 1. Kale

Block	1940 3	1941 5	1942 4	1943 2	1944 1	
No dung Dung			7.15 7.62	Crop failed	4.94 4.27	
Mean Increase	7.34	9.35	7.38 0.47		4.60 -0.67	

Cycle 2. Sugar Beet.

	1945 - Block 3				1946 - Block 5			
	Clean beet	Tops	Total Sugar	Sugar %	Clean beet	Tops	Total sugar	Sugar %
No dung Dung	6.2 7.9	2.8 3.4	21.9 27.2	17.47 17.20	7.29 8.12	7.29 7.45	27.9 31.7	19.14 19.53
Mean Increase	7.0 1.7	3.1 0.6	24.6 5.3	17.34 -0.27	7.70 0.83	7.37 0.16	29.8 3.8	19.34 0.39
Previous rotation:								
Ley								
Lucerne	7.2	3.0	25.4	17.60	8.16	7.88	31.2	19.10
Arable with kale	6.9	3.1	23.6	17.08	7.24	6.86	28.3	19.57

^HDung was applied four years previously, e.g. in 1938 on Block 4, etc.

1947 - Block 4

No dung	4.97	2.88	20.7	20.82
Dung	8.63	4.82	35.4	20.52
Mean	6.80	3.85	28.1	20.67
Increase	3.66	1.94	14.7	-0.30
Previous rotation:				
Ley				
Lucerne	7.42	4.21	30.9	20.82
Arable with kale	6.18	3.49	25.3	20.52

Ley and Arable Rotations Experiment

Yields of Indicator Crops

Potatoes - effect of dung in current year

Total tubers, tons per acre

Percentage ware

Prior to rotations

1938 - Block 4		1939 - Block 2		(1938)	(1939)
No dung	11.41	10.89			85.2
Dung	15.47	12.52			88.6
Mean	13.44	11.69		89.2	86.9
Increase	4.06 (± 0.32)	1.63 (± 0.11)			3.4 (± 1.45)

1940 - Block 1

	After Hay	After Kale	Mean		After Hay	After Kale	Mean
		$\pm 0.64^1$				$\pm 2.0^1$	
No dung	8.23	6.75	7.49		82.0	76.6	79.3
Dung	9.83	7.85	8.84		82.4	79.5	81.0
Mean	± 0.59	9.02	7.30	8.16	± 1.70	82.2	78.0
Resp.	± 0.50	1.60	1.10	1.35	± 2.36	0.4	2.9
			± 0.36				± 1.67

Fourth year of cycles

Previous crop rotation				
	Arable			
	Lu-	with		
	Ley	Cerne	Hay	Kale
1941 - Block 3			$\pm 0.45^1$	
No dung	15.19	11.15	11.82	9.35
Dung	14.38	14.23	14.40	12.19
Mean	± 0.41	14.79	12.69	13.11
Resp.	± 0.37	-0.80	3.08	2.58
			2.84	1.93
				± 0.18
1942 - Block 5			$\pm 0.37^1$	
No dung	13.70	13.22	12.52	10.83
Dung	14.66	15.40	14.75	13.14
Mean	± 0.35	14.18	14.31	13.64
Resp.	± 0.26	9.96	2.18	2.23
			2.31	1.92
				± 0.13

Previous crop rotation

Arable				
	Lu-	with		
	Ley	Cerne	Hay	Kale
			$\pm 1.47^1$	
			87.6	84.1
			83.4	82.2
			86.2	85.0
			84.2	86.6
				± 0.55
			± 1.32	86.9
			84.6	83.8
			83.8	84.4
			84.4	84.9
			± 1.27	-1.4
			0.9	0.8
			4.4	1.2
				± 0.64
			$\pm 1.16^1$	
			87.0	85.0
			85.0	81.4
			81.4	84.6
			± 1.27	86.1
			87.2	84.2
			84.2	86.2
				± 0.64
			± 0.97	87.2
			85.6	86.1
			86.1	82.8
			82.8	85.4
			± 1.27	0.4
			1.1	2.2
			2.2	2.8
			2.8	1.6
				± 0.64

Standard errors (1) for comparisons other than vertical ones.

Potatoes - effect of dung in current year.

Fourth year of cycles (continued)

Total tubers, tons per acre					Percentage ware						
Previous crop rotation					Previous crop rotation						
Arable					Arable						
Ley	Lu- cerne	with Hay	Kale	Mean	Ley	Lu- cerne	with Hay	Kale	Mean		
<u>1943 - Block 4</u>					<u>±2.60¹</u>						
No dung	8.84	9.76	8.08	7.43	8.53	80.4	79.8	76.6	71.4	77.0	
Dung	10.31	13.52	9.50	10.63	10.99	81.3	87.1	83.6	81.4	83.4	
Mean <u>±0.40</u>	9.58	11.64	8.79	9.03	9.76	<u>±1.14</u>	80.8	83.4	80.1	76.4	80.2
Resp. <u>±1.11</u>	1.47	3.76	1.42	3.20	2.46	<u>±4.67</u>	0.9	7.3	7.0	10.0	6.4
					<u>±0.56</u>					<u>±2.34</u>	
<u>1944 - Block 2</u>					<u>±0.91¹</u>						
No dung	12.08	13.20	10.56	11.22	11.76	89.4	87.5	86.2	82.2	86.3	
Dung	13.04	14.64	11.76	12.24	12.92	89.4	89.0	88.1	85.5	88.0	
Mean <u>±0.57¹</u>	12.56	13.92	11.16	11.73	12.34	<u>±0.82</u>	89.4	88.2	87.2	83.8	87.2
Resp. <u>±0.45</u>	0.96	1.44	1.20	1.02	1.16	<u>±0.76</u>	0.0	1.5	1.9	3.3	1.7
					<u>±0.22</u>					<u>±0.38</u>	
<u>1945 - Block 1</u>					<u>±1.55¹</u>						
No dung	13.28	12.96	12.46	11.06	12.44	87.0	86.8	84.7	85.0	85.9	
Dung	13.32	12.68	11.89	12.48	12.59	86.8	85.0	85.8	83.2	85.2	
Mean <u>±0.39</u>	13.30	12.82	12.18	11.77	12.52	<u>±0.71</u>	86.9	85.9	85.2	84.1	85.6
Resp. <u>±1.35</u>	0.04	-0.28	-0.57	1.42	0.15	<u>±2.76</u>	-0.2	-1.8	1.1	-1.8	-0.7
					<u>±0.68</u>					<u>±1.38</u>	
<u>1946 - Block 3</u>					<u>±0.52¹</u>						
No dung	11.37	9.59	8.94	8.32	9.56	90.2	89.1	87.6	89.7	89.2	
Dung	13.28	11.50	11.93	8.88	11.40	90.6	92.3	90.8	93.2	91.7	
Mean <u>±0.86</u>	12.32	10.54	10.44	8.60	10.48	<u>±0.47</u>	90.4	90.7	89.2	91.4	90.4
Resp. <u>±0.69</u>	1.91	1.91	2.99	0.56	1.84	<u>±0.46</u>	0.4	3.2	3.2	3.5	2.5
					<u>±0.35</u>					<u>±0.23</u>	
<u>1947 - Block 5</u>					<u>±0.64¹</u>						
No dung	8.42	8.74	6.28	8.03	7.87	93.3	92.8	92.4	93.2	92.9	
Dung	9.06	9.62	8.11	8.86	8.91	94.1	93.2	93.2	94.0	93.6	
Mean <u>±0.35</u>	8.74	9.18	7.20	8.44	8.39	<u>±0.53</u>	93.7	93.0	92.8	93.6	93.3
Resp. <u>±0.28</u>	0.64	0.88	1.83	0.83	1.04	<u>±0.70</u>	0.8	0.4	0.8	0.8	0.7
					<u>±0.14</u>					<u>±0.35</u>	

Standard errors (1) for comparisons other than vertical ones.

In 1946 and 1947 read "Arable with sugar beet" for "Arable with kale".

Ley and Arable Rotations Experiment

Potatoes

Standard errors per plot

			Total tubers			Percentage ware
			d.f.	tons per acre	% of mean	
1938 - Block 4	Whole plot	7	0.68	5.08		
	Sub-plot	7	0.64	4.79		
1939 - Block 2	Whole plot	7	1.00	8.52		1.49
	Sub-plot	7	0.22	1.86		2.90
1940 - Block 1	Whole plot	6	1.18	14.5		3.40
	Sub-plot	6	0.72	8.75		3.34

Fourth year of cycles

1941 - Block 3	Whole plot	4	0.57	4.47	1.87
	Sub-plot	4	0.37	2.86	1.27
1942 - Block 5	Whole plot	4	0.49	3.62	1.38
	Sub-plot	4	0.26	1.95	1.27
1943 - Block 4	Whole plot	4	0.57	5.82	1.61
	Sub-plot	4	1.11	11.3	4.67
1944 - Block 2	Whole plot	4	0.80	6.51	1.17
	Sub-plot	4	0.45	3.65	0.76
1945 - Block 1	Whole plot	4	0.55	4.40	1.00
	Sub-plot	4	1.35	10.8	2.76
1946 - Block 3	Whole plot	4	1.21	11.5	0.66
	Sub-plot	4	0.69	6.6	0.46
1947 - Block 5	Whole plot	4	0.50	6.0	0.75
	Sub-plot	4	0.28	3.3	0.70

Barley - residual effect of dung applied to potatoes in the previous year.

Grain, cwt. per acre.			Straw, cwt. per acre.	
<u>1938 - Block 5</u>		<u>1939 - Block 4</u>	(1938)	(1939)
No dung		15.3		23.1
Dung		16.2		28.6
Mean Response	7.66	15.8 0.9 (± 0.22)	12.6	25.8 (± 1.08) 5.5
<u>1940 - Block 2</u>				
No dung	6.4		7.9	
Dung	8.5		11.4	
Mean Response	7.5 2.1	(± 0.80)	9.7 3.5	(± 1.02)
<u>1941 - Block 1</u>				
Crop previous to potatoes				
	Hay	Kale	Mean	
	$\pm 1.16^1$			$\pm 1.28^1$
No dung	7.0	10.6	8.8	10.0
Dung	7.6	9.7	8.6	10.1
Mean ± 0.87	7.3	10.1	8.7	13.2
Resp. ± 1.56	0.6	-0.9	-0.2	11.6
			± 1.03	12.2
			± 1.52	
			0.1	0.6
			1.0	± 1.07

Standard errors (1) for comparisons other than vertical ones

Ley and Arable Rotations Experiment

Barley - residual effect of dung applied to potatoes in previous year.

Grain, cwt. per acre

Straw, cwt. per acre

Fifth years of cycles

Crop previous to potatoes

Crop previous to potatoes

Arable Lu- with Ley cerne Hay Kale Mean					Arable Lu- with Ley cerne Hay Kale Mean							
<u>1942 - Block 3</u> <u>$\pm 2.27^1$</u>					<u>$\pm 2.46^1$</u>							
No dung	18.6	22.6	13.3	9.5	16.0	21.7	26.2	16.3	12.1	19.1		
Dung	21.8	21.3	18.1	17.8	19.8	25.0	25.7	21.7	20.7	23.3		
Mean	<u>± 1.42</u>	20.2	21.9	15.7	13.7	17.9	<u>± 1.82</u>	23.4	26.0	19.0	16.4	21.2
Resp.	<u>± 3.55</u>	3.2	-1.3	4.8	8.3	3.8	<u>± 3.31</u>	3.3	-0.5	5.4	8.6	4.2
						<u>± 1.78</u>						<u>± 1.66</u>
<u>1943 - Block 5</u> <u>$\pm 2.96^1$</u>					<u>$\pm 3.34^1$</u>							
No dung	13.2	14.9	14.0	22.7	16.2	23.8	24.1	20.7	34.7	25.8		
Dung	12.8	14.5	17.4	16.2	15.2	19.2	22.8	26.7	23.6	23.1		
Mean	<u>± 2.28</u>	13.0	14.7	15.7	19.4	15.7	<u>± 2.19</u>	21.5	23.4	23.7	29.2	24.4
Resp.	<u>± 3.79</u>	-0.4	-0.4	3.4	-6.5	-1.0	<u>± 5.05</u>	-4.6	-1.3	6.0	-11.1	-2.7
						<u>± 1.90</u>						<u>± 2.52</u>
<u>1944 - Block 4</u> <u>$\pm 3.24^1$</u>					<u>$\pm 4.38^1$</u>							
No dung	18.1	22.5	19.0	24.2	21.0	22.7	20.1	21.8	23.2	21.9		
Dung	15.7	19.7	23.0	17.4	18.9	19.2	23.5	29.2	20.8	23.2		
Mean	<u>± 2.21</u>	16.9	21.1	21.0	20.8	20.0	<u>± 2.78</u>	20.9	21.8	25.5	22.0	22.6
Resp.	<u>± 4.76</u>	-2.4	-2.8	4.0	-6.8	-2.1	<u>± 6.77</u>	-3.5	3.4	7.4	-2.4	1.3
						<u>± 2.38</u>						<u>± 3.38</u>
<u>1945 - Block 2</u> <u>$\pm 2.07^1$</u>					<u>$\pm 1.80^1$</u>							
No dung	13.2	17.6	11.6	13.7	14.0	15.3	18.9	11.9	14.8	15.2		
Dung	11.1	20.0	14.5	15.4	15.2	13.8	20.9	16.8	17.8	17.3		
Mean	<u>± 1.17</u>	12.2	18.8	13.0	14.6	14.6	14.6	19.9	14.4	16.3	16.2	
Resp.	<u>± 3.43</u>	-2.1	2.4	2.9	1.7	1.2	-1.5	2.0	4.9	3.0	2.1	
						<u>± 1.72</u>						
<u>1946 - Block 1</u> <u>$\pm 1.44^1$</u>					<u>$\pm 1.80^1$</u>							
No dung	17.4	18.8	14.0	14.1	16.1	20.4	23.7	17.0	19.2	20.1		
Dung	16.4	16.4	15.6	15.7	16.0	18.5	20.1	19.6	19.8	19.5		
Mean	<u>± 0.80</u>	16.9	17.6	14.8	14.9	16.0	<u>± 0.44</u>	19.4	21.9	18.3	19.5	19.8
Resp.	<u>± 2.40</u>	-1.0	-2.4	1.6	1.6	-0.1	<u>± 3.48</u>	-1.9	-3.6	2.6	0.6	-0.6
						<u>± 1.20</u>						<u>± 1.74</u>

Standard errors (1) for comparisons other than vertical ones.

Bf/1.16

Barley - residual effect of dung applied to potatoes in the previous year.

Grain, cwt. per acre

Straw, cwt. per acre

Fifth years of cycles (continued)

Crop previous to potatoes

Crop previous to potatoes

	Arable					Mean	Arable					Mean
	Ley	Lu- cerne	with Hay	Sugar Beet			Ley	Lu- cerne	with Hay	Sugar Beet		
<u>1947 - Block 3</u>												$\pm 1.21^1$
No dung	15.5	12.5	12.3	10.4	12.7		15.8	15.4	13.3	14.9	14.8	
Dung	18.7	16.5	15.6	12.1	15.7		21.0	20.3	18.6	17.3	19.3	
Mean ± 0.99	17.1	14.5	14.0	11.2	14.2		18.4	17.8	16.0	16.1	17.1	
Resp. ± 1.39	3.2	4.0	3.3	1.7	3.0		5.2	4.9	5.3	2.4	4.5	
					± 0.69							

Standard errors (1) for comparisons other than vertical ones

Ley and Arable Rotations Experiment

Barley

Standard errors per plot

		Barley - Grain			Barley - Straw		
		cwt.	per	% of	cwt.	per	% of
		d.f.	acre	mean	d.f.	acre	mean
1938 - Block 5	Whole plot	7	1.79	23.4	7	2.62	20.8
	Sub-plot	8	1.67	21.8	8	1.50	11.9
1939 - Block 4	Whole plot	7	1.55	9.85	7	1.29	4.98
	Sub-plot	6 ^x	0.44	2.76	6 ^x	2.17	8.39
1940 - Block 2	Whole plot	7	2.34	31.5	7	2.35	24.3
	Sub-plot	7	1.60	21.5	7	2.04	21.1
1941 - Block 1	Whole plot	6	1.74	20.0	6	2.06	17.4
	Sub-plot	6	2.21	25.4	6	2.15	18.2
<u>Fifth year of cycles</u>							
1942 - Block 3	Whole plot	4	2.01	11.2	4	2.57	12.1
	Sub-plot	4	3.55	19.9	4	3.31	15.6
1943 - Block 5	Whole plot	4	3.22	20.5	4	3.09	12.7
	Sub-plot	4	3.79	24.1	4	5.05	20.6
1944 - Block 4	Whole plot	4	3.12	15.6	4	3.93	17.4
	Sub-plot	4	4.76	23.9	4	6.77	30.0
1945 - Block 2	Whole plot	4	1.65	11.3			
	Sub-plot	4	3.43	23.4			
1946 - Block 3	Whole plot	4	1.13	7.0	4	0.62	3.15
	Sub-plot	4	2.40	15.0	4	3.48	17.6
1947 - Block 3	Whole plot	4	1.39	9.8			
	Sub-plot	4	1.39	9.8			

^x One missing sub-plot.

MARKET GARDEN EXPERIMENT

Woburn, Lansome Field (begun in 1942)

The use of heavy dressings of organic manures for making a market garden soil, and the effect of sulphate of ammonia.

The four crops follow a two-year rotation, each of the two "series" into which the area is divided bearing two crops in a year, while the series carry different crops at any one season, except that in the first year the whole area was under winter cabbage.

System of replication: 2 series, each consisting of 4 randomized blocks of 10 plots each, certain interactions being partially confounded with block differences.

Area of each plot: 0.0125 acre. (In 1943, peas and beet 0.0105 acre)

Cropping:

- | | |
|----------|--|
| 1st year | Globe beet (sown April, lifted July)
Winter cabbage (transplanted August, cut December-March) |
| 2nd year | Peas (sown March-April, pulled June-July)
Leeks (transplanted July, lifted January-March) |

Treatments

Organic manures: Dung, sewage sludge compost (composted town refuse in 1942 and 1943), sewage sludge (West Middlesex) and vegetable compost, each at 15 and 30 tons per acre.

Organics were applied at 4 and 8 tons per acre to winter cabbage in the first year.

Sulphate of ammonia:

With organic manures: None, 0.6 cwt. N per acre

In absence of organics: None, 0.6, 1.2, 1.8 cwt. N per acre

Basal manuring:

Superphosphate, 0.4 cwt. P₂O₅ per acre (triple superphosphate in 1945 and 1946).

Muriate of potash, 0.5 cwt. K₂O per acre.

In 1943, 3200 lb. per acre carbonate of lime was applied to winter cabbage.

Time of application of manures:

Organic manures and basal dressings are given in a single dose in early spring before sowing peas and beet, except that in the first year the winter cabbage received a reduced dressing of organics and the full amount of basal manures. The sulphate of ammonia is divided between crops as follows (cwt. N per acre):

	With organics	In the absence of organics
Globe beet and peas	0, 0.2	0, 0.2, 0.4, 0.6
Cabbage and leeks	0, 0.4	0, 0.4, 0.8, 1.2

In the first year, the winter cabbage received the whole dressing of sulphate of ammonia.

Crop Notes

Winter Cabbage

Previous crop: Globe Beet (Cabbage in 1942)

Year	Series	Variety	Planted out	Harvested
1942	A	Christmas Cabbage	Aug. 13	Nov. 20 - Feb. 17
1943	B	January King	Sept. 2	Dec. 2 - March 16
1944	A	January King	Aug. 23	March 14, April 9
1945	B	January King	July 24 and Aug. 10	Dec. 1 - March 6
1946	A	Christmas Drumhead and Savoy	Aug. 12	Nov. 11 - 15
1947	B	Failed, owing to dry weather.		

Leeks (Cabbage in 1942)

Previous crop: Peas (Cabbage in 1942)

1942	B	January King ¹	Sept. 21	March 19
1943	A	Musselburgh	Aug. 6 - 20	March 6
1944	B	Musselburgh	July 18, 28	Feb. 22 - April 13
1945	A	Musselburgh	Aug. 7-14	March 5-12
1946	B	Musselburgh	Aug. 10, 30	May 15-20
1947	A	Musselburgh	July 18	Feb. 12-27

Globe Beet. Variety: Crimson Globe

Previous crop: Leeks (Cabbage in 1942)

Year	Series	Sown	Harvested
1943	B	April 27	July 19 - August 20
1944	A	April 16	July 24 - August 14
1945	B	April 20	July 5 - 30
1946	A	April 3	July 24 - August 6
1947	B	May 27	August 8 - 13

Peas. Variety: Kelvedon Wonder

Previous crop: Winter Cabbage

1943	A	May 4 ²	July 12 - 14
1944	B	April 11	July 5 - 11
1945	A	April 4, 18	June 25 - July 17
1946	B	March 25, May 13 ³	July 6, 23 - 30
1947	A	April 25	June 26 - 30

(1) No leek plants available

(2) First sowing, April 10, failed

(3) Two sowings, March 15 and April 20, failed

Market Garden Experiment

Bg/1.3

Globe Beet and Peas

	Sulphate of Ammonia, cwt. N per acre			None 0.2 Mean			None 0.2 Mean			None 0.2 Mean		
	None	0.2	Mean	None	0.2	Mean	None	0.2	Mean	None	0.2	Mean
Organic Manures t.p.a.	Globe Beet Total produce ¹ : tons/acre			Globe Beet Bulbs ² : tons/acre			Peas Green peas: cwt/acre					
1943	(Series B)			(Series A)								
O	2.65	3.53	3.09	±1.246	±0.881		±4.37	±3.09				
D 15	7.56	3.56	5.56				22.1	16.7	19.4			
D 30	8.06	8.80	8.43				30.4	28.9	29.7			
CTR 15	4.91	5.81	5.36				15.4	27.7	21.5			
CTR 30	5.43	7.86	6.64				24.3	17.7	21.0			
SS 15	5.29	5.11	5.20				21.6	30.1	25.8			
SS 30	6.98	6.66	6.82				20.1	28.3	24.2			
VC 15	5.46	6.92	6.19				31.8	26.9	29.3			
VC 30	8.34	7.24	7.79				15.4	23.8	19.6			
Mean	6.50 ³	6.49 ³	5.86 ⁴				29.4	33.0	31.2			
S.E. per plot	1.763 or 30.1%			23.5 ³ 27.0 ³ 24.0 ⁴			6.18 or 25.8%					
1944	(Series A)			(Series B)								
O	1.33	1.10	1.21	±0.775	±0.548		±2.61	±1.84				
D 15	4.15	4.26	4.21				11.2	7.1	9.1			
D 30	3.07	5.77	4.42				9.3	8.0	8.7			
CSS 15	2.71	1.51	2.11				8.3	14.2	11.2			
CSS 30	1.96	5.01	3.48				10.6	9.9	10.2			
SS 15	2.94	3.78	3.36				10.4	11.2	10.8			
SS 30	4.90	4.13	4.52				11.2	10.8	11.0			
VC 15	1.21	2.99	2.10				7.3	8.9	8.1			
VC 30	4.68	5.76	5.22				4.9	13.6	9.3			
Mean	3.20 ³	4.15 ³	3.22 ⁴				9.8	11.5	10.6			
S.E. per plot	1.09% or 33.0%			9.0 ³ 11.0 ³ 10.2 ⁴			3.68 or 36.1%					

(1) Excludes totally unmarketable produce and includes tops

(3) Excludes "No organic manure".

(4) Mean of all plots.

All standard errors have 17 d.f.

Symbols: O No organic manure CTR Composted town refuse
 D Dung CSS Composted sewage sludge
 SS Sewage sludge VC Vegetable compost

Globe Beet and Peas

	Sulphate of Ammonia, cwt. N per acre								
	None	0.2	Mean	None	0.2	Mean	None	0.2	Mean
Organic Manures t.p.a.	Globe Beet Total produce ¹ : tons/acre			Globe Beet Bulbs: tons/acre			Peas Green peas: cwt/acre		
<u>1945</u>		(Series B)			(Series B)			(Series A)	
		±0.748	±0.529		±0.457	±0.323		±7.79	±5.51
0	3.03	4.91	3.97	1.51	2.73	2.12	48.3	37.8	43.0
D	15	6.28	5.79	6.03	3.51	3.32	3.41	46.2	42.9
D	30	6.24	7.02	6.63	3.45	3.63	3.54	38.0	47.8
CSS	15	6.12	4.55	5.33	3.41	2.85	3.13	46.4	32.7
CSS	30	5.30	6.55	5.93	2.91	3.52	3.22	43.6	50.0
SS	15	5.40	5.95	5.68	2.87	3.29	3.08	41.8	49.9
SS	30	7.20	5.67	6.44	3.81	3.18	3.50	44.3	49.2
VC	15	5.80	5.13	5.46	3.25	3.16	3.21	46.5	38.7
VC	30	7.46	7.11	7.28	4.13	3.95	4.04	50.4	50.9
Mean		6.23 ³	5.97 ³	5.68 ⁴	3.42 ³	3.36 ³	3.15 ⁴	44.6 ³	45.2 ³
S.E. per plot		1.058 or 18.6%			0.646 or 20.5%			11.02 or 25.1%	
<u>1946</u>		(Series A)			(Series B)				
		±0.648	±0.458		±8.61	±6.09			
0		0.51	0.88	0.69	20.2	37.6	28.9		
D	15	2.32	2.77	2.55	35.5	31.6	33.6		
D	30	4.01	3.64	3.82	40.2	36.6	38.4		
CSS	15	0.97	1.44	1.21	38.6	36.8	37.7		
CSS	30	1.95	1.63	1.79	33.6	18.2	25.9		
SS	15	1.22	1.23	1.22	33.1	37.7	35.4		
SS	30	1.32	2.32	1.82	36.6	30.2	33.4		
VC	15	1.31	1.55	1.43	54.0	40.1	47.0		
VC	30	3.01	2.90	2.96	32.1	30.4	31.2		
Mean		2.01 ³	2.18 ³	1.84 ⁴	38.0 ³	32.7 ³	33.7 ⁴		
S.E. per plot		0.916 or 49.7%			12.18 or 36.1%				

(1) Excludes totally unmarketable produce and includes tops.

In 1946 many plants went to seed.

(2) Excludes unmarketable produce.

(3) Excludes "No organic manure".

(4) Mean of all plots.

All standard errors have 17 d.f.

Globe Beet and Peas

	Sulphate of Ammonia, cwt. N per acre			Sulphate of Ammonia, cwt. N per acre			Sulphate of Ammonia, cwt. N per acre		
	None	0.2	Mean	None	0.2	Mean	None	0.2	Mean
Organic Manures t.p.a.	Globe Beet Total produce ¹ : tons/acre			Globe Beet Bulbs ² : tons/acre			Peas Green peas cwt/acre		
1947	(Series B)			(Series B)			(Series A)		
O	1.43	0.86	1.14	0.87	0.46	0.66	22.3	17.1	19.7
D 15	3.35	2.90	3.12	1.57	1.52	1.54	27.4	25.7	26.6
D 30	5.52	5.25	5.39	3.02	3.04	3.03	26.9	27.0	26.9
CSS 15	1.67	2.63	2.15	0.97	1.38	1.18	28.6	25.8	27.2
CSS 30	3.41	2.53	2.97	1.91	1.24	1.57	26.4	23.4	24.9
SS 15	0.92	1.12	1.02	0.54	0.63	0.58	25.6	21.0	23.3
SS 30	0.60	1.12	0.86	0.20	0.62	0.41	25.0	27.0	26.0
VC 15	2.61	3.18	2.90	1.51	1.74	1.63	23.2	17.9	20.5
VC 30	3.19	3.23	3.21	1.87	1.75	1.81	27.8	26.2	27.0
Mean	2.66 ³	2.74 ³	2.34 ⁴	1.45 ³	1.49 ³	1.28 ⁴	26.4 ³	24.2 ³	24.3 ⁴
S.E. per plot	0.736 or 31.4%			0.438 or 34.2%			3.83 or 15.8%		

(1) Excludes totally unmarketable produce and includes tops.

(2) Excludes unmarketable produce

(3) Excludes "No organic manure"

(4) Mean of all plots

All standard errors have 17 d.f.

Yields on plots without organic Manure

Sulphate of Ammonia, cwt. N per acre

	None	0.2	0.4	0.6	S.E.
Globe Beet					
Total produce: tons per acre					
1943	2.65	3.53	2.78	4.16	±1.246
1944	1.33	1.10	1.89	3.25	±0.775
1945	3.03	4.91	3.70	4.35	±0.748
1947	1.43	0.86	0.48	0.89	±0.520
Bulbs: tons per acre					
1945	1.51	2.73	2.17	2.31	±0.457
1946	0.51	0.88	1.35	0.51	±0.648
1947	0.87	0.46	0.29	0.47	±0.310
Peas					
Green peas: cwt. per acre					
1943	22.1	16.7	16.7	19.9	±4.37
1944	11.2	7.1	10.1	16.2	±2.61
1945	48.3	37.8	39.6	33.4	±7.79
1946	20.2	37.6	28.4	23.0	±8.61
1947	22.3	17.1	20.4	21.4	±2.71

Winter Cabbage

	Sulphate of Ammonia, cwt.N per acre			cwt.N per acre			cwt.N per acre		
	None	0.6	Mean	None	0.6	Mean	None	0.6	Mean
Organic Manures t.p.a. ¹	Total produce: tons/acre			Plant number thousands per acre			Total produce tons/acre		
1942	(Series A)			(Series A)			(Series B)		
O	5.27	7.68	6.48	17.5	18.2	17.9	1.30	2.51	1.91
D	5.96 ^a	7.91 ^a	6.94 ^b	17.6 ^c	18.5 ^c	18.0 ^d	2.33 ^e	2.19 ^e	2.26 ^f
D	7.63 ^a	8.90 ^a	8.27 ^b	18.1 ^c	18.4	18.3	2.30 ^e	2.66 ^e	2.48 ^f
CTR	7.66	7.15	7.40	18.3	18.6	18.4	2.16	2.39	2.27
CTR	7.43	9.79	8.61	18.2	19.4	18.8	2.16	2.29	2.22
SS	8.75	9.60	9.18	18.8	19.1	19.0	2.14	2.43	2.28
SS	8.55	7.48	8.02	16.7	17.2	17.0	1.99	2.50	2.24
Mean	7.45 ²	8.46 ²	7.69 ³	17.9 ²	18.5 ²	18.2 ³	2.21 ²	2.41 ²	2.23 ³
S.E. per plot	1.479 or 19.2%			1.16 or 6.4%			0.424 or 19.0%		

Standard errors for D figures only: (a) 0.740 (b) 0.523 (c) 0.58 (d) 0.41
 (e) 0.212 (f) 0.150

- (1) Since there was no crop of globe beet and peas, organics and all the Sulphate of Ammonia (0.6 cwt. N per acre) were applied to the cabbages. No vegetable compost was available: instead dung was applied, so that the number of dunged plots was twice that in subsequent years.
- (2) Excludes "No organic manure".
- (3) Mean of all plots.

All standard errors have 17 d.f.

Symbols: O No organic manure. SS Sewage sludge
 D Dung VC Vegetable compost
 CTR Composted town refuse

Market Garden Experiment

Bg/1.7

Winter Cabbage and Leeks

Organic Manures t.p.a. ⁴	Winter Cabbage (Series B)				Leeks (Series A)			
	None	0.4	Mean	Sulphate of Ammonia, cwt. N per acre	None	0.4	Mean	None
1943	^{±0.336}	^{±0.238}	^{±0.67}	^{±0.47}	^{±0.270}	^{±0.191}	^{±0.70}	
O	1.26	2.94	2.10	15.2	16.6	15.9	1.60	^{±0.99}
D	2.86	3.24	3.05	16.3	17.3	16.8	1.56	42.8
D	3.12	4.04	3.58	17.0	16.1	16.6	2.00	44.4
CTR	2.73	2.88	2.81	17.4	15.9	16.7	1.46	44.6
CTR	2.01	3.16	2.58	13.6	17.4	15.5	0.96	44.2
SS	2.81	3.32	3.07	16.2	17.1	16.6	1.92	44.5
SS	3.61	3.04	3.32	16.7	17.3	17.0	1.50	44.5
VC	3.09	3.27	3.18	17.6	16.0	16.8	2.04	44.5
VC	3.57	3.50	3.54	17.1	17.8	17.4	1.57	44.1
Mean	2.98 ²	3.31 ²	3.00 ³	16.5 ²	16.9 ²	16.6 ³	1.65 ²	43.9 ³
S.E. per plot	0.475 or 15.8%			0.95 or 5.7%			0.382 or 22.4%	1.41 or 3.2%

(2) Excludes "No organic manure"

(3) Mean of all plots.

(4) Organics applied to previous crop of globe beet and peas.

All standard errors have 17 d.f.

Symbols: O No organic manure. SS Sewage sludge.
D Dung. VC Vegetable compost. CTR Composted town refuse.

Winter Cabbage and Leeks

Organic Manures t.p.a. ¹	Winter Cabbage (Series A)				Leeks (Series B)			
	None	0.4	Mean	Sulphate of Ammonia, cwt. N per acre	None	0.4	Mean	Plant number: thous./acre
1944	^{±0.417}	^{±0.295}	^{±0.76}	^{±0.54}	^{±0.479}	^{±0.339}	^{±3.34}	^{±2.36}
O	0.49	1.51	1.00	13.7	16.3	15.0	1.24	27.1
D	1.17	3.41	2.29	16.0	18.6	17.3	1.98	34.8
D	1.53	3.92	2.72	16.9	17.9	17.4	3.44	31.2
CSS	0.68	1.98	1.33	15.2	18.0	16.6	1.50	36.0
CSS	0.78	2.08	1.43	14.0	16.0	15.0	2.53	34.5
SS	2.17	4.09	3.13	14.9	16.8	15.8	1.01	31.9
SS	4.27	4.32	4.29	15.9	17.0	16.4	0.77	29.1
VC	0.91	2.28	1.60	17.0	17.0	17.0	1.67	26.8
VC	1.17	2.88	2.03	14.8	17.0	15.9	3.21	28.0
Mean	1.59 ²	3.12 ²	2.28 ³	15.6 ²	17.3 ²	16.3 ³	2.02 ²	35.6
S.E. per plot	0.589 or 25.8%		1.07 or 6.6%			0.678 or 35.8%	1.89 ³	32.1 ²
							4.73 or 15.2%	31.2 ³

(1) Organics applied to previous crop of globe beet and peas.

(2) Excludes "No organic manure".

(3) Mean of all plots.

All standard errors have 17 d.f.

Symbols: O No organic manure. SS Sewage sludge.
 D Dung VC Vegetable compost.

CSS Composted sewage sludge

Winter Cabbage and Leeks

Organic Manures t.p.a. 1	Winter Cabbage (Series B)				Leeks (Series A)									
	Total produce: tons/acre		Plant number: thous./acre		Total produce: tons/acre		Plant number: thous./acre							
	None	0.4	Mean	Sulphate of Ammonia, cwt. N per acre	None	0.4	Mean	None	0.4	Mean	None	0.4	Mean	
1945														
0	±0.956	+0.676	5.47	±1.00	±0.71	1.17	1.20	±0.151	±2.48	±1.75	1.18	47.6	41.6	
D	4.26	6.67	5.47	16.5	16.6	1.73	1.33	1.53	44.8	44.6	44.8	41.7	43.2	
D	7.82	7.34	7.58	19.2	17.6	1.80	1.26	1.53	42.5	38.4	42.5	38.4	40.4	
CSS	6.97	7.53	7.25	17.5	16.2	1.43	1.09	1.26	43.7	42.6	43.7	42.6	43.2	
CSS	5.85	6.95	6.40	17.6	17.4	17.5	1.25	1.13	1.19	41.1	35.5	41.1	38.3	
CSS	6.77	9.30	8.04	18.2	17.5	17.8	1.25	1.13	1.19	41.1	35.5	41.1	38.3	
SS	6.74	7.54	7.14	16.5	15.8	16.2	1.07	0.88	0.98	38.1	33.9	38.1	36.0	
SS	8.57	8.95	8.76	17.5	17.5	17.5	1.49	1.25	1.37	39.6	35.4	39.6	37.5	
VC	6.01	7.23	6.62	16.9	17.3	17.1	1.39	1.31	1.35	45.4	41.8	45.4	43.6	
VC	7.97	8.97	8.47	17.3	17.6	17.4	1.97	1.47	1.72	43.1	36.8	43.1	36.8	
Mean	7.09 ²	7.98 ²	7.42 ³	17.6 ²	17.1 ²	17.3 ³	1.52 ²	1.21 ²	1.33 ³	42.3 ²	38.3 ²	42.3 ²	38.3 ²	
S.E. per plot	1.352 or 18.4%			1.42 or 8.2%			0.301 or 22.6%			3.50 or 8.6%				

(1) Organics applied to previous crop of globe beet and peas.

(2) Excludes "No organic manure".

(3) Mean of all plots

All standard errors have 17 d.f.

Symbols: O No organic manure. SS Sewage sludge. CSS Composted sewage sludge.
 D Dung VC Vegetable compost.

Winter Cabbage and Leeks

Organic Manures t.p.a.	Winter Cabbage (Series A)			Sulphate of Ammonia, cwt. N per acre			Leeks (Series B)		
	None	0.4	Mean	None	0.4	Mean	None	0.4	Mean
0	1.404	+0.286	+0.32	+0.22	+0.391	+0.276	+1.17	+1.17	+0.83
D	1.45	2.27	1.86	17.5	17.9	17.7	2.10	2.01	42.6
D	2.38	4.00	3.19	17.3	18.1	17.7	2.75	2.74	40.6
D	2.56	4.81	3.69	17.8	17.9	17.8	3.76	3.88	43.6
CSS	1.47	3.72	2.60	17.1	18.0	17.5	2.42	3.06	40.3
CSS	2.06	3.55	2.80	17.3	18.0	17.7	2.27	3.19	42.4
SS	2.62	4.49	3.56	17.5	17.8	17.7	2.82	2.82	42.8
SS	4.47	5.99	5.23	17.5	17.2	17.4	4.09	3.71	39.9
VC	1.17	2.82	1.99	17.8	16.5	17.2	2.61	3.58	42.4
VC	2.25	2.92	2.58	16.8	17.9	17.4	3.42	3.03	42.8
Mean	2.372	4.04 ²	3.10 ³	17.4 ²	17.7 ²	17.6 ³	3.02 ²	3.25 ²	42.2 ²
S.E. per plot	0.572 or 18.5%		0.45 or 2.6%			0.553 or 18.8%	2.94 ³		42.3 ²

(1) Organics applied to previous crop of globe beet and peas.

(2) Excludes "No organic manure".

(3) Mean of all plots.

All standard errors have 17 d.f.

Symbols: O No organic manure. SS Sewage sludge. CSS Composted sewage sludge.
 D Dung VC Vegetable compost.

Winter Cabbage and Leeks

Organic Manures t.p.a. ¹	1947	Sulphate of Ammonia, cwt. N per acre					
		None	0.4	Mean	None	0.4	Mean
		Leeks			Leeks		
		Total produce: tons/acre			Plant number; thous./acre		
O		±0.197	±0.140		±0.30	±0.57	
D 15		1.68	1.82	1.75	42.5	42.6	42.6
D 30		1.80	1.92	1.86	42.2	40.6	41.4
CSS 15		2.24	2.03	2.13	42.9	42.8	42.8
CSS 30	(Cabbage failed)	1.92	1.74	1.83	42.5	42.1	42.3
SS 15		1.97	2.01	1.99	42.4	42.6	42.5
SS 30		1.93	1.79	1.86	42.5	42.6	42.5
VC 15		1.93	1.87	1.90	43.2	43.7	43.5
VC 30		1.81	1.88	1.84	43.5	42.6	43.0
Mean		2.01	2.32	2.16	42.7	44.4	43.6
S.E. per plot		1.95 ²	1.94 ²	1.89 ³	42.7 ²	42.7 ²	42.6 ³
		0.279 or 14.8%			1.14 or 2.7%		

(1) Organics applied to previous crop or globe beet and peas.

(2) Excludes "No organic manure".

(3) Mean of all plots

All standard errors have 17 d.f.

Symbols: O No organic manure. CTR Composted town refuse.
 D Dung. SS Sewage sludge.
 VC Vegetable compost.

Yields on plots without organic manure

	Sulphate of Ammonia, cwt. N per acre									
	None	0.4	0.8	1.2	S.E.	None	0.4	0.8	1.2	S.E.
	Total produce: tons per acre					Plant number: thous. per acre				
1942 ⁴										
	Winter Cabbage									
Series A	5.27	7.68	8.37	5.33	±1.046	17.5	18.2	19.4	17.1	±0.82
Series B	1.30	2.51	2.16	1.70	±0.300	No figures available				
1943	1.26	2.94	3.03	2.57	±0.336	15.2	16.6	16.9	16.1	±0.67
1944	0.49	1.51	3.38	2.62	±0.417	13.7	16.3	17.8	15.7	±0.76
1945	4.26	6.67	6.88	8.33	±0.956	16.5	16.8	16.7	18.4	±1.00
1946	1.45	2.27	2.75	4.23	±0.404	17.5	17.9	17.9	17.6	±0.32
	Leeks									
1943	1.23	1.96	1.90	1.56	±0.270	41.9	43.7	44.2	44.0	±0.99
1944	1.54	0.94	1.17	1.93	±0.479	27.1	29.0	28.1	31.0	±3.34
1945	1.17	1.20	1.21	1.17	±0.213	47.6	41.6	41.8	42.5	±2.48
1946	1.91	2.10	2.01	2.69	±0.391	42.6	40.6	46.4	43.1	±1.17
1947	1.68	1.82	1.85	1.24	±0.197	42.5	42.6	42.0	41.8	±0.80

(4) The dressings of nitrogen in 1942 were None, 0.6, 1.2, 1.8 cwt. per acre, since no globe beet or peas were grown.

All standard errors have 17 d.f.

CHEMICAL ANALYSES OF FERTILIZERS

Three, Four and Six Course Rotations, 1939-1947

Crop, Year	% N Nitrate of soda	% Total P ₂ O ₅ Super-phosphate	% Total P ₂ O ₅ Mineral ²⁵	% K ₂ O Sulphate of potash ³	% K ₂ O Muriate of potash
1939 a		17.0	25.9		53.2
b	16.0	16.9		49.5	52.2
1940 a		16.8	25.9		52.2
b	15.7	17.9		50.3	53.9
1941 a		18.9	25.7		57.0
b	15.7	18.0		49.8	53.8
1942 a		20.0	25.7		53.8
b	16.1	20.3		49.9	53.7
1943 a		20.3	26.4		53.7
b	15.9	20.4		49.8	57.4
1944 a		19.6 ⁴	27.0		57.7
b	16.1	19.6		49.8	57.4
1945 a		19.2	33.2		57.8
b	16.2	18.8		51.2	57.1
1946 a		19.0	33.3		60.6
b	15.8	19.6		50.7	59.9
1947	15.6	19.9	33.1		60.2

Analysis of Sulphate of ammonia 21% N assumed.

1. b refers to spring dressings in the 3-Course Rotation Experiment; a to all other dressings in the 3-, 4- and 6-Course Rotation Experiments. "Crop year 1939" includes 1938 autumn sowings.
2. Mineral phosphate (used in 4-Course only); Gafsa 1939-44, Moroccan 1945-47.
3. Sulphate of potash, applied in spring to potatoes in 3-Course until 1946, replaced by Muriate of potash in 1947.
4. 18.86 for Superphosphate in 4-Course.

Chemical Analysis of Manures used in Three and Four Course Rotations.1939-47Wheat Straw to Three and Four Course Rotations

Crop Year	% Organic Matter	% N	% P ₂ O ₅	% K ₂ O
1939	83.65	0.603	0.311	1.614
1940	77.00	0.541	0.244	1.600
1940 b	84.30	0.537	0.194	1.382
1941	82.87	0.445	0.142	1.185
1942	79.08	0.409	0.138	0.817
1943	84.02	0.525	0.170	0.950
1944	81.00	0.532	0.144	0.620
1945	82.40	0.299	0.088	0.688
1946	81.41	0.450	0.250	0.954
1947	80.38	0.555	0.171	1.011


Adco Compost to Three and Four Course Rotations


1939	12.74	0.398	0.316	0.339
1940	13.73	0.404	0.235	0.332
1941	15.79	0.398	0.277	0.170
1942	14.44	0.398	0.257	0.146
1943	19.00	0.487	0.319	0.180
1944	20.10	0.528	0.284	0.230
1945	21.80	0.434	0.242	0.271
1946	18.04	0.527	0.372	0.322
1947	15.90	0.504	0.372	0.290

Dung to Four Course Rotation

1939	16.45	0.500	0.273	0.709
1940	17.62	0.435	0.164	0.414
1941	14.65	0.520	0.138	0.528
1942	18.08	0.510	0.253	0.858
1943	23.26	0.710	0.300	1.020
1944	35.20	0.846	0.288	1.355
1945	29.65	0.904	0.292	1.608
1946	14.29	0.515	0.203	0.691
1947	14.26	0.453	0.213	0.445

[#]'b' refers to 3-Course Spring dressings





Rothamsted Field Experiments, 1939-47, Vol. I

CORRIGENDA

Misleading faults caused by defective correcting material used in the reproduction are listed.

Page	
A/1.6	1943 Plot 3 Grain Mean, 14.3; 1944 Plot 18 Grain Mean, 12.5; 1944 Plot 20 Grain Mean, 17.3.
A/1.7	1945 Plot 17 Grain Mean, 11.0; straw figures are 29.9, 15.9, 12.4 15.1, 18.3
A/2.3	1940 Plot 7-2, 35.6
A/5.4	1945 Plot 8 Roots, No Cross Dressing, 3.19
A/6.4	1939 Plot 1 Limed 2nd Crop, 11.8
A/6.6	1943 Plot 11-1 Limed 2nd crop, 9.8; Plot 12 Limed Total, nil.
A/6.9	1949 Plot 4 ¹ Unlimed, 2.0
A/6.10	1941 heading L (only), 1939 Sp. 105 Unlimed 0.1
Ba/1.3	1942 Standard error (a), 0.517.
Ba/1.7	The four "In seed bed" rows immediately above the standard errors should be deleted.
Ba/3.2	Wheat Variety 1939-45 Yeoman 1946,47 Squareheads Master
Ba/4.1	Area of each plot at Woburn, 0.27 acre.
Ba/4.2	1947 Clover Sown 15.4.46
Ba/4.3	1939 Sugar % Woburn N increment, -0.05
Bc/1.10	1946 Dung response, Deep ploughing, -0.10
Bc/1.12	1944 Dung response, Phosphate absent, -1.4; 1945 Dung response, Deep Ploughing, -0.9 (delete "Dung Abs" figure).
Bc/1.13	1945 Delete 2nd decimal places without rounding-off.
Bc/1.14	1945 Dung response, Potash absent, 2.4 (delete "Dung Pres." figure)
Bc/1.16	1946 Delete all figures on same line as "Ploughing" except the standard errors ±1.77 and ±2.51.
Bc/1.20	1947 All potash responses should be positive.
Bc/1.21	1944 Shallow ploughing, Potash in seed bed, 77.4
Be/1.1	Sulphate of ammonia treatment rates, 0.4, 0.8 cwt. N per acre.
Be/1.6	1947 Standard error of mean (bottom line) ±0.54
Be/1.7	1947 Cabbages weight, response to Sulph. Amm. to Green crops in absence of Straw, +0.20; 1946 Plant Number, responses to Sulph. Amm. to Green Crops in abs. and pres. of straw, -1.5 and +0.5.
Be/1.10	Insert headings "Tares" and "Mustard".
Bf/1.3	Lucerne Block 3 1st Cycle Third year, No third cut.
Bf/1.4	1938 Sown Mar. 9. Harvested Aug. 25.
Bf/1.6	Block 2 First year Total increase, -0.02
Bf/1.17	1942 Barley Straw S.E. per sub-plot, 3.31